

3-2017

# Pion Distribution Amplitude and Quasidistributions

A. V. Radyushkin

Old Dominion University, aradyush@odu.edu

Follow this and additional works at: [http://digitalcommons.odu.edu/physics\\_fac\\_pubs](http://digitalcommons.odu.edu/physics_fac_pubs)

 Part of the [Astrophysics and Astronomy Commons](#), [Elementary Particles and Fields and String Theory Commons](#), and the [Quantum Physics Commons](#)

---

## Repository Citation

Radyushkin, A. V., "Pion Distribution Amplitude and Quasidistributions" (2017). *Physics Faculty Publications*. 62.  
[http://digitalcommons.odu.edu/physics\\_fac\\_pubs/62](http://digitalcommons.odu.edu/physics_fac_pubs/62)

## Original Publication Citation

Radyushkin, A. V. (2017). Pion distribution amplitude and quasidistributions. *Physical Review D*, 95(5), 056020. doi:10.1103/PhysRevD.95.056020

**Pion distribution amplitude and quasidistributions**

A. V. Radyushkin

*Physics Department, Old Dominion University, Norfolk, Virginia 23529, USA*  
*Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA*  
 (Received 17 January 2017; published 27 March 2017)

We extend our analysis of quasidistributions onto the pion distribution amplitude. Using the formalism of parton virtuality distribution amplitudes, we establish a connection between the pion transverse momentum dependent distribution amplitude  $\Psi(x, k_\perp^2)$  and the pion quasidistribution amplitude (QDA)  $\mathcal{Q}_\pi(y, p_3)$ . We build models for the QDAs from the virtuality-distribution-amplitude-based models for soft transverse momentum dependent distribution amplitudes, and analyze the  $p_3$  dependence of the resulting QDAs. As there are many models claimed to describe the primordial shape of the pion distribution amplitude, we present the  $p_3$ -evolution patterns for models producing some popular proposals: Chernyak-Zhitnitsky, flat, and asymptotic distribution amplitude. Our results may be used as a guide for future studies of the pion distribution amplitude on the lattice using the quasidistribution approach.

DOI: 10.1103/PhysRevD.95.056020

**I. INTRODUCTION**

The parton distribution functions (PDFs)  $f(x)$ , and two-body distribution amplitudes (DAs)  $\varphi(x)$  are related to matrix elements of bilocal operators on the light cone  $z^2 = 0$ , which prevents a straightforward calculation of these functions in a lattice gauge theory formulated in the Euclidean space. The usual way out is to calculate their moments. In particular, high precision lattice calculations of the second moment of the pion distribution amplitude  $\varphi_\pi(x)$  were reported in Ref. [1]. However, recently, X. Ji [2] suggested a method allowing us to calculate PDFs and DAs as functions of  $x$ . To this end, he proposes to use purely spacelike separations  $z = (0, 0, 0, z_3)$ .

The matrix elements of equal-time bilocal operators produce distributions  $Q(y, p_3)$  in the momentum  $p_3$  component (quasidistributions). The crucial point is that they tend to the light-cone distributions  $f(y)$ ,  $\varphi(y)$  in the  $p_3 \rightarrow \infty$  limit. In case of PDFs, the results of lattice calculations of the parton quasidistributions (PQDs) were reported in Refs. [3–8]. It is expected [9] that PQDs  $Q(y, p_3)$  should have a mild perturbative evolution [10–13] with respect to  $p_3$  for large  $p_3$ . However, the values of  $p_3$  used in the cited lattice calculations are not very large, and the observed strong variation of PQDs with  $p_3$  does not have a perturbative form.

In our recent paper [14] we have studied nonperturbative evolution of PQDs using the formalism of *virtuality distribution functions* [15,16]. We found that PQDs can be obtained from the transverse momentum dependent distributions (TMDs)  $\mathcal{F}(x, k_\perp^2)$ . Then we built models for the nonperturbative evolution of PQDs using simple models for TMDs. Our results are in qualitative agreement with the  $p_3$ -evolution patterns obtained in lattice calculations [3–8] and also in diquark spectator models [17–19].

As emphasized in Ref. [14], because of the relation between PQDs and TMDs, the nonperturbative evolution of PQDs reflects the  $k_\perp$  dependence of the TMDs  $\mathcal{F}(x, k_\perp^2)$ , and thus its study provides a new approach to the investigation of the three-dimensional structure of hadrons.

Our goal in the present paper is to perform a similar analysis of the pion *quasidistribution amplitude* (QDA)  $\mathcal{Q}_\pi(y, p_3)$  that produces the pion DA  $\varphi_\pi(y)$  in the large- $p_3$  limit. The basic ingredients of our analysis are *virtual distribution amplitudes* (VDAs) and *transverse momentum dependent amplitudes* (TMDAs) introduced in Refs. [15,16].

The paper is organized as follows. We start in Sec. II with an introductory overview of the basic concepts involved. First, we recall a covariant definition of the twist-2 pion distribution amplitude. After that, we discuss its definition within the light-front (LF) formalism. Then we outline the basics of the VDA/TMDA approach. In Sec. III, we discuss the quasidistribution amplitudes. In particular, we show that QDAs are completely determined by TMDAs through a rather simple transformation. Since the basic relations between the parton distributions are rather insensitive to complications brought by spin, in Sec. III we refer to a simple scalar model. In Sec. IV, we discuss modifications related to quark spin and the gauge nature of gluons in quantum chromodynamics (QCD). In Sec. V we discuss the VDA-based models for soft TMDAs, and present our results for nonperturbative evolution of QDAs obtained in these models. The large- $p_3$  limit of perturbative evolution is discussed in Sec. VI. Our conclusions are given in Sec. VII.

**II. PION DISTRIBUTION AMPLITUDE****A. Covariant definition**

The pion DA  $\varphi_\pi(x, \mu^2)$  was originally introduced [20] as a function  $\varphi_\pi(x, \mu^2)$  whose  $x^n$  moments

$$f_n(\mu^2) = \int_0^1 x^n \varphi_\pi(x, \mu^2) dx \quad (1)$$

are given by reduced matrix elements of twist-2 local operators

$$\begin{aligned} & i^{n+1} R_{\mu^2} \langle 0 | \bar{d}(0) \gamma_5 \{ \gamma_\nu D_{\nu_1} \dots D_{\nu_n} \} u(0) | \pi^+, P \rangle \\ & = \{ P_\nu P_{\nu_1} \dots P_{\nu_n} \} f_n(\mu^2). \end{aligned} \quad (2)$$

As usual  $\{\dots\}$  denotes the twist-2 projection of a Lorentz structure, i.e., symmetrization of indices and subtraction of traces. Since matrix elements of local operators with  $n > 0$  diverge, one needs to supply them by a renormalization procedure denoted above by  $R_{\mu^2}$ , with  $\mu^2$  being the renormalization scale. In QCD, the standard choice of  $R_{\mu^2}$  is based on the dimensional regularization and the modified minimal subtraction scheme  $\overline{\text{MS}}$ . As a result of such a renormalization, the zeroth moment  $f_0(\mu^2)$  does not have  $\mu^2$  dependence since the anomalous dimension of the axial current is zero. Hence,  $f_0(\mu^2)$  for all  $\mu^2$  is equal to the pion decay constant  $f_\pi$

$$f_0(\mu^2) = \int_0^1 \varphi_\pi(x, \mu^2) dx = f_\pi \quad (3)$$

known experimentally,  $f_\pi \approx 130$  MeV.

This definition of DA is oriented on the use of the operator product expansion and a description of the pion in terms of the twist-2 DA  $\varphi_\pi(x, \mu^2)$  that gives the collinear distribution of the pion momentum  $p$  among its two valence constituents. The dependence of  $\varphi_\pi(x, \mu^2)$  on  $\mu^2$  is governed by perturbative evolution [21–23] and does not reflect the primordial (nonperturbative) pion's structure in the direction transverse to  $p$ .

As is well known, for very large  $\mu^2$ , the pion DA tends to the ‘‘asymptotic DA’’  $\varphi_\pi^{\text{as}}(x) = 6f_\pi x(1-x)$  [24]. In general,  $\varphi_\pi(x, \mu^2)$  may differ from its asymptotic form. Over the years, several forms were proposed for the pion DA ‘‘at low normalization point,’’ e.g., Chernyak-Zhitnitsky DA  $\varphi_\pi^{\text{CZ}}(x) = 30f_\pi x(1-x)(1-2x)^2$  [25], ‘‘flat DA’’  $\varphi_\pi^{\text{flat}}(x) = f_\pi$  [26–30], ‘‘root DA’’  $\varphi_\pi^{\text{root}}(x) = 8f_\pi \sqrt{x(1-x)}/\pi$  [31], etc.

## B. Light-front formalism definition

A different definition [23] is used in the LF quantization framework, where the pion distribution amplitude  $\phi_\pi(x, \mu^2)$  is understood as the  $k_\perp$  integral

$$\phi_\pi(x, \mu^2) = \frac{\sqrt{6}}{(2\pi)^3} \int_{k_\perp^2 \leq \mu^2} \psi(x, k_\perp) d^2 k_\perp \quad (4)$$

of the light-front wave function (LFWF)  $\psi(x, k_\perp)$ . We intentionally use here a different notation  $\phi_\pi(x, \mu^2)$  to

emphasize the fact that  $\psi(x, k_\perp)$  is an object of the *Hamiltonian light-front framework*, while the pion DA  $\varphi_\pi(x, \mu^2)$  in Eq. (1) is defined within the *covariant Lagrangian formulation* of the quantum field theory (QFT).

Another difference is the use of a straightforward cutoff  $k_\perp^2 \leq \mu^2$  rather than a more sophisticated  $\overline{\text{MS}}$ -like subtraction. As a result,  $\phi_\pi(x, \mu^2)$  has a nonperturbative evolution with  $\mu^2$  even if the perturbative evolution is absent. Take a simple example  $\psi(x, k_\perp) \sim \phi(x) e^{-k_\perp^2/\Lambda^2}$ . Then the zeroth  $x$  moment of  $\phi_\pi(x, \mu^2)$  has the  $\sim [1 - e^{-\mu^2/\Lambda^2}]$  dependence, i.e. it is not constant, reaching  $f_\pi$  in the  $\mu^2 \rightarrow \infty$  limit only.

Of course, if one has in mind only the applications in which nonperturbative part of the  $\mu^2$  dependence may be ignored, then  $\phi_\pi(x, \mu^2)$  of the LF definition is very similar to the covariantly defined  $\varphi_\pi(x, \mu^2)$ , and the difference between them may be treated as the use of different renormalization schemes.

As a matter of fact, in actual LF calculations one encounters LFWFs integrated to some process-dependent scale  $\mu$ , i.e., the choice of the renormalization prescription and the scale  $\mu$  is dictated by diagrams. Moreover, if the relevant  $\mu^2$ 's are not extremely large, the simple example above shows that one may need to take into account the nonperturbative  $\mu^2$  dependence of  $\phi_\pi(x, \mu^2)$  reflecting the transverse momentum behavior of the LFWF  $\psi(x, k_\perp)$ , i.e., the three-dimensional structure of the pion, which may be essential for some processes.

In particular, the photon-pion transition form factor involves  $\phi_\pi(x, \mu^2 = x^2 Q^2)/[xQ^2]$ , i.e., LFWF  $\psi(x, k_\perp)$  integrated over  $k_\perp$  until  $xQ$  [32,33]. As a result, the remaining  $x$  integral in the LF formula has a finite  $Q^2 \rightarrow 0$  limit: the infrared small- $x$  divergence is eliminated by a cutoff provided by  $\phi(x, \mu^2 = x^2 Q^2)$ . On the other hand, a formula involving  $\overline{\text{MS}}$ -based DA  $\varphi(x, \mu^2)$  with a fixed scale  $\mu^2$  is singular in the  $Q^2 \rightarrow 0$  limit. One may question the applicability of the LF formula down to  $Q^2 = 0$ , but at least it does not give an infinite result for a quantity that is known to be finite. For this reason, the LF formula looks as a more attractive tool for modeling the form factor behavior at moderate  $Q^2$  than the perturbative QCD  $1/Q^2$  twist expansion.

Still, a problem with the LF formalism is that LFWFs are not directly connected with the usual objects of the covariant field theory, such as matrix elements of local or nonlocal operators.

In our papers [15,16], we have developed the formalism of VDAs that is fully based on the covariant field theory concepts. In the VDA approach, the pion is described by the TMDA which has a direct connection with the objects of the covariant QFT. On the other hand, just like the LF wave functions, the TMDAs give a three-dimensional description of the pion structure.

### C. Pion TMDA

To omit inessential complications related to spin, we illustrate the ideas underlying TMDAs using a simple example of a scalar theory. The key element of our approach [15] is the *VDA representation*

$$\langle 0|\psi(0)\psi(z)|p\rangle = \int_0^\infty d\sigma \int_0^1 dx \times \Phi(x, \sigma) e^{-ix(pz) - i\sigma(z^2 - i\epsilon)/4} \quad (5)$$

that basically reflects the fact that the matrix element  $\langle 0|\psi(0)\psi(z)|p\rangle$  depends on  $z$  through  $(pz)$  and  $z^2$ . It may be treated as a double Fourier representation with respect to these variables.

The main nontrivial feature of this representation is in its specific limits of integration over  $x$  and  $\sigma$ . They hold for any contributing Feynman diagram [16], so we assume that this property is true in general. Note that starting with the first loop, the diagram contributions are nonanalytic in  $z^2$  due to  $\ln z^2$  factors, but the VDA representation, unlike the Taylor expansion in  $z^2$ , is valid nevertheless.

While the VDA representation is a fully covariant expression, it is convenient to use a frame in which the pion momentum  $p$  is purely longitudinal  $p = (E, \mathbf{0}_\perp, P)$ . Choosing some special cases of  $z$ , one can get representations for several parton functions, all in terms of one and the same universal VDA  $\Phi(x, \sigma)$ . In particular, choosing  $z$  on the light front  $z_+ = 0$  and with  $z_\perp = 0$  (i.e., taking  $z = z_-$ ) gives the twist-2 distribution amplitude  $\varphi(x)$

$$\langle 0|\psi(0)\psi(z_-)|p\rangle = \int_0^1 dx \varphi(x) e^{-ixp_+ z_-}. \quad (6)$$

Comparing this relation with the VDA representation we have

$$\varphi(x) = \int_0^\infty \Phi(x, \sigma) d\sigma, \quad (7)$$

provided that the  $z^2 \rightarrow 0$  limit is finite, e.g., in the super-renormalizable  $\varphi^3$  theory. In the renormalizable  $\varphi^4$  theory, the function  $\Phi(x, \sigma)$  has a  $\sim 1/\sigma$  hard part, and the integral (7) is logarithmically divergent, reflecting the perturbative evolution of the DA in such a theory. In this case, one may arrange a regularization of the  $\sigma$  integral characterized by some parameter  $\mu^2$ . Then  $\varphi(x) \rightarrow \varphi(x, \mu^2)$ .

Light-cone singularities are avoided if we choose a spacelike  $z$ , e.g., take  $z$  that has  $z_-$  and  $z_\perp$  components only. Then we can introduce the transverse momentum dependent distribution amplitude  $\Psi(x, k_\perp^2)$  as a Fourier transform

$$\langle 0|\psi(0)\psi(z_-, z_\perp)|p\rangle = \int_0^1 dx e^{-ixp_+ z_-} \times \int d^2 k_\perp \Psi(x, k_\perp^2) e^{i(k_\perp z_\perp)} \quad (8)$$

of the matrix element with respect to  $z_-$  and  $z_\perp$ . Because of the rotational invariance in the  $z_\perp$  plane, TMDA depends on  $k_\perp^2$  only, the fact already reflected in the notation. The TMDA may be written in terms of the VDA as

$$\Psi(x, k_\perp^2) = \frac{i}{\pi} \int_0^\infty \frac{d\sigma}{\sigma} \Phi(x, \sigma) e^{-i(k_\perp^2 - i\epsilon)/\sigma}. \quad (9)$$

The integrated TMDA

$$\mathfrak{f}(x, \mu^2) \equiv \pi \int_0^{\mu^2} dk_\perp^2 \Psi(x, k_\perp^2) \quad (10)$$

is analogous to the  $\mu^2$ -dependent pion distribution amplitude  $\phi(x, \mu^2)$  of the LF formalism [but, of course, being an object of the covariant QFT,  $\mathfrak{f}(x, \mu^2)$  does not coincide with it]. In terms of the VDA,

$$\mathfrak{f}(x, \mu^2) = \int_0^\infty d\sigma [1 - e^{-i(\mu^2 - i\epsilon)/\sigma}] \Phi(x, \sigma). \quad (11)$$

Since it is defined by a straightforward cutoff,  $\mathfrak{f}(x, \mu^2)$  evolves with  $\mu^2$  even if the limit  $\mu^2 \rightarrow \infty$  is finite, e.g., in a super-renormalizable theory. The evolution equation

$$\mu^2 \frac{d}{d\mu^2} \mathfrak{f}(x, \mu^2) = \pi \mu^2 \Psi(x, \mu^2) \quad (12)$$

follows from the definition (10). When the TMDA  $\Psi(x, k_\perp^2)$  vanishes faster than  $1/k_\perp^2$  (such a TMDA will be called “soft”), evolution essentially stops at large  $\mu^2$ .

In a renormalizable theory, it makes sense to treat  $\Phi(x, \sigma)$  as a sum of a soft part  $\Phi^{\text{soft}}(x, \sigma)$ , generating a nonperturbative evolution of  $\mathfrak{f}(x, \mu^2)$ , and a  $\sim 1/\sigma$  hard tail. To avoid nonperturbative evolution, one may choose an  $\overline{\text{MS}}$ -type construction, e.g., regularize the  $\sigma$  integral in Eq. (7) by a  $\sigma^{-\epsilon}$  factor and then subtract  $1/\epsilon$  poles.

However, just like in the LF formalism, the objects that appear in actual calculations are exactly the integrated TMDAs rather than their  $\overline{\text{MS}}$ -type sisters. In particular, the photon-pion transition form factor is given in the VDA approach by the  $x$  integral of  $\mathfrak{f}(x, \mu^2)/[xQ^2]$  taken at  $\mu^2 = xQ^2$  [15], i.e., it involves TMDA  $\Psi(x, k_\perp^2)$  integrated over  $k_\perp^2$  until  $xQ^2$ . As a result, the TMDA formula has a finite  $Q^2 \rightarrow 0$  limit. Furthermore, using simple models for soft TMDAs one can get a very close description of experimental data by the nonperturbative evolution of the integrated TMDA [16].

For very large  $\mu^2$ , the perturbative evolution dominates and eventually brings  $\mathfrak{f}(x, \mu^2)$  to its asymptotic form  $6f_\pi x(1-x)$ . The question, however, is what kind of shape  $\mathfrak{f}(x, \mu^2)$  has at low scales  $\mu \sim 1$  GeV, and also how this shape changes with  $\mu^2$ . As we have discussed, this non-perturbative  $\mu^2$  evolution reflects the  $k_\perp$  dependence of the soft part of the pion TMDA.

Below, we shall see that there is another function, the pion *quasidistribution amplitude*  $Q_\pi(y, P)$  whose  $P$  dependence is also determined by the  $k_\perp$  dependence of the pion TMDA. The quasidistributions have been introduced recently by X. Ji [2] to facilitate a calculation of light-front functions (PDFs, DAs, etc.) on the lattice.

### III. QUASIDISTRIBUTION AMPLITUDE

#### A. Definition

The basic proposal of Ref. [2] is to consider equal-time bilocal operators corresponding to  $z = (0, 0, 0, z_3)$  (or, for brevity,  $z = z_3$ ). Incorporating the VDA representation, we have

$$\langle 0 | \psi(0) \psi(z_3) | p \rangle = \int_0^\infty d\sigma \int_{-1}^1 dx \Phi(x, \sigma) e^{ixp_3 z_3 + i\sigma z_3^2/4}. \quad (13)$$

Using again the frame in which  $p = (E, 0_\perp, P)$ , and introducing the pion quasidistribution amplitude through

$$\langle 0 | \psi(0) \psi(z_3) | p \rangle = \int_{-\infty}^\infty dy Q_\pi(y, P) e^{-iyPz_3}, \quad (14)$$

we get a relation between QDA and VDA,

$$Q_\pi(y, P) = \sqrt{\frac{iP^2}{\pi}} \int_0^\infty \frac{d\sigma}{\sqrt{\sigma}} \int_0^1 dx \Phi(x, \sigma) e^{-i(x-y)^2 P^2/\sigma}. \quad (15)$$

It is easy to see that, for large  $P$ , we have

$$\sqrt{\frac{iP^2}{\pi\sigma}} e^{-i(x-y)^2 P^2/\sigma} = \delta(x-y) + \frac{\sigma}{4P^2} \delta''(x-y) + \dots \quad (16)$$

and  $Q_\pi(y, P \rightarrow \infty)$  tends to the integral (7) leading to  $\varphi_\pi(y)$ . This observation suggests that one may be able to extract the ‘‘light-cone’’ distribution amplitude  $\varphi_\pi(y)$  from the studies of the purely ‘‘spacelike’’ function  $Q_\pi(y, P)$  for large  $P$  [2].

#### B. Evolution

Again, to study the  $P$  evolution of  $Q_\pi(y, P)$  it makes sense to split  $\Phi(x, \sigma)$  into the soft part, for which the integral over  $\sigma$  is finite, and the hard tail that generates perturbative evolution.

The nonperturbative evolution of  $Q_\pi^{\text{soft}}(y, P)$  with respect to  $P$  has the area-preserving property. Namely, since

$$\int_{-\infty}^\infty dy e^{-i(x-y)^2 P^2/\sigma} = \sqrt{\frac{\pi\sigma}{iP^2}} \quad (17)$$

we have

$$\int_{-\infty}^\infty dy Q_\pi^{\text{soft}}(y, P) = \int_0^1 dx \varphi_\pi^{\text{soft}}(x) = f_\pi. \quad (18)$$

In other words,  $Q_\pi^{\text{soft}}(y, P)$  for any  $P$  has the same area normalization as  $\varphi_\pi^{\text{soft}}(x)$ . In this respect, the pion QDA pleasantly differs from the integrated TMDA  $\mathfrak{f}^{\text{soft}}(x, \mu^2)$  whose zeroth moment is  $\mu^2$  dependent.

Similarly, we have the momentum sum rule

$$\int_{-\infty}^\infty dy y Q_\pi^{\text{soft}}(y, P) = \int_0^1 dx x \varphi_\pi^{\text{soft}}(x). \quad (19)$$

#### C. Relation to TMDA

Comparing the VDA representation (16) for  $Q_\pi(y, P)$  with that for the TMDA  $\Psi(x, k_\perp^2)$  (9) (note that they are valid both for soft and hard parts) we conclude that

$$Q_\pi(y, P) = \int_{-\infty}^\infty dk_1 \int_0^1 dx P \Psi(x, k_1^2 + (x-y)^2 P^2). \quad (20)$$

Thus, the quasidistribution amplitude  $Q_\pi(y, P)$  is completely determined by the form of the TMDA  $\Psi(x, k_\perp^2)$ .

This formula may be also obtained if one takes  $z = (0, z_1, 0, z_3)$  in the VDA representation and introduces the momentum  $k_1$  conjugate to  $z_1$ . Then

$$\begin{aligned} & \int_{-\infty}^\infty dy e^{iyPz_3} \langle 0 | \psi(0) \psi(z_1, z_3) | p \rangle \\ &= \int_{-\infty}^\infty dk_1 e^{-ik_1 z_1} \int_0^1 dx \Psi(x, k_1^2 + (x-y)^2 P^2). \end{aligned} \quad (21)$$

Taking  $z_1 = 0$  gives Eq. (20). Furthermore, introducing the variable  $k_3 \equiv (x-y)P$ , we have

$$Q_\pi(y, P) = \int_{-\infty}^\infty dk_1 \int_{-yP}^{(1-y)P} dk_3 \Psi(y + k_3/P, k_1^2 + k_3^2). \quad (22)$$

Thus,  $Q_\pi(y, P)$  is given by an integral over a stripe of width  $P$  in the two-dimensional  $(k_1, k_3)$  plane. When  $P \rightarrow \infty$  for a fixed nonzero  $y$ , the stripe covers the whole  $(k_1, k_3)$  plane. Moreover, for a soft TMDA  $\Psi(x, k^2)$  that rapidly decreases outside a region  $k^2 \lesssim \Lambda^2$ , only the values of  $k_3 \lesssim \Lambda$  are essential, and for large  $P$  one may

approximate the first argument of the TMDA by  $y$ . Hence, the  $P \rightarrow \infty$  limit gives  $\varphi_\pi^{\text{soft}}(y)$ .

For comparison, the integrated TMDA  $\mathfrak{f}(y, \mu^2)$  is obtained by integrating  $\Psi(y, k_\perp^2)$  over a circle of radius  $\mu$  in the  $k_\perp$  plane. Again, the circle covers the whole plane when  $\mu \rightarrow \infty$ , and  $\mathfrak{f}^{\text{soft}}(y, \mu^2) \rightarrow \varphi_\pi^{\text{soft}}(y)$ .

Thus, while the patterns of the nonperturbative evolution of  $Q_\pi^{\text{soft}}(y, P)$  and  $\mathfrak{f}^{\text{soft}}(y, \mu^2)$  are different, they become more and more close for large  $P$  and  $\mu$ , eventually producing the same function  $\varphi_\pi^{\text{soft}}(y)$ .

## IV. QCD

### A. Spinor quarks

In the spinor case, one deals with the matrix element

$$B^\alpha(z, p) \equiv \langle 0 | \bar{\psi}(0) \gamma_5 \gamma^\alpha \psi(z) | p \rangle. \quad (23)$$

It may be decomposed into  $p^\alpha$  and  $z^\alpha$  parts:  $B^\alpha(z, p) = p^\alpha B_p(z, p) + z^\alpha B_z(z, p)$ , or in the VDA representation

$$B^\alpha(z, p) = \int_0^\infty d\sigma \int_{-1}^1 dx \times [p^\alpha \Phi(x, \sigma) + z^\alpha Z(x, \sigma)] e^{-ix(pz) - i\sigma(z^2 - ie)/4}. \quad (24)$$

If we take  $z = (z_-, z_\perp)$  in the  $\alpha = +$  component of  $\mathcal{O}^\alpha$ , the purely higher-twist  $z^\alpha$  part drops out and we can introduce the TMDA  $\Psi(x, k_\perp^2)$  that is related to the VDA  $\Phi(x, \sigma)$  by the scalar formula (9).

In the QDA case, the easiest way to avoid the effects of the  $z^\alpha$  admixture is to take the time component of  $B^\alpha(z = z_3, p)$  and define

$$B^0(z_3, p) = p^0 \int_{-1}^1 dx Q_\pi(y, P) e^{iyPz_3}. \quad (25)$$

The connection between  $Q_\pi(y, P)$  and  $\Phi(x, \sigma)$  is given then by the same formula (15) as in the scalar case. As a result, we have the sum rules (18) and (19) corresponding to charge and momentum conservation. Furthermore, the quasidistribution amplitude  $Q_\pi(y, P)$  is related to TMDA  $\Psi(x, k_\perp^2)$  by the scalar conversion formula (20).

### B. Gauge fields

In QCD, for  $\pi^+$  one should take the operator

$$\mathcal{O}^+(0, z; A) \equiv \bar{d}(0) \gamma_5 \gamma^\alpha \hat{E}(0, z; A) u(z) \quad (26)$$

involving a straight-line path-ordered exponential

$$\hat{E}(0, z; A) \equiv P \exp \left[ igz_\nu \int_0^1 dt A^\nu(tz) \right] \quad (27)$$

in the quark (adjoint) representation. As is well known, its Taylor expansion has the same structure as that for the original  $\bar{\psi}(0) \gamma_5 \gamma^\alpha \psi(z)$  operator, with the only change that one should use covariant derivatives  $D^\nu = \partial^\nu - igA^\nu$  instead of the ordinary  $\partial^\nu$  ones.

Again, the  $z^\alpha$  admixture is avoided if the pion quasidistribution amplitude is defined through the time component of  $\mathcal{O}^\alpha$ . Then we have the same relation between the VDA and QDA as in the scalar case. Because of Eq. (18), this results in the area-preserving property for the soft part

$$\int_{-\infty}^\infty dy Q^{\text{soft}}(y, P) = f_\pi. \quad (28)$$

Also, due to Eq. (19) we have the momentum sum rule

$$\int_{-\infty}^\infty dy (y - 1/2) Q_q^{\text{soft}}(y, P) = 0. \quad (29)$$

Since the VDA  $\Phi(x, \sigma)$  is defined through the matrix element of a gauge-invariant operator, it is gauge invariant also. For this reason, TMDA  $\Psi(x, k_\perp^2)$  is a gauge-invariant object as well. It should not be confused with the  $k_T$ -dependent (and gauge-dependent) ‘‘underintegrated distributions’’ that appear in perturbative loop calculations based on the Sudakov decomposition of the integration momentum  $k$ .

## V. MODELS FOR SOFT PART

### A. Models

To get an idea about patterns of the nonperturbative evolution of the QDAs, we need some explicit models of the  $k_\perp$  dependence of soft TMDAs  $\Psi(x, k_\perp^2)$ . We will use here the same models as in our papers [14, 16]. While TMDAs are functions of two independent variables  $x$  and  $k_\perp^2$ , we take, for simplicity, the case of factorized models

$$\Psi(x, k_\perp^2) = \varphi_\pi(x) \psi(k_\perp^2), \quad (30)$$

in which  $x$  dependence and  $k_\perp$  dependence appear in separate factors.

If we assume a Gaussian dependence on  $k_\perp$ ,

$$\Psi^G(x, k_\perp^2) = \frac{\varphi_\pi(x)}{\pi \Lambda^2} e^{-k_\perp^2/\Lambda^2}, \quad (31)$$

the conversion formula (20) results in

$$Q_\pi^G(y, P) = \frac{P}{\Lambda \sqrt{\pi}} \int_0^1 dx \varphi_\pi(x) e^{-(x-y)^2 P^2/\Lambda^2}. \quad (32)$$

In the space of impact parameters  $z_\perp$ , the Gaussian model gives a  $e^{-z_\perp^2 \Lambda^2/4}$  falloff that is too fast for large  $z_\perp$ . As an alternative extreme case, we take a model with the  $1/(1 + z_\perp^2 \Lambda^2/4)$  dependence on  $z_\perp$ , whose falloff at large

$z_{\perp}$  is too slow. It corresponds to the “slow” model for the TMDA

$$\Psi^S(x, k_{\perp}^2) = 2\varphi_{\pi}(x) \frac{K_0(2|k_{\perp}|/\Lambda)}{\pi\Lambda^2} \quad (33)$$

that has a logarithmic singularity for small  $k_{\perp}$  reflecting a too slow falloff for large  $z_{\perp}$ . For the QDA, we have

$$Q_{\pi}^S(y, P) = \frac{P}{\Lambda} \int_0^1 dx \varphi_{\pi}(x) e^{-2|x-y|P/\Lambda}. \quad (34)$$

Note that the Gaussian model and the slow model have the same  $\sim(1 - z_{\perp}^2\Lambda^2/4)$  behavior for small  $z_{\perp}$ , i.e., they correspond to the same value of the  $\langle 0|\varphi(0)\partial^2\varphi(0)|P\rangle$  matrix element (in the scalar case), provided that one takes the same value of  $\Lambda$  in both models. For large  $z_{\perp}$ , however, the Gaussian model has a falloff that is too fast, while the falloff of the slow model is too slow. Thus, they look like two extreme cases, and provide a good illustration of the nonperturbative evolution of the pion QDA, with expectation that other models would produce results somewhere in between these two cases.

## B. Numerical results

To compare evolution patterns induced by the Gaussian and slow models, we take the ansatz (30) with  $\varphi_{\pi}(x)$  having a drastic shape of the Chernyak-Zhitnitsky (CZ) DA  $\varphi_{\pi}^{\text{CZ}}(x) = 30f_{\pi}x(1-x)(1-2x)^2$ . As one can see from Fig. 1, for  $P/\Lambda = 1$  the Gaussian model shows no indication of humps visible for higher  $P/\Lambda$  ratios. In the slow model, small humps are present even for  $P/\Lambda = 1$ . For high ratios  $P/\Lambda = 5$  and 10, the two models give close results, with strong humps.

Assuming  $\Lambda \sim 0.6$  GeV suggested by the VDA-based fits of the photon-pion transition form factor in Ref. [16], we expect that  $P \sim 3$  GeV would be required to support (or rule out) the CZ-type shape of the pion DA.

It is also interesting to note that the nonperturbative evolution pattern here is exactly opposite to the perturbative one. In the latter case, the humps of the initially CZ-shaped DA become less pronounced as the normalization scale increases and eventually disappear, with the DA tending to the asymptotic  $\sim x(1-x)$  shape.

To compare patterns of the QDA’s nonperturbative evolution for different shapes of the limiting DA, we take three models for  $\varphi_{\pi}(x)$ : Chernyak-Zhitnitsky  $\varphi_{\pi}^{\text{CZ}}(x)$ , flat  $\varphi_{\pi}^{\text{flat}}(x) = f_{\pi}$ , and asymptotic  $\varphi_{\pi}^{\text{as}}(x) = 6f_{\pi}x(1-x)$ . The results in the Gaussian and the slow models are rather similar. To avoid plotting too many graphs, we take, for definiteness, the slow model. Then, for the flat limiting DA we have

$$\frac{1}{f_{\pi}} Q_{\pi}^{\text{S.flat}}(y, P) = \frac{P}{\Lambda} \int_0^1 dx e^{-2|x-y|P/\Lambda}. \quad (35)$$

This integral can be calculated analytically. Writing  $y = (1 + \eta)/2$  in terms of a symmetric variable  $\eta$ , we get

$$\begin{aligned} \frac{1}{f_{\pi}} Q_{\pi}^{\text{S.flat}}(y, P) &= (1 - e^{-P/\Lambda} \cosh(P\eta/\Lambda))\theta(|\eta| \leq 1) \\ &+ \sinh(P/\Lambda) e^{-P|\eta|/\Lambda} \theta(|\eta| \geq 1). \end{aligned} \quad (36)$$

Similar, but more lengthy expressions may be obtained for two other models. As one can see from Fig. 2, for small  $P = \Lambda$  we have very close curves. For larger  $P = 3\Lambda$  the difference becomes visible, and for large  $P = 5\Lambda$  and  $P = 10\Lambda$  the curves shown in Fig. 3 are distinctly different. In fact, the  $P = 10\Lambda$  curves are very close to their limiting forms. Again, the nonperturbative evolution pattern in case of the flat DA is opposite to the perturbative one: as  $P$  increases,  $Q_{\pi}^{\text{flat}}(y, P)$  broadens from a rather narrow function for  $P = \Lambda$  and becomes almost constant for  $P = 10\Lambda$ .




FIG. 1. Quasidistribution amplitude  $Q_{\pi}^{\text{CZ}}(y, P)$  for  $P/\Lambda = 1, 3, 5, 10$  in the Gaussian (left) and slow models (right).




FIG. 2. Quasidistribution amplitudes  $Q_\pi(y, P)$  in the slow model for  $P = \Lambda$  (left) and  $P = 3\Lambda$  (right) evolving to CZ, flat and asymptotic DAs.




FIG. 3. Quasidistribution amplitudes  $Q_\pi(y, P)$  in the slow model for  $P = 5\Lambda$  (left) and  $P = 10\Lambda$  (right) evolving to CZ, flat and asymptotic DAs.

## VI. LEADING-ORDER HARD TAIL

The nonperturbative evolution of  $Q_\pi(y, P)$  essentially stops for  $P/\Lambda \gtrsim 20$ , and for larger values of  $P$  the dominant role is played by the perturbative evolution induced by the hard part. In our papers [15,16], it was suggested to take a purely soft TMDA (or VDA) as a starting approximation, and then “generate” hard tail by adding one-gluon exchanges. The only new parameter is the overall factor  $\alpha_s$ , while the  $k_\perp$  dependence of the hard tail of the TMDA  $\Psi(x, k_\perp^2)$  is completely determined by the soft part.

For large  $k_\perp$ , the generated hard part of the TMDA has a  $\sim 1/k_\perp^2$  behavior, but its explicit functional form is much more complicated. In particular, it is finite in the  $k_\perp \rightarrow 0$  limit [16]. The infrared cutoff for the naive  $1/k_\perp^2$  extrapolation is provided by the finite size of the pion encoded in the parameters, like  $\Lambda$ , present in the soft TMDA. Postponing the analysis of the interplay between the nonperturbative and perturbative evolution for future studies, we just outline below the VDA treatment of the hard tail.

For large  $\sigma$ , the lowest-order (in  $\alpha_s$ ) hard tail has the form

$$\Phi^{\text{hard}}(x, \sigma) = \Delta(x)/\sigma, \quad (37)$$

with  $\Delta(x)$  given by

$$\Delta(x) = \int_0^1 dz V(x, z) \varphi_\pi^{\text{soft}}(z), \quad (38)$$

where  $V(x, z)$  is the perturbative evolution kernel [21–23]. The asymptotic form (37) corresponds to a  $\sim 1/k_\perp^2$  TMDA, which is singular for  $k_\perp = 0$ . As explained above, this singularity is absent in the exact (rather complicated) expression for the hard tail. For illustration purposes, we take now the simplest regularization  $1/k_\perp^2 \rightarrow 1/(k_\perp^2 + m^2)$ . It corresponds to the change  $1/\sigma \rightarrow e^{-im^2/\sigma}/\sigma$  in the hard part of VDA,

$$\Phi^{\text{hard}}(x, \sigma) \rightarrow \frac{\Delta(x)}{\sigma} e^{-im^2/\sigma}. \quad (39)$$



To proceed with the conversion formula, one needs the integral over  $\sigma$

$$I(x, y, P) = \int_0^\infty \frac{d\sigma}{\sqrt{\pi\sigma}} \frac{P}{\sigma} e^{-(x-y)^2 P^2 / \sigma - m^2 / \sigma} = \frac{1}{\sqrt{(x-y)^2 + m^2 / P^2}}. \quad (40)$$

This gives the hard part of the quasidistribution amplitude

$$Q_\pi^{\text{hard}}(y, P) = \int_0^1 dx \frac{\Delta(x)}{\sqrt{(x-y)^2 + m^2 / P^2}}. \quad (41)$$

It generates evolution with respect to  $P^2$  in the form

$$P^2 \frac{d}{dP^2} Q_\pi^{\text{hard}}(y, P) = \frac{m^2}{2P^2} \int_0^1 dx \frac{\Delta(x)}{[(x-y)^2 + m^2 / P^2]^{3/2}}. \quad (42)$$

Taking the  $m/P \rightarrow 0$  limit we have

$$\frac{m^2}{2P^2} \int_0^1 dx \frac{V(x, z)}{[(x-y)^2 + m^2 / P^2]^{3/2}} = V(y, z) + \mathcal{O}(m^2 / P^2), \quad (43)$$

i.e., for large  $P^2$  the quasidistribution amplitude evolves according to the perturbative evolution equation with respect to  $P^2$ . This evolution is completely determined by the form of the soft DA  $\varphi^{\text{soft}}(x)$ . When the model for the latter is fixed, the particular choice of the soft TMDA  $\Psi^{\text{soft}}(x, k_\perp^2)$  does not affect the form of the hard part and the perturbative  $P$  evolution of the pion QDA  $Q(y, P)$ .

## VII. CONCLUSIONS

In this paper, we extended the approach of Ref. [14], where we have been dealing with the parton distribution

functions, the basic ingredients of perturbative QCD analysis of hard inclusive processes. Now we have dealt with the pion distribution amplitude, the basic ingredient of hard exclusive processes involving the pion. We applied the formalism of virtuality distribution amplitudes to study the  $p_3$  dependence of quasidistribution amplitudes  $Q_\pi(y, p_3)$ .

Just like in Ref. [14], we have established a simple relation between QDAs and TMDAs that allows us to derive models for QDAs from the models for TMDAs. Unlike the PDF case, there are many drastically different models claimed to describe the primordial shape of the pion DA. We have presented the  $p_3$ -evolution patterns for models producing some popular proposals: Chernyak-Zhitnitsky, flat, and asymptotic DAs. Our results may be used as a guide for future studies of the pion distribution amplitude on the lattice using the quasidistribution approach.

As our estimates show, one would need  $P$  of the order of a few GeV for the nonperturbative evolution to settle. It is natural to expect that perturbative evolution will be rather important at such scales. Thus, an interesting and technically challenging question for future studies is the interplay between the nonperturbative and perturbative evolution of the pion quasidistribution amplitude.

Another interesting problem for future studies is the analysis of more complicated models for TMDAs, in particular, models with nonfactorized  $k_\perp$  and  $x$  dependence. As the simplest generalization of the models used in the present paper, one can take  $x$ -dependent functions  $\Lambda(x)$  instead of the constant values  $\Lambda$ .

## ACKNOWLEDGMENTS

This work is supported by Jefferson Science Associates, LLC under U.S. DOE Contract No. DE-AC05-06OR23177 and by U.S. DOE Grant No. DE-FG02-97ER41028.

- 
- [1] V. M. Braun, S. Collins, M. Göckeler, P. Pérez-Rubio, A. Schäfer, R. W. Schiel, and A. Sternbeck, Second Moment of the pion light-cone distribution amplitude from lattice QCD, *Phys. Rev. D* **92**, 014504 (2015).
- [2] X. Ji, Parton Physics on a Euclidean Lattice, *Phys. Rev. Lett.* **110**, 262002 (2013).
- [3] H. W. Lin, Calculating the  $x$  dependence of hadron parton distribution functions, *Proc. Sci.*, LATTICE2013 (2014) 293.
- [4] C. Alexandrou, K. Cichy, V. Drach, E. Garcia-Ramos, K. Hadjiyiannakou, K. Jansen, F. Steffens, and C. Wiese, Lattice calculation of parton distributions, *Phys. Rev. D* **92**, 014502 (2015).
- [5] H. W. Lin, Bjorken- $x$  dependence of hadronic structure from lattice QCD, *Few Body Syst.* **56**, 455 (2015).
- [6] J. W. Chen, S. D. Cohen, X. Ji, H. W. Lin, and J. H. Zhang, Nucleon helicity and transversity parton distributions from lattice QCD, *Nucl. Phys.* **B911**, 246 (2016).
- [7] C. Alexandrou, K. Cichy, K. Hadjiyiannakou, K. Jansen, F. Steffens, and C. Wiese, A lattice calculation of parton distributions, *Proc. Sci.*, DIS2016 (2016) 042.
- [8] C. Alexandrou, Parton distribution functions from Lattice QCD, *Few Body Syst.* **57**, 621 (2016).
- [9] X. Xiong, X. Ji, J. H. Zhang, and Y. Zhao, One-loop matching for parton distributions: Nonsinglet case, *Phys. Rev. D* **90**, 014051 (2014).

- [10] V. N. Gribov and L. N. Lipatov, Deep inelastic e p scattering in perturbation theory, *Yad. Fiz.* **15**, 781 (1972) [*Sov. J. Nucl. Phys.* **15**, 438 (1972)].
- [11] L. N. Lipatov, The parton model and perturbation theory, *Yad. Fiz.* **20**, 181 (1974) [*Sov. J. Nucl. Phys.* **20**, 94 (1975)].
- [12] G. Altarelli and G. Parisi, Asymptotic freedom in parton language, *Nucl. Phys.* **B126**, 298 (1977).
- [13] Y. L. Dokshitzer, Calculation of the structure functions for deep inelastic scattering and  $e^+e^-$  annihilation by perturbation theory in quantum chromodynamics, *Zh. Eksp. Teor. Fiz.* **73**, 1216 (1977) [*Sov. Phys. JETP* **46**, 641 (1977)].
- [14] A. Radyushkin, Nonperturbative evolution of parton quasi-distributions, *Phys. Lett. B* **767**, 314 (2017).
- [15] A. V. Radyushkin, Virtuality distributions in application to  $\gamma\gamma^* \rightarrow \pi^0$  transition form factor at handbag level, *Phys. Lett. B* **735**, 417 (2014).
- [16] A. V. Radyushkin, Virtuality and transverse momentum dependence of the pion distribution amplitude, *Phys. Rev. D* **93**, 056002 (2016).
- [17] L. Gamberg, Z. B. Kang, I. Vitev, and H. Xing, Quasi-parton distribution functions: A study in the diquark spectator model, *Phys. Lett. B* **743**, 112 (2015).
- [18] I. Vitev, L. Gamberg, Z. Kang, and H. Xing, A study of quasi-parton distribution functions in the diquark spectator model, *Proc. Sci.*, QCDEV2015 (2015) 045.
- [19] A. Bacchetta, M. Radici, B. Pasquini, and X. Xiong, Reconstructing parton densities at large fractional momenta, *Phys. Rev. D* **95**, 014036 (2017).
- [20] A. V. Radyushkin, Deep elastic processes of composite particles in field theory and asymptotic freedom, [arXiv: hep-ph/0410276](https://arxiv.org/abs/hep-ph/0410276).
- [21] A. V. Efremov and A. V. Radyushkin, Asymptotical behavior of pion electromagnetic form-factor in QCD, *Teor. Mat. Fiz.* **42**, 147 (1980) [*Theor. Math. Phys.* **42**, 97 (1980)].
- [22] A. V. Efremov and A. V. Radyushkin, Factorization and asymptotical behavior of pion form-factor in QCD, *Phys. Lett.* **94B**, 245 (1980).
- [23] G. P. Lepage and S. J. Brodsky, Exclusive processes in quantum chromodynamics: Evolution equations for hadronic wave functions and the form-factors of mesons, *Phys. Lett.* **87B**, 359 (1979).
- [24] A. V. Efremov and A. V. Radyushkin, Report No. JINR-E2-11535, 1978.
- [25] V. L. Chernyak and A. R. Zhitnitsky, Exclusive decays of heavy mesons, *Nucl. Phys.* **B201**, 492 (1982); Erratum: Exclusive decays of heavy mesons, *Nucl. Phys.* **B214**, 547 (E) (1983).
- [26] F. M. Dittes and A. V. Radyushkin, Radiative Corrections to the pion form-factor in quantum chromodynamics, *Yad. Fiz.* **34**, 529 (1981) [*Sov. J. Nucl. Phys.* **34**, 293 (1981)].
- [27] I. V. Anikin, A. E. Dorokhov, and L. Tomio, On high  $Q^2$  behavior of the pion form-factor for transitions  $\gamma^*\gamma \rightarrow \pi^*0$  and  $\gamma^*\gamma^* \rightarrow \pi^0$  within the nonperturbative approach, *Phys. Lett. B* **475**, 361 (2000).
- [28] E. Ruiz Arriola and W. Broniowski, Pion light cone wave function and pion distribution amplitude in the Nambu-Jona-Lasinio model, *Phys. Rev. D* **66**, 094016 (2002).
- [29] A. V. Radyushkin, Shape of pion distribution amplitude, *Phys. Rev. D* **80**, 094009 (2009).
- [30] M. V. Polyakov, On the pion distribution amplitude shape, *JETP Lett.* **90**, 228 (2009).
- [31] S. V. Mikhailov and A. V. Radyushkin, Nonlocal condensates and QCD sum rules for pion wave function, *Pis'ma Zh. Eksp. Teor. Fiz.* **43**, 551 (1986) [*JETP Lett.* **43**, 712 (1986)].
- [32] G. P. Lepage and S. J. Brodsky, Exclusive processes in perturbative quantum chromodynamics, *Phys. Rev. D* **22**, 2157 (1980).
- [33] I. V. Musatov and A. V. Radyushkin, Transverse momentum and Sudakov effects in exclusive QCD processes:  $\gamma^*\gamma\pi^0$  form-factor, *Phys. Rev. D* **56**, 2713 (1997).