

2022

Lower Extremity Joint Stiffness During Running in Adolescents with Autism Spectrum Disorder

Lauren A. Luginsland MS
Old Dominion University

Justin A. Haegele PhD
Old Dominion University

Hunter J. Bennett PhD
Old Dominion University

Follow this and additional works at: https://digitalcommons.odu.edu/gradposters2022_education



Part of the [Sports Sciences Commons](#)

Recommended Citation

Luginsland, Lauren A. MS; Haegele, Justin A. PhD; and Bennett, Hunter J. PhD, "Lower Extremity Joint Stiffness During Running in Adolescents with Autism Spectrum Disorder" (2022). *College of Education & Professional Studies (Darden) Posters*. 4.
https://digitalcommons.odu.edu/gradposters2022_education/4

This Book is brought to you for free and open access by the 2022 Graduate Research Achievement Day at ODU Digital Commons. It has been accepted for inclusion in College of Education & Professional Studies (Darden) Posters by an authorized administrator of ODU Digital Commons. For more information, please contact digitalcommons@odu.edu.



LOWER EXTREMITY JOINT STIFFNESS DURING RUNNING IN ADOLESCENTS WITH AUTISM SPECTRUM DISORDER

Lauren A. Luginsland, Hunter J. Bennett, Justin A. Haegele
Old Dominion University, Norfolk, VA, USA
Contact Email: llugi001@odu.edu



Introduction

- Autism spectrum disorder (ASD) is a neurodevelopmental disorder characterized by a variety of physiological and behavioral characteristics. ASD affects 1 in 44 children in the United States¹
- Running is the most common form of physical activity for girls and the second most common form for boys aged 12 to 15 years.² This is consistent for autistic adolescents, who enjoy solitary activities, such as running, more than team-based sports³⁻⁵
- Autistic adolescents have reported elevated levels of fear for sustaining injury, being bullied, and fear of exclusion within their physical education classes compared to their non-autistic peers⁵
- Late adolescents undergo rapid skeletal growth leaving individuals more vulnerable to injury risk. Inadequate joint stiffness is one of several factors that may increase injury risk⁶
- The purpose of this study was to examine ankle and knee joint stiffness in autistic adolescents and non-autistic matched controls at self-selected and matched running speeds

Significance

- Investigating loading and joint stiffness of the lower extremity in autistic persons during running may provide points of emphasis for therapeutic and rehabilitation interventions

Methods

- Twenty-two (n=22) autistic adolescents (15 males, aged 14±2 years BMI 22.24±5.76 kg/m²) and seventeen (n=17) age, sex, and body mass index healthy non-autistic controls (CON) participated
- Participants ran at two matched speeds: self-selected speed (SS) and a standardized speed of 3.0 m/s (3.0)
- Three-dimensional motion capture (Vicon Vantage) and force platform (Bertec) data were recorded. Figure 1 provides an example of the progression from motion capture of a human participant (top) to skeletal model (bottom)
- Joint stiffness (unitless) was calculated as the quotient of the change in joint moment (normalized to body mass*leg length) and the change in joint angle (radians) during the energy absorption phase (Fig 2)
- Stiffness and changes in joint moments were analyzed using 2 (group) x 2 (speed) ANOVA

Table 1. Ensemble joint stiffnesses and change in joint moments (mean±SD)

	Joint Stiffness				Joint Moments			
	ASD SS	ASD 3.0	CON SS	CON 3.0	ASD SS	ASD 3.0	CON SS	CON 3.0
Knee	5.84±0.86	6.44±0.80	6.80±1.05	6.97±1.32	2.47±0.35	2.67±0.38	2.80±0.33	2.90±0.42
Ankle	7.49±1.15	7.33±1.43	8.88±2.35	8.77±2.28	2.28±0.35	2.38±0.38	2.81±0.42	2.82±0.43

Figure 1. Example of data collection, motion capture data in Vicon Nexus then analyzed in Visual 3D





Figure 2. Examples of knee (left) and ankle (right) joint stiffness waveforms



Results

- There were no significant interactions between groups and speeds (p>0.05) for any variable
- Autistic adolescents had reduced knee and ankle joint stiffness (all p<0.020; Table 1)
- Running at the 3.0m/s standardized speed resulted in increased knee joint stiffness compared to self-selected (p=0.010)
- Autistic adolescents had reduced changes in knee and ankle moments (p<0.003; Table 1)
- Running at the 3.0m/s standardized speed increased knee moments over self-selected speed (p=0.004)

Discussion

- Autistic adolescents typically display increased joint stiffness compared to controls.⁸ However, our expectation of a similar increase in joint stiffness during running was incorrect.
- Joint stiffness during running has typically been viewed as greater stiffness indicates an increase in injury risk⁷; thus, reduced stiffness could be beneficial
- However, decreased joint stiffness could be indicative of a less efficient running style whereby the elastic recoil is not being optimally utilized by the knee and ankle musculature
- We cannot ignore the implications of prior training on running mechanics. We did not ascertain participants' physical activity; however, we previously tracked physical activity engagement, finding no differences between groups in low and moderate to vigorous physical activity⁹

References

- Maenner, et al., 2021. *Surveillance Summaries*, 70(11); 1-16
- Fakhouri et al., 2014. *NCHS Data Brief*. 141; 1-8
- Healy et al., 2017. *Autism Research*. 47; 49-57
- Stanish et al., 2015. 32(4); 302-317
- Haegele et al., 2022. *Autism*. 26(1); 51-61
- Butler et al., 2003. *Clin Biomech*. 18(6); 511-517
- Brazier et al., 2014. *Strength and Cond*. 36(5); 103-12
- Eggleston et al., 2018. *Human Movement Sci*. 62; 25-33
- Haegele et al., 2021. *Disability and Health*. 14(3)