

**Old Dominion University
ODU Digital Commons**

Community & Environmental Health Faculty
Publications

Community & Environmental Health

10-2017

A Social-Ecological Framework of Theory, Assessment, and Prevention of Suicide

Robert J. Cramer

Old Dominion University, rccramer@odu.edu

Nester D. Kapusta

Follow this and additional works at: https://digitalcommons.odu.edu/commhealth_fac_pubs

 Part of the [Community Health and Preventive Medicine Commons](#), [Mental and Social Health Commons](#), and the [Psychology Commons](#)

Repository Citation

Cramer, Robert J. and Kapusta, Nester D., "A Social-Ecological Framework of Theory, Assessment, and Prevention of Suicide" (2017).
Community & Environmental Health Faculty Publications. 26.
https://digitalcommons.odu.edu/commhealth_fac_pubs/26

Original Publication Citation

Cramer, R. J., & Kapusta, N. D. (2017). A social-ecological framework of theory, assessment, and prevention of suicide. *Frontiers in Psychology*, 8, 1756. doi:10.3389/fpsyg.2017.01756



A Social-Ecological Framework of Theory, Assessment, and Prevention of Suicide

Robert J. Cramer^{1*} and Nestor D. Kapusta²

¹ School of Community and Environmental Health Sciences, Old Dominion University, Norfolk, VA, United States, ² Suicide Research Group, Department for Psychoanalysis and Psychotherapy, Medical University of Vienna, Vienna, Austria

OPEN ACCESS

Edited by:

Raffaella Calati,
Centre Hospitalier Universitaire de
Montpellier, France

Reviewed by:

Timo Partonen,
National Institute for Health and
Welfare, Finland
Gianluca Serafini,
Department of Neuroscience, San
Martino Hospital, University of Genoa,
Italy

*Correspondence:

Robert J. Cramer
rcramer@odu.edu

Specialty section:

This article was submitted to
Clinical and Health Psychology,
a section of the journal
Frontiers in Psychology

Received: 10 August 2017

Accepted: 22 September 2017

Published: 09 October 2017

Citation:

Cramer RJ and Kapusta ND (2017) A
Social-Ecological Framework of
Theory, Assessment, and Prevention
of Suicide. *Front. Psychol.* 8:1756.
doi: 10.3389/fpsyg.2017.01756

The juxtaposition of increasing suicide rates with continued calls for suicide prevention efforts begs for new approaches. Grounded in the Centers for Disease Control and Prevention (CDC) framework for tackling health issues, this personal views work integrates relevant suicide risk/protective factor, assessment, and intervention/prevention literatures. Based on these components of suicide risk, we articulate a Social-Ecological Suicide Prevention Model (SESPM) which provides an integration of general and population-specific risk and protective factors. We also use this multi-level perspective to provide a structured approach to understanding current theories and intervention/prevention efforts concerning suicide. Following similar multi-level prevention efforts in interpersonal violence and Human Immunodeficiency Virus (HIV) domains, we offer recommendations for social-ecologically informed suicide prevention theory, training, research, assessment, and intervention programming. Although the SESPModel calls for further empirical testing, it provides a suitable backdrop for tailoring of current prevention and intervention programs to population-specific needs. Moreover, the multi-level model shows promise to move suicide risk assessment forward (e.g., development of multi-level suicide risk algorithms or structured professional judgments instruments) to overcome current limitations in the field. Finally, we articulate a set of characteristics of social-ecologically based suicide prevention programs. These include the need to address risk and protective factors with the strongest degree of empirical support at each multi-level layer, incorporate a comprehensive program evaluation strategy, and use a variety of prevention techniques across levels of prevention.

Keywords: suicide, prevention, social-ecological model, risk assessment

Suicide rates in the United States are increasing in the last decade from 11.0 (per 100,000) in 2004 to 13.4 in 2014 (Drapeau and McIntosh, 2014). Most recent data summarized by the Centers for Disease Control and Prevention (CDC) echoes this pattern with detailed analyses showing that trend inclines may be moderated by factors such as gender, age, and method (Centers for Disease Control and Prevention, 2016a). What is particularly concerning is that the increasing suicide rate is occurring in the presence of a 2012 national suicide prevention strategy put forth by the U.S. Surgeon General's Office, which is based on four broad strategic directions reflecting a multi-level perspective: (1) create supportive environments promoting healthy, empowered persons, families, and communities; (2) enhance community-oriented prevention services; (3) promote timely, supportive services, and; (4) improve suicide-related surveillance data (United States Surgeon General's Office, 2012).

Despite progress in the effectiveness of suicide prevention efforts (Mann et al., 2005; Zalsman et al., 2016), suicide prevention still suffers from several critical limitations: the inability to predict suicidal behavior in individuals (Fowler, 2012; Chu et al., 2015; Chan et al., 2016), inconsistent suicide-related terminology (Skegg, 2005; Silverman and De Leo, 2016), lack of multi-level theoretical development (O'Connor, 2011; Barzilay and Apter, 2014), and insufficient implementation of multi-level prevention programs (Hegerl et al., 2008; van der Feltz-Cornelis et al., 2011). Informed by theoretical, risk/protective factor, and prevention program evidence, we articulate a conceptual multi-level framework for suicide prevention.¹ We further make recommendations concerning development of multi-level suicide risk theory, research, assessment and prevention.

THE STATE OF SUICIDE PREVENTION EFFORTS

The current scope of suicide prevention efforts spans primary prevention (e.g., public awareness campaigns), secondary prevention (e.g., gate-keeper training programs), tertiary prevention (e.g., psychotherapy), and postvention (e.g., survivor support groups). Extending these traditional categories on a mental health intervention spectrum, suicide prevention can be applied at a universal (i.e., to the general public), selective (e.g., groups defined by lifetime risk such as military personnel), and indicated prevention (i.e., high risk groups where risk is already elevated—e.g., psychiatric inpatients) levels (Institute of Medicine Committee on Prevention of Mental Disorders, 1994). A recent systematic review summarizes the overall state of effective suicide prevention programs across these levels (Zalsman et al., 2016). The findings support reduction of suicide-related thoughts and behaviors (i.e., ideation, attempts and completed suicide) for: (1) restricting access to lethal means (e.g., hot-spots for jumping), (2) school-based awareness programs, (3) lithium and clozapine use, and (4) psychotherapeutic efforts for depression. Authors also noted a lack of current evidence for an array of other prevention approaches (e.g., gatekeeper training, physician and public education).

Of the few approaches that cut across more than one of multiple potential levels of prevention, recent efforts by the United States (US) National Action Alliance for Suicide Prevention have focused on initiatives (e.g., Zero Suicide, Vision Zero) toward the goal of absolute elimination of suicide (Erlich, 2016). Components of these approaches range from improving follow-up practices with patients post-discharge and maintaining contact with at-risk persons to enhancing infrastructure (e.g., personnel, training content) and prevention resources. Two promising tertiary suicide prevention strategies common to clinical psychiatry are Dialectical Behavior Therapy (DBT) (Comtois and Linehan, 2006) and the Collaborative Assessment and Management of Suicide (CAMS) (Ellis et al., 2015). A noteworthy gap is that, even clinical or targeted approaches

showing potential effectiveness are unable to simultaneously target the individual through societal level influences on suicide risk.

A MULTI-LEVEL UNDERSTANDING OF SUICIDE PREVENTION

We echo other calls in the literature for a multi-level public health approach to suicide prevention (Dahlberg and Krug, 2002; van der Feltz-Cornelis et al., 2011). The CDC provides valuable guidance based on the assumption that prevention efforts for any health or disease issue require integrated multi-level efforts within a Social-Ecological Model (SEM) (Centers for Disease Control Prevention, 2017). The SEM is a four tier framework for organizing risk and protective factors, which then inform corresponding prevention strategies. From macro to micro levels, the four strata are: societal, community, relational, and individual levels. Societal factors concern larger scale issues such as social and cultural norms, policies, and other guiding rules or laws. Community level influences are those circumscribed to a certain region like neighborhood centers, schools, workplaces and healthcare providers. Relational factors are those defined by direct person-to-person interaction such as social support or withdrawal, peers, and family. Individual level factors pertain to person characteristics such as demographics, attitudes, health conditions, and others. The SEM has been meaningfully applied to a range of health issues and prevention programs such as health literacy (McCormack et al., 2017) and vaccine usage (Kumar et al., 2009).

We see at least three straightforward benefits of such a multi-level schema. First, suicide risk and protective factor literature tends to be fragmented by SEM level. That is, even where summaries of risk factors are provided, they are often limited to one or two SEM levels. An SEM of suicide prevention, therefore, provides a potentially comprehensive framework for organizing risk and protective factor knowledge; as such, it is a working template for adding new factors, as well as integrating levels to examine how upper level factors may moderate the influence of lower level factors, and vice versa.

Following from enhanced organization of factors, a second benefit is that an SEM of suicide prevention can provide grounding for multi-level intervention and prevention program design and implementation. This idea has been demonstrated by closely-related comprehensive approaches to prevention of gun violence prevention (Rubens and Shehadeh, 2014) and campus sexual assaults (Centers for Disease Control Prevention, 2016b). For instance, Rubens and Shehadeh organized potential interventions and preventions for gun violence in the US along levels of the SEM, noting potential strategies ranging from individual (e.g., parent-child relationships) to societal (e.g., financial liability for those violating gun safety norms) approaches (Rubens and Shehadeh, 2014). Finally, articulation of a multi-level approach to suicide prevention can provide a framework for the re-organization of current theories of suicide. That is, to date causal theories of suicide consistently fail to fully integrate multi-level perspectives. It is our hope that a

¹Authors recognize not every construct can be defined due to the brevity of personal views format articles.

social-ecological view of suicide prevention would spur growth and effectiveness in theory and practice.

TOWARD A SOCIAL-ECOLOGICAL MODEL OF SUICIDE PREVENTION

In support of a multi-level approach to suicide prevention, Caine proposed to frame suicide prevention within an SEM model in terms of its shared risk with interpersonal violence (Caine, 2013). However, the resulting ecological model of shared risk was limited in scope in terms of merely listing sample risk and protective factors in common for both suicide and interpersonal violence. Extending this approach we articulate a comprehensive picture of risk and protective factors associated with at least one aspect of suicide-related thoughts and behavior, yielding the SESPM.

Search Strategy and Selection Criteria

² In order to balance comprehensiveness of sources cited, while also recognizing brevity of this manuscript format, we did the following to identify sources to inform the SESPM. We searched Pubmed, Medline, Psychinfo, and Psycharticles using combinations of the following phrases: "suicide," "risk factor," "protective factor," "prevention," "intervention," "review," and "meta-analysis" while focusing on articles from 1980 to present. Reviews and meta-analyses were given priority because we aimed to provide a big picture review (see Table 1). We further used Google Scholar to identify pertinent content from the following major professional organizations: American Foundation for Suicide Prevention, Suicide Prevention Resources Center, American Association of Suicidology, World Health Organization, CDC, and Substance Abuse and Mental Health Services Administration. Once a full set of key sources was identified, we dropped sources that were completely redundant with others.

Selection of the final integrated body of existing evidence represented scoping/conceptual summaries (Bryan and Rudd, 2006; Van Orden et al., 2010; Drapeau and McIntosh, 2014; Bernard et al., 2015), systematic reviews/meta-analyses (Serafini et al., 2012; Calear et al., 2016; Chan et al., 2016; Ma et al., 2016; Zalsman et al., 2016; Franklin et al., 2017), mortality risk studies (Björksenstam et al., 2015, 2016; Madsen et al., 2017), measure development (Linehan et al., 1983), and policy analysis (Anestis and Anestis, 2015) into a unified SESPM framework, as presented in Table 1. In doing so, we differentiate factors widely applicable across groups vs. those that tend to demonstrate population-specific associations with suicide (e.g., military veterans, youth, lesbian, gay, bisexual and transgender [LGBT] persons). For example, concerning LGBT youth, literature consistently links population-specific experiences of internalized stigma and victimization as associated with suicide risk; moreover, sexual orientation minority status itself is linked with elevated suicide risk (Haas et al., 2011; Duncan and Hatzenbuehler, 2014). The need for attention to nuance

of even strong risk factors varying by population is further illustrated by primary psychiatric diagnoses linked to suicide. For instance, standardized mortality risk (SMR) and other research documents the exacerbated prominence of depression, bipolar, and cluster B personality disorders (e.g., borderline, antisocial) in enhancing risk for death by suicide among psychiatric patients (Björksenstam et al., 2015, 2016; Madsen et al., 2017). To illustrate, cluster B disorders are associated with SMRs in this population as high as 33–34 (Björksenstam et al., 2015; Madsen et al., 2017). Cannabis use and dependence, another diagnostically relevant disorder category, has been shown to be associated with exacerbated suicide risk, especially among adolescents and particularly when the cannabis is associated with experiencing of other psychiatric conditions (e.g., psychosis) (Serafini et al., 2012). Thus, cannabis use or dependence may also serve as poor coping or a pathway to suicide among adolescent youth.

We also note many factors with the strongest, most consistent associations with suicide risk (see Table 1), defining strongest and consistent in terms of effect sizes and odds ratios related to suicide-related thoughts or behaviors, direct associations with suicide (e.g., serving as a mediator), as well as those that are highlighted by clinical and prevention experts as those requiring attention across populations (e.g., depression). In all, SEM levels with the strongest support tend to be individual and interpersonal/relational levels. For instance, at the individual or interpersonal levels we note risk factors with the strongest associations with suicide-related thoughts and behavior such as a prior suicide attempt, diagnosis of depression or bipolar disorders, and suicide contagion. To illustrate, hopelessness provides a clear example of an individual psychological risk factor with considerable support; hopelessness has been identified as an independent risk factor for suicide requiring clinical assessment (Bryan and Rudd, 2006), and empirical data raises the potential that hopelessness may serve as a pathway to suicide-related thoughts and behavior explaining the influence of other risk factors (e.g., thinking styles) (Abramson et al., 1998). Moreover, protective factors such as presence, use and perception of positive social support is denoted as among the strongest factors (see Table 1 for full list of demarcated factors with strongest research support). As such, from a public health education standpoint, the integrated summary may serve to reinforce the key factors to include in dissemination efforts by public organizations.

Given the fact that suicidology is an ongoing research field, the proposed SESPM is not intended to be exhaustive. The purpose of the SESPM is a guide to move research and prevention forward, as well as to provide a framework for understanding nuance in suicide prevention. To illustrate the latter point, examination of the multi-level organization identifies several levels at which for example firearm-related factors may influence suicide risk. While firearm access or ownership is associated with elevated suicide risk (Anglemeyer et al., 2014), this link may be moderated by other individual (e.g., safe storage) (McCarten et al., 2001), relational (e.g., restricted means counseling) (Stanley et al., 2016), and societal (e.g., firearm restriction laws) (Anestis and Anestis, 2015) factors. The SESPM provides a summary of literature in order to build toward better mediation,

²Our paper is not intended to be a systematic review; therefore, full search criteria and other details are not reported.

TABLE 1 | Compilation of major suicide risk and protective factors organized by levels of centers for disease control and prevention's social-ecological model.

Risk factors	Protective factors
Societal: Economic downturn/depression Living location with less restrictive firearm laws Seasonal variation Stigma about mental health and treatment Air pollutants Viruses/parasites Poverty Mountain region of the US Western and southern US	Societal: Healthy economy Living location with more restrictive firearm laws Mental health funding Northeast US
Community: Exposure to community violence Local suicide epidemic Barriers to healthcare access	Community: Crisis support lines/hotlines Healthcare/mental healthcare access Effective mental healthcare Trained gate keepers Community involvement School-based support and intervention programming*
Interpersonal/Relationship: Living in household with firearm Exposure to suicide/contagion Family violence Family conflict Family history of mental illness Family history of suicide/attempts Relationship instability Death of a loved one Severing of romantic relationship Social isolation/withdrawal Combat exposure*	Interpersonal/Relationship: Presence of social support Use of social support Perceived social support Concerns suicide is harmful to child/family Sense of responsibility to family Healthy long-term committed relationship/marriage Help-seeking behavior Children present in the home Pregnancy* Pulling together Caring letters Social connectedness Contact with caregivers* Support for connection with healthcare providers Cognitive-behavioral therapy Dialectical-behavior therapy Collaborative assessment and management of suicide (CAMS)
Individual: Biological Male sex (completions)/Female sex (attempts)* Serotonin dysfunction Family history of suicidal behavior	Individual: Biological SSRI usage Lithium/mood stabilizer treatment Clozapine usage
Socio-Demographic Gender (e.g., Transgender status) Lesbian, gay, bisexual or other sexual orientation minority identity* Religiosity/spirituality (i.e., suicide as a resolution to problems)* Native American ethnicity* Hispanic ethnicity* Asian/Pacific Islander ethnicity* Whites (compared to non-Whites)* Older adult age* Middle adult age* High risk professions (e.g., military, law enforcement)* Firearm ownership (and unlocked, loaded)* Incarceration* High perceived/subjective stress Job loss/unemployment Financial strain Recent discharge from psychiatric hospital* Bullying/bias crime victimization*	Socio-Demographic Heterosexual sexual orientation Religiosity/spirituality (i.e., beliefs about suicide being wrong)*

(Continued)

TABLE 1 | Continued

Risk factors	Protective factors
<p>Psychiatric</p> <p>Mental health diagnoses/symptoms such as depression, bipolar, post-traumatic stress disorder, anxiety, and active phase schizophrenia</p> <p>Personality disorders such as Borderline Personality</p> <p>Substance use/abuse (e.g., cannabis)</p> <p>Alcohol use/abuse</p>	<p>Psychiatric</p> <p>Treatment motivation</p>
<p>Psychological</p> <p>Prior suicide attempt</p> <p>Current suicidal thinking</p> <p>Presence of suicidal intent</p> <p>Presence of suicide plan</p> <p>Access to/presence of lethal means</p> <p>Preparatory behaviors (e.g., giving away prized possessions)</p> <p>Prior or current non-suicidal self-injury</p> <p>History of other suicide (e.g., ideation)</p> <p>Hopelessness</p> <p>Low self-control/high impulsivity</p> <p>Aggression</p> <p>Agitation</p> <p>Emotion dysregulation</p> <p>Severe mood change</p> <p>Childhood abuse</p> <p>Feelings of burdensomeness</p> <p>Rejection/thwarted belonging</p> <p>Chronic illness*</p> <p>Acute health symptoms*</p> <p>Fatigue</p> <p>Sleep disturbance/disorders</p> <p>Neuroticism</p> <p>Introversion</p> <p>Limited openness to experience</p> <p>Perfectionism</p> <p>Homelessness*</p> <p>Low self-esteem</p> <p>Shame</p> <p>Physical pain tolerance</p> <p>Fearlessness of suicide/death</p> <p>Thinking errors/negative thinking</p> <p>Psychache/psychic pain</p> <p>Internalized stigma*</p>	<p>Psychological</p> <p>Coping skills</p> <p>Problem solving skills</p> <p>Moral objections to suicide</p> <p>Survival beliefs/desire to live</p> <p>Fear of suicide/death</p> <p>Fear of social disapproval</p> <p>Optimism</p> <p>Hopefulness/positive future orientation</p> <p>Life satisfaction</p> <p>Intact reality testing</p> <p>High self-esteem/self-efficacy</p> <p>Resiliency</p> <p>Extraversion</p> <p>Additional reasons for living</p>

*Risk or protective factor demonstrating unique importance for a specific population.

Bold italics font, strongest risk/protective factor for suicide risk.

moderation, and causal research, as well as multi-level prevention efforts.

RECOMMENDATIONS FOR SESPM-INFORMED RESEARCH, THEORY AND PROGRAMMING

Consequently, we advocate five next steps for the suicidology field.

Empirical Testing and Adaptation

We adopt the view that the SESPM is both preliminary and fluid, suggesting prevention may need to account for population- or context-specific considerations. For instance, the SESPM

itself may vary by population or culture. To illustrate, it is well known that risk factors such as Human Immunodeficiency Virus (HIV) status itself (Carrico et al., 2007) and internalized HIV-related stigma (Cramer et al., 2015) play particularly salient roles in suicide risk, whereas other factors may be less important for this group. Additionally, the SESPM offers a clear organizational approach to future systematic reviews and hierarchical approaches to meta-analysis or regression. In all, with future empirical testing, the SESPM may need refinement or adaptation by population over time. We advocate in a first step to conduct risk and protective factor meta-analyses to develop appropriate SESPM templates for risk groups. In a second step, this quantitative information about the weight of risk factors should be used in population trials consequently (see Figure 1).




TABLE 2 | Sample suicide prevention and intervention strategies by level of the social-ecological model.

Social-ecological model level	Sample intervention and prevention programming	Sample theory of suicide
Societal	1. Firearm laws or regulations concerning storage, mental health background checks, etc. 2. Public awareness campaign targeting mental health and therapy stigma reduction. 3. Suicide-specific federal funding initiatives.	Sociological theory of suicide
Community	1. Crisis support lines. 2. Free mental health screenings provided by community mental health centers or in clinics treating high risk populations. 3. School-based programs targeting diversity-related social norms, mental health care access, or suicide awareness.	Military transition theory
Relational	1. Group psychotherapy. 2. Individual psychotherapy. 3. Gate keeper training.	Interpersonal-psychological theory of suicide
Individual	1. Adoption of positive health behaviors (e.g., exercise, food choices, sleep hygiene) 2. Mental health literacy courses. 3. Positive coping skills training/adoption.	Cubic model of suicide

A Framework for Public Health Education and Training Efforts

Promising suicide prevention education and training programming exists for the public (Teo et al., 2016) and medical/health professionals (Cramer et al., 2017), yet these areas are in need of further study (Zalsman et al., 2016). Moreover, empirically-tested educational prevention strategies often lack consistent structural framing. That is, content and modalities of these trainings vary, often neglecting content such

as community and societal level risk factors and prevention efforts. As has been done in the development of other public health prevention approaches such as HIV prevention (Baral et al., 2013), we encourage development of research/data summaries, educational materials, and training content to be organized by SESPM levels. For example, graduate training or continuing education programs may address established suicide prevention-related competencies (Rudd et al., 2008; Cramer et al., 2013) by SESPM level. While a full program-wide

review of such competencies is beyond the scope of this piece, using structured training approaches like observed structured clinical examinations (OSCEs), health professions literature (Hung et al., 2012; Cramer et al., 2016) highlights necessary skills for health providers like knowing empirically-indicated risk/protective factors and intervention/support possibilities. Established training models such as patient simulation or online-mediated courses can integrate such skill development and factual content into a SESPM framework. The end goal of such an approach would be that SESPM-educated health clinicians may be able to make better use of their multi-level knowledge and skills in working with at-risk individuals or designing stronger prevention programs.

Multi-Level Suicide Risk Theory

Historically, public health and health science prevention efforts have lacked adequate theoretical grounding. Recent health professions literature argues that effective prevention efforts require strong grounding to bolster effective health behavior change (Im, 2015; Prestwich et al., 2015; Krieger, 2016). The advantages of theory-informed public health include conceptualization of multi-level prevention/intervention programming, transdisciplinary communication, and accounting for practical societal and scientific influences (e.g., funding, political issues). Interestingly, social and behavioral science literature focuses on suicide as the subject of theoretical speculation, although there are varying levels of empirical testing and support across these theories.

Although a full review of all contemporary suicide risk theories is beyond the scope of this work, we provide an example theory with short description for each SESPM level for illustrative purposes (see **Table 2**). From top down, example theories can be seen in societal (e.g., *Le Suicide*) (Durkheim, 1897), community (e.g., Military Transition Theory) (Castro and Kintzle, 2014), relational (e.g., Interpersonal-Psychological Theory of Suicide) (Joiner, 2005; Van Orden et al., 2010), and individual (e.g., Cubic Model of Suicide) (Shneidman, 1981) level perspectives. In his seminal text *Le Suicide*, sociologist Emil Durkheim theorized suicide as a reaction to the intersection of social integration (the clustering of people in social groups) and regulation (the extent of rituals and customs being influenced by societal norms) (Lester, 1999). Rooted in a litany of causes for suicide among US veterans, Military Transition Theory highlights suicide risk as a function of factors unique to the community of military personnel reintegrating into civilian life (Castro and Kintzle, 2014). A three-stage transition is posited: approaching, managing, and assessing the transition. These stages imply a degree of multi-level influence in that they require the person to navigate and evaluate individual, familial, work and other challenges.

The Interpersonal-Psychological Theory of Suicide (IPTS) posits that suicide thinking is a function of self-perceptions in relation to others; negative interpersonal cognitions occur in two forms: thwarted belonging and perceived burdensomeness (Van Orden et al., 2010). Ideation transitions to an attempt when the individual has developed sufficient habituation to pain and fearlessness of death in order to commit the act. In this way, the IPTS may be considered both individual and relational in nature. Finally, Shneidman articulated an individual theory of impulsive

suicide (Shneidman, 1981). The impulsive act, calculated to be fatal by the attempter, is thought to occur in the presence of acute psychological states of stress, agitation and psychache (i.e., emotional pain).

Relying on the SESPM organizational framework of risk and protective factors (see **Table 1**), we believe a valuable next step in the theoretical development and testing is a *multi-level or social-ecological theory of suicide* in both population-based and clinical trials, as well as in interventions (see **Figure 1**) in order to inform theory development and prevention programming. The “x” lines refer to the idea that suicide research suffers from the problem of repeated over simplification of studies bypassing a comprehensive use or development of a multi-level model. Therefore, as illustrated in **Figure 1**, it is critical to understand that the proposed SESPM, a multi-level organizational framework, does not constitute a true theory of suicide by itself. However, following the pathways outlined in the Figure, a consistent and causal theory of suicidal behavior should be deductible from such a framework, and therefore be empirically testable (Horvath, 2016). A social-ecological theory of suicide, for instance, may specify individual attitudes, traits and mental health symptoms as primary, direct predictors of imminent suicide risk. Complementing that testable hypotheses, societal or community level factors, as well as sub-population variation, may serve as directly influence chronic risk, while also playing moderating roles concerning how individual and relational factors affect imminent or acute risk.

Enhancing Suicide Risk Assessment Methods

Despite a proliferation of suicide risk assessment tools (Lotito and Cook, 2015), recent evidence suggests limited ability to predict future suicidal behavior (Chan et al., 2016; Large et al., 2016). Self-report instruments suffer an identical limitation much of suicide-related theory and risk/protective factor summaries have: a lack of accounting multi-level understanding. For example, the most commonly used self-report tools often assess the frequency and nature of past/present/future ideation and attempts, strongly correlated mental health symptoms (e.g., depression, hopelessness), or protective factors (e.g., reasons for living). While such information is clinically useful to fill in gaps not otherwise captured in interview (Lotito and Cook, 2015), it is still limited in scope.

Several assessment structures exist in the literature, sometimes hinting at the need to address multi-level facets. Such attempts have led to different recommendations for the development of suicide risk assessment tools, but none of these strategies have seen effective implementation yet. For example, suicide risk assessment is often based on lists of symptoms without an integrated perspective (Kral and Sakinofsky, 1994), therefore proposing a model that comprises both background and subjective suicide risk factors. The former are the socio-demographic indices associated with increased risk which are based on different populations and cultures, and are prone to change over time. The latter, background risk factors, can inform the clinician about a patient's general level of risk, while the assessment of individual factors focusses on emotions, cognition, idiosyncratic meanings, general mental state, and experience.

Suicide risk assessment methods can also be based on factors falling into (1) individual, (2) clinical, (3) interpersonal, (4) situational, and (5) demographic categories, thus encompassing some of the SESPM levels suggested herein (Simon, 2011).

In total, the variation in recommended methods suggests that a one-size-fits-all solution to suicide risk estimation is an ill fit. Such methodological complexities might be responsible for the result of the most recent meta-analysis of suicide risk assessment scales, which concluded that there is insufficient evidence to support the use of risk scales and tools in clinical practice due to the rather low positive predictive value (PPV) of the scales, which ranged between only 1.3 and 16.7% (Chan et al., 2016), with 87% false positives, a clinically imprecise, economically intensive and unnecessarily stigmatizing proportion. We argue that the heterogeneity and confusion about suicide risk assessment methods has its primary origin in the lack of a unified and empirically testable theory of suicidal behavior.

We posit that the next meaningful steps in suicide risk assessment tool development may lie in two areas: (1) a psychometrically-validated structured professional judgment (SPJ) of key multi-level risk factors, such as the Screening Tool for Assessing Risk of Suicide (STARS) protocol (Hawgood and De Leo, 2016) (which does not fully account for multi-level influences) and (2) a multi-level suicide risk assessment algorithm. For example, although a potentially time, resource and funding intense project, we recommend development of a suicide risk assessment tool for use by mental and medical health professionals that addresses risk and protective factors across all four layers of the SESPM. Such a new approach may be translated into a SPJ tool, an approach to mental health assessment that provides semi-structured rating forms to be used by trained health professionals. In addition, post-interview, more rigorous interviewer-rated checklists, could help to refine and validate the SPJ tool. Nowadays, online implementation and translation in different languages allows for the development of a globally available instrument for suicide risk assessment within an SESPM model for ongoing refinement. Violence risk literature provides examples of well-validated SPJs accounting for three SEM levels, accounting for empirically-indicated risk and protective factors for interpersonal violence, including individual (e.g., affective stability, substance use), relational (e.g., treatment compliance, personal/social support), and community (e.g., living situation, professional services) level issues (Douglas et al., 2013). Community level risk is further accounted for by the relevance of these factors to the setting of evaluation (e.g., inpatient hospital vs. outpatient clinic). Using the SESPM, it is plausible that a suicide SPJ could be developed by first identifying and testing a lengthy set of risk of factors, exposing the preliminary instrument testing in emergency room, outpatient clinic and inpatient hospital settings. After initial reduction and psychometric evaluation of the instrument, further testing would be required for longitudinal and cross-cultural validation. Likely, a culturally-adapted version of such an instrument would be quite useful.

Alternatively, it may be beneficial to develop a risk assessment algorithm, which is stratified by available knowledge (i.e., different sets of risk factors might be relevant for males vs.

females, adolescents vs. elderly, etc.). As such, a computer-based suicide risk estimation algorithm may be developed in which a clinician can collect comprehensive multi-level patient and situational information, entering the information into a weighted equation. The first step in such effort is mathematical identification of a weighted formula, likely based on a validated multi-level theory of suicide that has to be developed. First approaches in the direction of a suicide risk algorithm have been provided; however, the presented tool was limited to six individual level factors (e.g., age, self-harm history) suitable for depressed persons for the prediction of suicidal ideation only (Liu et al., 2016). With simultaneous systematic review of multi-level risk and protective factors for suicide ideation, attempts, and completed suicide, identification of relative weights for an SESPM-based algorithm based suicide risk assessment tool should be attainable.

Development of Multi-Level Prevention Programming

First and foremost, SESPM-based research is needed to inform best practices for prevention programming. Thus, our recommendations for development of multi-level programming are provided with the caveat that further research and theory development are required. As a starting point, multi-level suicide prevention should address (1) general practitioner education concerning depression and suicide; (2) increased access to care for high-risk groups, and; (3) emphasis on restricting access to lethal means (van der Feltz-Cornelis et al., 2011). While we agree with the importance of these practical recommendations, we further posit that design of an ideal multi-level approach to suicide prevention would possess the following characteristics: (1) incorporation of the risk and protective factors with strongest empirical support relative to the population (e.g., general population vs. high risk psychiatric inpatients); (2) use of prevention strategies at each SESPM level; (3) inclusion of a multi-level program evaluation strategy including data gathered from patients and other stakeholders (e.g., therapists, policy makers, etc.)—patient data would include suicide and self-harm, whereas additional patient and stakeholder information could cover subjective and objective patient-oriented outcomes more generally; (4) grounding in relevant theory to inform mechanisms of change; and (5) presence of prevention efforts using at least primary and secondary prevention techniques where possible.

LIMITATIONS AND CONCLUSIONS

The SESPM perspective holds a number of limitations warranting attention. For example, one shortcoming of the piece is that the present summary of factors did not rise to the level of rigor as formal systematic reviews (nor was it our intention to do so). Moreover, the SESPM is not exhaustive; rather it is intended to provide a flexible framework for additional research and program development moving forward. Another limitation of the present discussion can be seen in failing to conduct meta-regression or analysis; future scholarship may offer

very important confirmation or modification of the framework via such analyses. A notable limitation of our SESPM-based recommendations is the labor, time and resources necessary for education and training efforts; however, we argue that the cumbersome processes involved in many of the SESPM-based recommendations is ultimately worthwhile in the long run because multi-level suicide prevention efforts may save lives and improve quality of life beyond what is currently within the capabilities of public health and clinical mental health fields. Also, while the same limitations concerning time, cost and resources certainly apply to risk assessment tool development recommendations, we believe that the scientific progress in suicide prevention is not a question of experienced clinical rating vs. algorithm building, but both approaches mutually informing each other to create new insight.

We have articulated background, structure and recommendations for a SESPM. The bottom line of our perspective is that we agree with a sentiment that has been expressed in prior suicidology literature: scholars and

practitioners alike need to expand how we think about suicide. The SESPM represents a valuable step in moving from a hyper-focus on individual-level suicide risk prediction toward a comprehensive multi-level perspective on suicide prevention. We welcome further dialogue, research and development moving forward.

AUTHOR CONTRIBUTIONS

Both authors contributed equally to this paper in performing literature search, conceptualizing, drafting, and revising the manuscript.

FUNDING

This conceptual analysis received no further funding from third parties besides provision of the academic environment at Old Dominion University, Norfolk, USA and Medical University of Vienna, Austria.

REFERENCES

- Abramson, L. Y., Alloy, L. B., Hogan, M. E., Whitehouse, W. G., Cornette, M., Akhavan, S., et al. (1998). Suicidality and cognitive vulnerability to depression among college students: a prospective study. *J. Adolesc.* 21, 473–487.
- Anestis, M. D., and Anestis, J. C. (2015). Suicide rates and state laws regulating access and exposure to handguns. *Am. J. Public Health* 105, 2049–2058. doi: 10.2105/AJPH.2015.302753
- Anglemyer, A., Horvath, T., and Rutherford, G. (2014). The accessibility of firearms and risk for suicide and homicide victimization among household members: a systematic review and meta analysis. *Ann. Intern. Med.* 160, 101–110. doi: 10.7326/M13-1301
- Baral, S., Logie, C. H., Gross, A., Wirtz, A. L., and Beyer, C. (2013). Modified social ecological model: a tool to guide the assessment of the risks and risk contexts of HIV epidemics. *BMC Public Health* 17, 482–489. doi: 10.1186/1471-2458-13-482
- Barzilay, S., and Aptek, A. (2014). Psychological models of suicide. *Arch. Suicide Res.* 18, 295–312. doi: 10.1080/13811118.2013.824825
- Bernard, V., Geoffroy, P. A., and Bellivier, F. (2015). Seasons, circadian rhythms, sleep and suicidal behaviors vulnerability. *Encephale* 41, S29–S37. doi: 10.1016/S0013-7006
- Björkensztam, C., Ekseleius, L., Berlin, M., Björkensztam, E., Gerdin, B. (2016). Suicide risk and risk method in patients with personality disorders. *J. Psychiatry Res.* 83, 29–36. doi: 10.1016/j.jpsychires.2016.08.008
- Björkensztam, E., Björkensztam, C., Holm, H., Gerdin, B., and Ekselius, L. (2015). Excess cause-specific mortality in in-patient-treated individuals with personality disorder: 25-year nationwide population-based study. *Brit. J. Psychiatry* 207, 339–345. doi: 10.1192/bj.psy.114.149583
- Bryan, C. J., and Rudd, M. D. (2006). Advances in the assessment of suicide risk. *J. Clin. Psychol.* 62, 185–200. doi: 10.1002/jclp.20222
- Caine, E. D. (2013). Forging an agenda for suicide prevention in the United States. *Am. J. Public Health* 103, 822–829. doi: 10.2105/AJPH.2012.301078
- Calear, A. L., Christensen, H., Freeman, A., Fenton, K., Busby Grant, J., van Spijker, B., et al. (2016). A systematic review of psychosocial suicide prevention interventions for youth. *Eur. J. Child Adolesc. Psychiatry* 25, 467–482. doi: 10.1007/s00787-015-0783-4
- Carrico, A. W., Johnson, M. O., Morin, S. F., Remien, R. H., Charlebois, E. D., Steward, W. T., et al. (2007). Correlates of suicidal ideation among HIV positive persons. *AIDS* 21, 1199–1203. doi: 10.1097/QAD.0b013e3281532c96
- Castro, C. A., and Kintzle, S. (2014). Suicides in the military: the post-modern combat veteran and the Hemingway effect. *Curr. Psychiatry Rep.* 16, 460–468. doi: 10.1007/s11920-014-0460-1
- Centers for Disease Control and Prevention. (2016a). *Increase in Suicide in the United States: NCHS Data Brief No. 241, April. 2016*. Available online at: <https://www.cdc.gov/nchs/products/databriefs/db241.htm> (Accessed January 31, 2017).
- Centers for Disease Control and Prevention. (2016b). *Sexual violence on campus: Strategies for prevention*. 2016. Available online at: <https://www.cdc.gov/violenceprevention/pdf/campussvprevention.pdf> (Accessed January 31, 2017).
- Centers for Disease Control and Prevention. (2017). *The Social-Ecological Model: A Framework for Prevention*. Available online at: <https://www.cdc.gov/violenceprevention/overview/social-ecologicalmodel.html> (Accessed January 13, 2017).
- Chan, M. K., Bhatti, H., Meader, N., Stockton, S., Evans, J., O'Connor, R. C., et al. (2016). Predicting suicide following self-harm: systematic review of risk factors and risk scales. *Br. J. Psychiatry* 209, 277–283. doi: 10.1192/bj.psy.2015.170050
- Chu, C., Klein, K. M., Buchman-Schmitt, J. M., Hom, M. A., Hagan, C. R., and Joiner, T. E. (2015). Routinized assessment of suicide risk in clinical practice: an empirically informed update. *J. Clin. Psychol.* 71, 1186–1200. doi: 10.1002/jclp.22210
- Comtois, K. A., and Linehan, M. M. (2006). Psychosocial treatments for suicidal behavior: a practice-friendly review. *J. Clin. Psychol.* 62, 161–170. doi: 10.1002/jclp.20220
- Cramer, R. J., Bryson, C. N., Eichorst, M. K., Keyes, L. N., and Ridge, B. E. (2017). Conceptualization and pilot testing of a core competency-based training workshop in suicide risk assessment and management. *J. Clin. Psychol.* 73, 233–238. doi: 10.1002/jclp.22329
- Cramer, R. J., Bryson, C. N., Stroud, C. H., and Ridge, B. E. (2016). A pilot test of a graduate course in suicide theory, risk assessment, and management. *Teach Psychol.* 43, 238–242. doi: 10.1177/0098628316649483
- Cramer, R. J., Colbourn, S. L., Gemberling, T. M., Graham, J., and Stroud, C. H. (2015). Substance related coping, HIV-related factors, and mental health among an HIV-positive sexual minority community sample. *AIDS Care* 27, 1063–1068. doi: 10.1080/09540121.2015.1024097
- Cramer, R. J., Johnson, S. M., McLaughlin, J., Rausch, E. M., and Conroy, M. A. (2013). Suicide risk assessment training for psychology doctoral programs: core competencies and framework for training. *Train Edu. Prof. Psychol.* 7, 1–11. doi: 10.1037/a0031836
- Dahlberg, L. L., and Krug, E. G. (2002). “Violence—a global public health problem,” in *World Report on Violence and Health*, eds E. Krug, L. L. Dahlberg, J. A. Mercy, A. B. Zwi, and R. Lozano (Geneva: World Health Organization), 1–56.
- Douglas, K. S., Hart, S. D., Webster, C. D., and Belfrage, H. (2013). *HCR-20V3: Assessing Risk of Violence – User Guide*. Burnaby, BC: Mental Health, Law, and Policy Institute, Simon Fraser University.
- Drapeau, C. W., and McIntosh, J. L. (2014). *U.S.A. Suicide: Official Final Data*. Washington, DC: American Association of Suicidology (Accessed January 31, 2017).

- Duncan, D. T., and Hatzenbuehler, M. (2014). Lesbian, gay, bisexual, and transgender hate crimes and suicidality among a population-based sample of sexual-minority adolescents in Boston. *Am. J. Public Health* 104, 272–278. doi: 10.2105/AJPH.2013.301424
- Durkheim, E. (1897). *Le Suicide*. Paris: Alcan.
- Ellis, T. E., Rufino, K. A., Allen, J. G., Fowler, J. C., and Jobes, D. A. (2015). Impact of a suicide specific intervention within inpatient psychiatric care: the Collaborative Assessment Andmanagement Of Suicide (CAMS). *Sui Life Threatening Behav.* 45, 556–566. doi: 10.1111/slbt.12151
- Erlich, M. D. (2016). Envisioning zero suicide. *Psychiatric Serv.* 67, 255. doi: 10.1176/appi.ps.201500334
- Fowler, J. C. (2012). Suicide risk assessment in clinical practice: pragmatic guidelines for imperfect assessments. *Psychotherapy* 49, 81–90. doi: 10.1037/a0026148
- Franklin, J. C., Ribeiro, J. D., Fox, K. R., Bentley, K. H., Kleiman, E. M., Huang, X., et al. (2017). Risk factors for suicidal thoughts and behaviors: a meta-analysis of 50 years of research. *Psychol. Bull.* 143, 187–232. doi: 10.1037/bul0000084
- Haas, A. P., Eliason, M., Mays, V. M., Mathy, R. M., Cochran, S. D., D'Augelli, A. R., et al. (2011). Suicide and suicide risk in lesbian, gay, bisexual, and transgender populations: review and recommendations. *J. Homosex.* 58, 10–51. doi: 10.1080/00918369
- Hawgood, J., and De Leo, D. (2016). Suicide prediction - a shift in paradigm is needed. *Crisis* 37, 251–255. doi: 10.1027/0227-5910/a000440
- Hegerl, U., Wittmann, M., Arensman, E., Van Audenhove, C., Bouleau, J. H., Van Der Feltz-Cornelis, C., et al. (2008). The European Alliance Against Depression (EAAD): a multifaceted, community-based action programme against depression and suicidality. *World J. Biol. Psychiatry* 9, 51–58. doi: 10.1080/15622970701216681
- Horvath, I. (2016). "Theory building in experimental design and research," in *Experimental Design Research*, ed P. Caish (Switzerland: Springer).
- Hung, E. K., Binder, R. L., Fordwood, S. R., Hall, S. E., Cramer, R. J., and McNeil, D. E. (2012). A method for evaluating competency in assessment and management of suicide risk. *Acad. Psychiatry* 36, 23–28. doi: 10.1176/appi.ap.10110160
- Im, E. O. (2015). The current status of theory evaluation in nursing. *J. Adv. Nurs.* 71, 2268–2278. doi: 10.1111/jan.12698
- Institute of Medicine Committee on Prevention of Mental Disorders. (1994). "Institute of medicine committee on prevention of mental disorders," in *Reducing Risks for Mental Disorders: Frontiers for Preventive Intervention Research*, eds P. J. Mrazek and R. J. Haggerty (Washington, DC: National Academies Press).
- Joiner, T. E. (2005). *Why People Die by Suicide*. Cambridge, MA: Harvard University Press.
- Kral, M. J., and Sakinofsky, I. (1994). Clinical model for suicide risk assessment. *Death Stud.* 18, 311–326.
- Krieger, N. (2016). Living and dying at the crossroads: Racism, embodiment, and why theory is essential to a public health of consequence. *Am. J. Public Health* 106, 832–833. doi: 10.2105/AJPH.2016.303100
- Kumar, S., Quinn, S. C., Kim, K. H., Musa, D., Hilyard, K. M., and Freimuth, V. S. (2009). The social ecological model as a framework for determinants of H1N1 influenza vaccine uptake in the US. *Health Edu. Behav.* 39, 229–243. doi: 10.1177/1090198111415105
- Large, M., Kaneson, M., Myles, N., Myles, H., Gunaratne, P., and Ryan, C. (2016). Meta-analysis of longitudinal cohort studies of suicide risk assessment of psychiatric patients: heterogeneity in results and lack of improvement over time. *PLOS ONE* *Forthcoming* 11:e0156322. doi: 10.1371/journal.pone.0156322
- Lester, D. (1999). The social causes of suicide: a closer look at Durkheim's Le suicide one hundred years later. *Death Stud.* 40, 307–321.
- Linehan, M. M., Goodstein, J. L., Nielsen, S. L., and Chiles, J. A. (1983). Reasons for staying alive when you are thinking of killing yourself: the Reasons for Living Inventory. *J. Consult. Clin. Psychol.* 51, 276–286.
- Liu, Y., Sareen, J., Bolton, J. M., and Wang, J. L. (2016). Development and validation of a risk prediction algorithm for the recurrence of suicidal ideation among general populationwith low mood. *J. Affect Dis.* 193, 11–17. doi: 10.1016/j.jad.2015.12.072
- Lotito, M., and Cook, E. (2015). A review of suicide risk assessment instruments and approaches. *Men Health Clin.* 5, 216–223. doi: 10.9740/mhc.2015.09.216
- Ma, J., Batterham, P. J., Calear, A. L., and Han, J. (2016). A systematic review of the predictions of the interpersonal-psychological theory of suicidal behavior. *Clin. Psychol. Rev.* 46, 34–45. doi: 10.1016/j.cpr.2016.04.008
- Madsen, T., Erlangsen, A., and Nordentoft, M. (2017). Risk estimates and risk factors related to psychiatric inpatient suicide-an overview. *Int. J. Environ. Res. Public Health* 14, 253–265. doi: 10.3390/ijerph14030253
- Mann, J. J., Aptek, A., Bertolote, J., Beautrais, A., Currier, D., Haas, A., et al. (2005). Suicide prevention strategies: a systematic review. *J. Am. Med. Assoc.* 294, 2064–2074. doi: 10.1001/jama.294.16.2064
- McCarten, J. M., Hoffmire, C. A., and Bossarte, R. M. (2001). Changes in overall and firearm veteran suicide rates by gender,—2010. *Am. J. Prev. Med.* 48, 360–364. doi: 10.1016/j.amepre.2014.10.013
- McCormack, L., Thomas, V., Lewis, M. A., and Rudd, R. (2017). Improving low health literacy and patient engagement: a social ecological approach. *Pat. Edu. Counsel.* 100, 8–13. doi: 10.1016/j.pec.2016.07.007
- O'Connor, R. C. (2011). The integrated motivational-volitional model of suicidal behavior. *Crisis* 32, 295–298. doi: 10.1027/0227-5910/a000120
- Prestwich, A., Webb, T. L., and Conner, M. (2015). Using theory to develop and test interventions to promote changes in health behaviour: evidence, issues, and recommendations. *Curr. Opin. Psychol.* 5, 1–15. doi: 10.1016/j.copsyc.2015.02.011
- Rubens, M., and Shehadeh, N. (2014). Gun violence in United States: in search for a solution. *Front. Public Health* 3:17. doi: 10.3389/fpubh.2014.00017
- Rudd, M. D., Cukrowicz, K. C., and Bryan, C. J. (2008). Core competencies in suicide risk assessment and management: implications for supervision. *Train. Edu. Prof. Psychol.* 2, 219–228. doi: 10.1037/1931-3918.2.4.219
- Serafini, G., Pompili, M., Innamorati, M., Rihmer, Z., Sher, L., and Girardi, P. (2012). Can cannabis increase suicide risk in psychosis? a critical review. *Curr. Pharm. Des.* 18, 5156–5187.
- Shneidman, E. S. (1981). A psychological theory of suicide. *Suicide Life Threat. Behav.* 11, 221–231.
- Silverman, M. M., and De Leo, D. (2016). Why there is a need for an international nomenclatureandclassification system for suicide. *Crisis* 37, 83–87. doi: 10.1027/0227-5910/a000419
- Simon, R. I. (2011). *Preventing Patient Suicide: Clinical Assessment and Management*. Washington, DC: American Psychiatric Publishing.
- Skegg, K. (2005). Self-harm. *Lancet* 366, 1471–1483. doi: 10.1016/S0140-6736(05)67600-3
- Stanley, I. H., Horn, M. A., Rogers, M. L., Anestis, M. D., and Joiner, T. E. (2016). Discussing firearm ownership and access as part of suicide risk assessment and prevention: means safety versus means restriction. *Arch. Suicide Res.* 13, 1–17. doi: 10.1080/13811118.2016.1175395
- Teo, A. R., Andrea, S. B., Sakakibara, R., Motohara, S., Matthieu, M. M., and Fetters, M. D. (2016). Brief gatekeeper training for suicide prevention in an ethnic minority population: a controlled intervention. *BMC Psychiatry* 16:211. doi: 10.1186/s12888-016-0924-4
- United States Surgeon General's Office. (2012). *National Strategy for Suicide Prevention: Goals and Objectives for Action*. Washington, DC: Department of Health and Human Services.
- van der Feltz-Cornelis, C. M., Sarchiapone, M., Postuvan, V., Volker, D., Roskar, S., Grum, A. T., et al. (2011). Best practice elements ofmultilevel suicide prevention strategies: a review of systematic reviews. *Crisis* 32, 319–333. doi: 10.1027/0227-5910/a000109
- Van Orden, K. A., Witte, T. K., Cukrowicz, K. C., Braithwaite, S. R., Selby, E. A., and Joiner, T. E. (2010). The interpersonal theory of suicide. *Psychol. Rev.* 117, 575–600. doi: 10.1037/a0018697
- Zalsman, G., Hawton, K., Wasserman, D., van Heeringen, K., Arensman, E., Sarchiapone, M., et al. (2016). Suicide prevention strategies revisited: 10year systematic review. *Lancet Psychol.* 3, 646–659. doi: 10.1016/S2215-0366(16)30030-X

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Cramer and Kapusta. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.