

5-2017

## Target and Double Spin Asymmetries of Deeply Virtual $\pi^0$ Production With a Longitudinally Polarized Proton Target and CLAS

A. Kim

H. Avakian

V. Burkert

K. Joo

W. Kim

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.odu.edu/physics\\_fac\\_pubs](https://digitalcommons.odu.edu/physics_fac_pubs)



Part of the [Astrophysics and Astronomy Commons](#), and the [Nuclear Commons](#)

---

### Original Publication Citation

Kim, A., Avakian, H., Burkert, V., Joo, K., Kim, W., Adhikari, K. P., . . . Zhang, J. (2017). Target and double spin asymmetries of deeply virtual  $\pi^0$  production with a longitudinally polarized proton target and CLAS. *Physics Letters B*, 768, 168-173. doi:10.1016/j.physletb.2017.02.032

This Article is brought to you for free and open access by the Physics at ODU Digital Commons. It has been accepted for inclusion in Physics Faculty Publications by an authorized administrator of ODU Digital Commons. For more information, please contact [digitalcommons@odu.edu](mailto:digitalcommons@odu.edu).

---

**Authors**

A. Kim, H. Avakian, V. Burkert, K. Joo, W. Kim, K. P. Adhikari, Z. Akbar, S. Anefalos Pereira, R. A. Badui, M. Battaglieri, S. Bültmann, A. Klein, S E. Kuhn, Y. Prok, B. Torayev, L. B. Weinstein, and J. Zhang



## Target and double spin asymmetries of deeply virtual $\pi^0$ production with a longitudinally polarized proton target and CLAS



A. Kim<sup>b,a,\*</sup>, H. Avakian<sup>c</sup>, V. Burkert<sup>c</sup>, K. Joo<sup>a</sup>, W. Kim<sup>b</sup>, K.P. Adhikari<sup>e,d</sup>, Z. Akbar<sup>f</sup>, S. Anefalos Pereira<sup>g</sup>, R.A. Badui<sup>h</sup>, M. Battaglieri<sup>i</sup>, V. Baturine<sup>c</sup>, I. Bedlinskiy<sup>j</sup>, A.S. Biselli<sup>k</sup>, S. Boiarinov<sup>c</sup>, P. Bosted<sup>l,c</sup>, W.J. Briscoe<sup>m</sup>, W.K. Brooks<sup>n</sup>, S. Bültmann<sup>e</sup>, T. Cao<sup>o</sup>, D.S. Carman<sup>c</sup>, A. Celentano<sup>i</sup>, S. Chandavar<sup>p</sup>, G. Charles<sup>q</sup>, T. Chetry<sup>p</sup>, L. Colaneri<sup>s,r</sup>, P.L. Cole<sup>t</sup>, N. Compton<sup>p</sup>, M. Contalbrigo<sup>u</sup>, O. Cortes<sup>t</sup>, V. Crede<sup>f</sup>, A. D'Angelo<sup>s,r</sup>, N. Dashyan<sup>v</sup>, R. De Vita<sup>i</sup>, E. De Sanctis<sup>g</sup>, C. Djalali<sup>o</sup>, H. Egiyan<sup>w,c</sup>, A. El Alaoui<sup>y,x,n</sup>, L. El Fassi<sup>x,d</sup>, P. Eugenio<sup>f</sup>, G. Fedotov<sup>z,o</sup>, R. Fersch<sup>l</sup>, A. Filippi<sup>aa</sup>, J.A. Fleming<sup>ab</sup>, A. Fradi<sup>q</sup>, M. Garcia<sup>ac</sup>, Y. Ghandilyan<sup>v</sup>, G.P. Gilfoyle<sup>ad</sup>, K.L. Giovanetti<sup>ae</sup>, F.X. Girod<sup>ac,c</sup>, W. Gohn<sup>a</sup>, E. Golovatch<sup>z</sup>, R.W. Gothe<sup>o</sup>, K.A. Griffioen<sup>l</sup>, L. Guo<sup>c,h</sup>, K. Hafidi<sup>x</sup>, C. Hanretty<sup>af</sup>, M. Hattawy<sup>q</sup>, D. Heddle<sup>c,ag</sup>, K. Hicks<sup>p</sup>, M. Holtrop<sup>w</sup>, Y. Ilieva<sup>m,o</sup>, D.G. Ireland<sup>ah</sup>, B.S. Ishkhanov<sup>z</sup>, D. Jenkins<sup>ai</sup>, H. Jiang<sup>o</sup>, H.S. Jo<sup>q</sup>, S. Joosten<sup>aj</sup>, D. Keller<sup>p,af</sup>, G. Khachatryan<sup>v</sup>, M. Khandaker<sup>ak,t</sup>, A. Klein<sup>e</sup>, F.J. Klein<sup>al</sup>, V. Kubarovsky<sup>am,c</sup>, S.E. Kuhn<sup>e</sup>, S.V. Kuleshov<sup>j,n</sup>, L. Lanza<sup>r</sup>, P. Lenisa<sup>u</sup>, H.Y. Lu<sup>o</sup>, I.J.D. MacGregor<sup>ah</sup>, N. Markov<sup>a</sup>, P. Mattione<sup>an</sup>, M.E. McCracken<sup>an</sup>, B. McKinnon<sup>ah</sup>, V. Mokeev<sup>z,c</sup>, A. Movsisyan<sup>u</sup>, E. Munevar<sup>c</sup>, P. Nadel-Turonski<sup>al,c</sup>, L.A. Net<sup>o</sup>, S. Niccolai<sup>q</sup>, M. Osipenko<sup>i</sup>, A.I. Ostrovidov<sup>f</sup>, M. Paolone<sup>aj</sup>, K. Park<sup>o,c</sup>, E. Pasyuk<sup>ao,c</sup>, W. Phelps<sup>h</sup>, S. Pisano<sup>q,g</sup>, O. Pogorelko<sup>j</sup>, J.W. Price<sup>ap</sup>, Y. Prok<sup>af,e</sup>, M. Ripani<sup>i</sup>, A. Rizzo<sup>s,r</sup>, G. Rosner<sup>ah</sup>, P. Rossi<sup>g,c</sup>, P. Roy<sup>f</sup>, C. Salgado<sup>ak</sup>, R.A. Schumacher<sup>an</sup>, E. Seder<sup>a</sup>, Y.G. Sharabian<sup>c</sup>, Iu. Skorodumina<sup>z,o</sup>, G.D. Smith<sup>ab</sup>, D. Sokhan<sup>ah</sup>, N. Sparveris<sup>aj</sup>, S. Stepanyan<sup>c</sup>, P. Stoler<sup>am</sup>, I.I. Strakovsky<sup>m</sup>, S. Strauch<sup>o</sup>, V. Sytnik<sup>n</sup>, M. Taiuti<sup>aq</sup>, B. Torayev<sup>e</sup>, M. Ungaro<sup>a,c</sup>, H. Voskanyan<sup>v</sup>, E. Voutier<sup>q</sup>, D.P. Watts<sup>ab</sup>, X. Wei<sup>c</sup>, L.B. Weinstein<sup>e</sup>, N. Zachariou<sup>o</sup>, L. Zana<sup>w,ab</sup>, J. Zhang<sup>e,c</sup>

<sup>a</sup> University of Connecticut, Storrs, CT 06269, United States

<sup>b</sup> Kyungpook National University, Daegu 41566, Republic of Korea

<sup>c</sup> Thomas Jefferson National Accelerator Facility, Newport News, VA 23606, United States

<sup>d</sup> Mississippi State University, Mississippi State, MS 39762-5167, United States

<sup>e</sup> Old Dominion University, Norfolk, VA 23529, United States

<sup>f</sup> Florida State University, Tallahassee, FL 32306, United States

<sup>g</sup> INFN, Laboratori Nazionali di Frascati, 00044 Frascati, Italy

<sup>h</sup> Florida International University, Miami, FL 33199, United States

<sup>i</sup> INFN, Sezione di Genova, 16146 Genova, Italy

<sup>j</sup> Institute of Theoretical and Experimental Physics, Moscow, 117259, Russia

<sup>k</sup> Fairfield University, Fairfield, CT 06824, United States

<sup>l</sup> College of William and Mary, Williamsburg, VA 23187-8795, United States

<sup>m</sup> The George Washington University, Washington, DC 20052, United States

<sup>n</sup> Universidad Técnica Federico Santa María, Casilla 110-V Valparaíso, Chile

<sup>o</sup> University of South Carolina, Columbia, SC 29208, United States

<sup>p</sup> Ohio University, Athens, OH 45701, United States

<sup>q</sup> Institut de Physique Nucléaire, CNRS/IN2P3 and Université Paris Sud, Orsay, France

<sup>r</sup> INFN, Sezione di Roma Tor Vergata, 00133 Rome, Italy

<sup>s</sup> Università di Roma Tor Vergata, 00133 Rome, Italy

<sup>t</sup> Idaho State University, Pocatello, ID 83209, United States

\* Corresponding author.

E-mail address: [kenjo@jlab.org](mailto:kenjo@jlab.org) (A. Kim).

- <sup>u</sup> INFN, Sezione di Ferrara, 44100 Ferrara, Italy  
<sup>v</sup> Yerevan Physics Institute, 375036 Yerevan, Armenia  
<sup>w</sup> University of New Hampshire, Durham, NH 03824-3568, United States  
<sup>x</sup> Argonne National Laboratory, Argonne, IL 60439, United States  
<sup>y</sup> LPSC, Université Grenoble-Alpes, CNRS/IN2P3, Grenoble, France  
<sup>z</sup> Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, 119234 Moscow, Russia  
<sup>aa</sup> INFN, Sezione di Torino, 10125 Torino, Italy  
<sup>ab</sup> Edinburgh University, Edinburgh EH9 3JZ, United Kingdom  
<sup>ac</sup> CEA, Centre de Saclay, Irfu/Service de Physique Nucléaire, 91191 Gif-sur-Yvette, France  
<sup>ad</sup> University of Richmond, Richmond, VA 23173, United States  
<sup>ae</sup> James Madison University, Harrisonburg, VA 22807, United States  
<sup>af</sup> University of Virginia, Charlottesville, VA 22901, United States  
<sup>ag</sup> Christopher Newport University, Newport News, VA 23606, United States  
<sup>ah</sup> University of Glasgow, Glasgow G12 8QQ, United Kingdom  
<sup>ai</sup> Virginia Tech, Blacksburg, VA 24061-0435, United States  
<sup>aj</sup> Temple University, Philadelphia 19122, United States  
<sup>ak</sup> Norfolk State University, Norfolk, VA 23504, United States  
<sup>al</sup> Catholic University of America, Washington, DC 20064, United States  
<sup>am</sup> Rensselaer Polytechnic Institute, Troy, NY 12180-3590, United States  
<sup>an</sup> Carnegie Mellon University, Pittsburgh, PA 15213, United States  
<sup>ao</sup> Arizona State University, Tempe, AZ 85287-1504, United States  
<sup>ap</sup> California State University, Dominguez Hills, Carson, CA 90747, United States  
<sup>aq</sup> Università di Genova, 16146 Genova, Italy

## ARTICLE INFO

## Article history:

Received 27 September 2016  
 Received in revised form 20 January 2017  
 Accepted 14 February 2017  
 Available online 22 February 2017  
 Editor: V. Metag

## Keywords:

Nucleon structure  
 CLAS collaboration  
 Deeply Virtual Meson Production  
 Chiral-odd generalized parton distributions  
 Longitudinally polarized target

## ABSTRACT

The target and double spin asymmetries of the exclusive pseudoscalar channel  $\bar{e}p \rightarrow ep\pi^0$  were measured for the first time in the deep-inelastic regime using a longitudinally polarized 5.9 GeV electron beam and a longitudinally polarized proton target at Jefferson Lab with the CEBAF Large Acceptance Spectrometer (CLAS). The data were collected over a large kinematic phase space and divided into 110 four-dimensional bins of  $Q^2$ ,  $x_B$ ,  $-t$  and  $\phi$ . Large values of asymmetry moments clearly indicate a substantial contribution to the polarized structure functions from transverse virtual photon amplitudes. The interpretation of experimental data in terms of generalized parton distributions (GPDs) provides the first insight on the chiral-odd GPDs  $\tilde{H}_T$  and  $\tilde{E}_T$ , and complement previous measurements of unpolarized structure functions sensitive to the GPDs  $H_T$  and  $\bar{E}_T$ . These data provide a crucial input for parametrizations of essentially unknown chiral-odd GPDs and will strongly influence existing theoretical calculations based on the handbag formalism.

© 2017 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP<sup>3</sup>.

The introduction of generalized parton distributions (GPDs) [1–3] defines a new important and far-ranging theoretical framework that allows for the description of the angular momentum components of quarks and gluons in the proton in terms of density distributions in both longitudinal momentum fraction and transverse spatial degrees of freedom. They provide information on the orbital motion of partons, rendering a three dimensional view of hadron structure [4,5]. Therefore, GPDs are the universal functions that offer an unprecedented opportunity to investigate the nucleon internal structure and provide insight into the hadron at the quark–gluon level.

At leading twist there are eight GPDs [5] for each quark flavor  $q$ : four correspond to parton helicity conserving (chiral-even) processes, denoted as  $H^q$ ,  $E^q$ ,  $\tilde{H}^q$ ,  $\tilde{E}^q$ , and the remaining four,  $H_T^q$ ,  $E_T^q$ ,  $\tilde{H}_T^q$ ,  $\tilde{E}_T^q$ , correspond to parton helicity-flip (chiral-odd) processes [6,7]. The conventional  $\bar{E}_T = 2\tilde{H}_T + E_T$  will be used as well hereafter. These GPDs can be accessed from the hard exclusive processes such as deeply virtual exclusive photon and meson electroproduction. Deeply virtual pseudoscalar meson electroproduction is sensitive to the chiral-odd GPDs which are less-known than their chiral-even counterparts, because they are not accessible in deeply virtual Compton scattering and, generally, their contributions are suppressed [8,9]. However, their knowledge opens a new avenue to study the partonic structure of the nucleon. In particular,  $H_T$  becomes the quark transversity structure function,  $h_1$ , in the forward limit, and it also integrates into the still unknown tensor charge; the  $\tilde{E}_T$  is related to the Boer–Mulders function with

its first moment interpreted as the proton's transverse anomalous magnetic moment [10,8].

The unpolarized cross section measurements [11,12] presented the first evidence that deeply virtual  $\pi^0$  electroproduction can be interpreted in terms of the chiral-odd GPDs. The inclusion of twist-3 components calculated using the chiral-odd GPD parametrizations leads to sizable transverse virtual photon amplitudes and brings theoretical calculations into agreement with experimental data. However, while the measurements of the unpolarized structure functions and beam spin asymmetries for deeply exclusive  $\pi^0$  production have been obtained by the CLAS [11–13] and Hall A Collaborations [14,15], there are no experimental data available on a longitudinally polarized target. This work presents the first extraction of target and double spin asymmetries for deeply virtual  $\pi^0$  production.

The experimental observables in  $\pi^0$  electroproduction are connected to the combinations of the different convolutions, defined as [9]

$$\langle F \rangle = \int dx \mathcal{H}(x, \xi, Q^2) F(x, \xi, t) \quad (1)$$

where  $F$  represents generic GPD,  $\mathcal{H}$  is a hard subprocess amplitude,  $t$  is a momentum transfer to the nucleon and  $\xi$  is a longitudinal momentum fraction transfer. The unpolarized cross sections contain the combinations of different generalized form factors and, similarly to DVCS, require the polarized structure functions to perform the separation of individual convolutions. The interpreta-

tion of single spin asymmetries, however, is more complicated in comparison with unpolarized cross-sections. Firstly, the measured asymmetries are the ratios of polarized structure functions and the unpolarized cross section, so the knowledge of unpolarized structure functions is necessary to isolate the polarized contribution. Secondly, the polarized structure functions are calculated as products of chiral-even and chiral-odd convolutions, complicating the separation of the different contributions. On the other hand, the double spin asymmetry is well suited for the extraction of chiral-odd GPDs, namely  $H_T$ , allowing clean separation of  $\tilde{E}_T$  and  $H_T$  in conjunction with unpolarized target measurements.

Spin asymmetries are defined as a ratio of the difference over the sum of cross sections for opposite helicity configurations and they can be expressed as:


$$A_{UL} = \frac{A_{UL}^{\sin \phi} \sin \phi + A_{UL}^{\sin 2\phi} \sin 2\phi}{1 + A_{UU}^{\cos \phi} \cos \phi + A_{UU}^{\cos 2\phi} \cos 2\phi}, \quad (2)$$

$$A_{LL} = \frac{A_{LL}^{\text{const}} + A_{LL}^{\cos \phi} \cos \phi}{1 + A_{UU}^{\cos \phi} \cos \phi + A_{UU}^{\cos 2\phi} \cos 2\phi}, \quad (3)$$

where the first index  $U$  ( $L$ ) stands for unpolarized (longitudinally polarized) beam, the second index  $U$  ( $L$ ) for the target polarization and  $\phi$  is azimuthal angle between the lepton and hadron scattering planes.  $A_{UU}^{\cos \phi}$  and  $A_{UU}^{\cos 2\phi}$  are connected to the unpolarized structure functions, common for the beam, target and double spin asymmetries, and  $A_{UL}^{\sin \phi}$ ,  $A_{UL}^{\sin 2\phi}$ ,  $A_{LL}^{\text{const}}$ ,  $A_{LL}^{\cos \phi}$  are connected to the polarized structure functions.

We present the first measurements of single target and double spin asymmetries for deeply virtual  $\pi^0$  electroproduction off the longitudinally polarized protons over a large phase space. The experiment was carried out in 2009 in Hall B at Jefferson Lab, using CLAS [16], a longitudinally polarized electron beam with average energy of 5.9 GeV and a longitudinally polarized solid ammonia target [17]. The target system, based on a 5 T superconducting magnet and a 1 K  $^4\text{He}$  refrigerating bath, was constructed to polarize protons in paramagnetically doped  $^{14}\text{NH}_3$  along the beam direction via the dynamic nuclear polarization method. Simultaneously the target's magnetic field serves as an effective shield from Møller electrons by focusing them towards the beam line, while allowing detection of photons from  $4^\circ$  and maintaining the minimum permitted angle for electrons and protons at  $21^\circ$ . The beam polarization was frequently monitored in Møller runs, via the measurement of the asymmetry of elastic electron–electron scattering. The target polarization was continuously monitored by a Nuclear Magnetic Resonance (NMR) system. In addition, data were collected using a  $^{12}\text{C}$  target for the purpose of unpolarized nuclear background studies.

The large acceptance of CLAS allowed simultaneous detection of all four final-state particles of the  $ep \rightarrow ep\pi^0$  and  $\pi^0 \rightarrow \gamma\gamma$  reactions. The scattered electron was identified by a reconstructed track in the drift chambers and matching it in time with signals in the same CLAS sector of the electromagnetic calorimeter (EC) and the Čerenkov counter. The cuts on EC energy deposition effectively suppressed the background from negative pions. The proton was identified as a positively charged particle track in the magnetic field of the superconducting toroidal magnet, passing through the drift chambers with the correct time-of-flight information from the scintillation counters. The neutral pion decay photons were detected in the EC and the inner calorimeter (IC), which was installed downstream of the target and dedicated to the detection of the photons emitted in the forward direction. The photons were detected in the angular range between  $4^\circ$  to  $17^\circ$  in the IC and for angles greater than  $21^\circ$  in the EC.




**Fig. 1.** (Color online.) Distributions of missing mass squared of the  $(ep)$  system for the reaction  $ep \rightarrow ep\pi^0$  before (a) and after (b) the exclusivity cuts are applied (except  $MM_X^2(ep)$  cut itself). The  $^{12}\text{C}$  data (points) are normalized to the  $^{14}\text{NH}_3$  data (line).

After the identification of the four particles, the exclusive events from the  $ep \rightarrow ep\pi^0$  reaction were selected. With the 4-momenta reconstructed for all final-state particles, the event kinematics is fully known, and energy and momentum conservation can be used to develop the *exclusivity cuts*. These constraints allow for the rejection of events from unpolarized nuclear background, different channels (e.g.  $\eta$ ,  $\rho$  or  $\omega$  meson production) and reactions with an additional particle present but undetected.

Three photon-detection topologies exist: (i) both photons detected in the IC, (ii) both photons in the EC and (iii) one photon in the IC and another in the EC. The experimental resolutions of the kinematic quantities for these topologies were different due to the superior IC resolution, and thus the *exclusivity cuts* were determined independently for each case. To ensure the exclusivity of  $\pi^0$  meson production we used the  $3\sigma$  cuts extracted from the Gaussian fits of the following four variables: the missing mass squared  $M_X^2(ep)$  of the  $(epX)$  system, the invariant mass of the two photons  $M_{\gamma\gamma}$ , the missing energy  $E_{ep\gamma\gamma}$  of the  $(ep\gamma\gamma)$  system and the angle  $\theta_{\pi^0 X}$  between the measured and the kinematically reconstructed  $\pi^0$  meson in the  $ep \rightarrow epX$  system.

Fig. 1 illustrates the effect of the exclusivity cuts on the missing mass of the  $ep$  system in  $ep \rightarrow epX$ . The contaminations from different meson production and nuclear background are greatly reduced, however even after the application of all exclusivity cuts, the events from nuclear background are still present. This remaining contamination from  $^{14}\text{N}$  was estimated using the data from carbon runs. The data from  $^{12}\text{C}$  target were normalized to  $^{14}\text{NH}_3$  by the total charge and corrected for the different areal densities of the target materials. The variations of the dilution factor with kinematics were too small to parameterize accurately, so constant dilution factors (0.9, 0.94, 0.91) were applied for three topologies (EC-EC, IC-IC and EC-IC). The contribution from unpolarized nuclear protons was less than 10% for all topologies. The small amount of background from accidental photons is visible in Fig. 2 under the invariant mass spectrum of the two photons  $M_{\gamma\gamma}$  for the three detection topologies. It was subtracted using the data in the sidebands  $(-4.5\sigma, -3\sigma) \cup (3\sigma, 4.5\sigma)$  of the  $M_{\gamma\gamma}$  distributions independently for each kinematic bin and helicity configuration. The latter is particularly important because it takes into account any polarization dependent background.

To ensure that the selected events were from the deep-inelastic regime, the kinematic cuts  $Q^2 > 1$  (GeV/c) $^2$  and  $W > 2$  GeV/c $^2$  were applied.  $W = \sqrt{(p+q)^2}$  is the  $\gamma^*p$  invariant mass, where  $q$  and  $p$  are the four-momenta of the virtual photon and nucleon, and  $Q^2 = -q^2$ . Then the data were divided into 110 four-dimensional kinematical bins for each of the 4 possible beam/tar-



**Fig. 2.** (Color online.) Distributions of invariant mass of the two-photon system for the three different detector configurations: IC-IC, EC-EC and EC-IC from left to right. The cyan areas represent the cuts used for event selection. The last two subfigures show the factors used to scale the number of events in the histograms to the first subfigure.

get helicity configurations. The target and double spin asymmetries were calculated for each kinematic bin as follows:

$$A_{UL} = \frac{\sum_i (n_i^{+-} + n_i^{-+}) - \sum_i (n_i^{++} + n_i^{--})}{P_t^- \sum_i f_i (n_i^{++} + n_i^{--}) + P_t^+ \sum_i f_i (n_i^{+-} + n_i^{-+})}, \quad (4)$$

$$A_{LL} = \frac{1}{P_b P_t^-} \frac{\sum_i (n_i^{+-} + n_i^{-+}) - \sum_i (n_i^{++} + n_i^{--})}{\sum_i f_i (n_i^{++} + n_i^{--}) + P_t^+ \sum_i f_i (n_i^{+-} + n_i^{-+})}, \quad (5)$$

where  $n^{\pm\pm}$  are the numbers of counts for each beam/target helicity configuration, normalized by the corresponding beam charge. The  $i$  index refers to the photon detection topology,  $f_i$  is the corresponding dilution factor, and  $P_t^\pm$  are the average values for the positive/negative target polarizations.

The average target polarizations  $P_t^\pm$  ( $P_t^+ \simeq 80\%$ ,  $P_t^- \simeq 74\%$ ) were extracted by dividing the product of beam and target polarizations  $P_b P_t$  by the beam polarization  $P_b$ . The former was determined by measuring the well-known spin asymmetry in elastic  $ep$  scattering [18,19]. The latter was measured a few times during the experiment using the Møller polarimeter in Hall B. The average value was determined to be  $84\% \pm 2\%$  using the beam polarization measurements weighted by all the events.

The extraction of target and double spin asymmetries for the exclusive  $ep \rightarrow ep\pi^0$  reaction includes several sources that could induce systematic uncertainties. The main source was the event selection procedure. The exclusivity cuts were modified from  $2.5\sigma$  to  $3.5\sigma$ , and the spin asymmetries were re-analyzed for every cut alteration. The corresponding variations of asymmetries were determined to be 4.4% on average. The sideband background subtraction procedure accounted for a systematic uncertainty of 1%. To avoid systematic uncertainty associated with NMR measurements the product of beam and target polarizations was extracted from the exclusive  $ep$  elastic scattering. Therefore, combined with the dilution factor, the uncertainties of the beam and the target polarizations lead to an overall normalization uncertainty of 3% for double spin asymmetry and 5% for target spin asymmetry. The acceptance and binning effects were studied through careful Monte-Carlo simulation, and both effects were found to be negligible. The individual uncertainties for each kinematic bin were added in quadrature, and their values, with an average of 4.5%, were found to be smaller than statistical uncertainties for the most of bins.

The target and double spin asymmetries for exclusive  $\pi^0$  production were measured over a wide kinematic range with  $1 < Q^2 < 5$  (GeV/c)<sup>2</sup>,  $0.1 < x_B < 0.6$ , and  $0 < -t < 2$  (GeV/c)<sup>2</sup>, where

$x_B = \frac{Q^2}{2pq}$  is the Bjorken variable,  $t = (p - p')^2$  is the momentum transfer to the nucleon, and  $p$  and  $p'$  are the initial and final four-momenta of the nucleon. The data were divided into two bins in the  $(Q^2, x_B)$  space, five bins in  $-t$  and eleven  $\phi$  bins with the measured asymmetries shown as a function of  $\phi$  in Fig. 3. The measurements exhibit strong azimuthal dependence for the target spin asymmetries with significant amplitudes of the  $\sin\phi$  moments and a large constant term for the double spin asymmetries.


The measured beam, target and double spin asymmetries were fitted simultaneously using six free parameters:  $A_{LU}^{\sin\phi}$ ,  $A_{UL}^{\sin\phi}$ ,  $A_{UL}^{\sin 2\phi}$ ,  $A_{LL}^{\cos\phi}$ ,  $A_{LL}^{\cos 2\phi}$  and  $A_{UU}^{\cos 2\phi}$  according to the Eqs. (2) and (3) to describe their azimuthal dependence. Simultaneous fit exploits the fact that the three asymmetries have the same denominator and constrains the common terms to be the same for the three different observables. The beam spin asymmetry was extracted in addition to the target and double spin asymmetries. This observable is important in the simultaneous fit to better constrain the unpolarized term  $A_{UU}^{\cos 2\phi}$  in the denominator common for the different polarized observables. Both  $A_{LU}^{\sin\phi}$  and  $A_{UU}^{\cos 2\phi}$  are by-products of the measurement and much better constrained from previous experiments with an unpolarized hydrogen target. Due to the limited statistics, the term  $A_{UU}^{\cos\phi} = \sqrt{2\epsilon(1+\epsilon)}\sigma_{LT}/\sigma_0$  was fixed using the structure functions reported by CLAS in [11]. The correlations of numerator terms with  $A_{UU}^{\cos\phi}$  and  $A_{UU}^{\cos 2\phi}$  were studied and found to be small for all asymmetry moments except  $A_{LL}^{\cos\phi}$  term. To verify the stability of moments extraction the fit was performed with and without  $A_{UU}^{\cos\phi}$  term. Only  $A_{LL}^{\cos\phi}$  exhibits positive correlation with unpolarized terms as evident from large systematic uncertainties on Fig. 4 driven by uncertainties on the denominator parameters.

In Fig. 4 the measured asymmetry moments for  $\pi^0$  electro-production are plotted as a function of  $-t$  in each  $(Q^2, x_B)$  bin, where each kinematic value is calculated as event weighted average. The theoretical predictions from two GPD-based approaches, GK [20] and GGL [21], are also included. They both calculate the contributions from the transverse virtual photon amplitudes using chiral-odd GPDs with  $-t$  dependence, incorporated from Regge phenomenology, but differ in the GPD parametrization methods. GGL provides the chiral-odd GPD parametrization via linear relations to chiral-even GPDs under parity and charge conjugation symmetries in a Regge-ized diquark model. This approach allows the model to overcome the issue that very few constraints on chiral-odd GPDs exist, while chiral-even GPDs can be relatively well-constrained using deep inelastic scattering, nucleon form-factor and DVCS measurements. In the GK model, chiral-odd GPDs are constructed from the double distributions and constrained using the latest results from lattice QCD [22] and transversity parton distribution functions [23] with the emphasis on  $H_T$  and  $\bar{E}_T$ , while the contributions from the other chiral-odd GPDs are considered negligible.

Within the approximation of GK model [20] the  $A_{LL}^{\cos\phi}$  component is defined as:

$$A_{LL}^{\cos\phi} \sigma_0 = \sqrt{1 - \epsilon^2} \frac{4\pi\alpha}{k} \frac{\mu_\pi^2}{Q^4} (1 - \xi^2) |\langle H_T \rangle|^2 \quad (6)$$

A rather straightforward interpretation is obtained for  $A_{LL}^{\cos\phi}$  component, which contains contributions from the chiral-odd GPDs only. It is expected to be determined by  $\langle H_T \rangle$  with a negligible contribution from  $\langle \bar{H}_T \rangle$ . The data do not decrease near threshold which is indicative of  $\langle H_T \rangle$  dominance, especially at low  $-t$  region, and both models display a rise of  $A_{LL}^{\cos\phi}$  in the same region since they rely on  $\langle H_T \rangle$  to describe it. Furthermore, the extraction of



**Fig. 3.** Target and double spin asymmetries for deep exclusive  $\pi^0$  production plotted as a function of  $\phi$  for each kinematic bin in  $(Q^2, x_B)$  space and  $-t$  range. The curves are simultaneous fit results described in the text. The shaded bands represent the overall systematic uncertainties. The latter exhibits strong variation with  $\phi$  in certain kinematic bins due to unseparable statistical fluctuations inflating the estimate of the systematic errors. The bins with low statistics are particularly affected by them.


$A_{LL}^{const}$  term is very stable due to the absence of  $\phi$  dependence and, therefore, provides a reliable experimental observable to constrain GPD  $H_T$ . Additionally, and perhaps more importantly, the double spin asymmetry measurements provide an independent test of the existing GPDs models probing the underlying assumption of  $H_T$  and  $\bar{E}_T$  dominance. The first approach to combine  $A_{LL}^{const}$  term with  $\sigma_0$  and  $\sigma_{TT}$  previously measured by CLAS was conducted in [24]. The constant term was calculated within GK model framework using the  $H_T$  and  $\bar{E}_T$  convolutions extracted from unpolarized cross section measurements [11,12] and compared to the measurements presented in this work. The agreement within error bars demonstrates the major contribution from  $\langle H_T \rangle$ .

The large magnitudes of  $A_{LL}^{\cos\phi}$  and  $A_{UL}^{\sin\phi}$  components suggest sizable contributions from the chiral-odd GPDs through the interference of transverse and longitudinal virtual photons amplitudes. The calculations of these contributions are complicated, largely due to the unknown phases between interfering terms [20,21]. The  $A_{UL}^{\sin\phi}$  term exhibits a relatively flat  $-t$  dependence similar to the observed dependence of the beam-spin asymmetry [13], but with a factor of three larger magnitude. Note that both terms are dominated by  $\langle \bar{H} \rangle^* \langle E_T \rangle$ , but the target spin asymmetry is also enhanced by  $\langle \bar{H} \rangle^* \langle \bar{H}_T \rangle$  and the beam spin asymmetry is reduced by a kinematic factor. The  $A_{UL}^{\sin 2\phi}$  component is determined by chiral-odd GPDs, namely  $E_T$  and  $\bar{H}_T$ , providing the means to disentangle these two GPDs and improve the parameterization of their combination  $\bar{E}_T$  [21].

In conclusion, for the first time target and double spin asymmetries from deeply virtual  $\pi^0$  meson production were extracted over a wide range of  $Q^2$ ,  $x_B$  and  $-t$ . The measurements shown in Fig. 4 are significantly different from zero in all kinematic bins. Our data provide a set of new observables in the kinematic range of  $t/Q^2$  where higher twist contributions may be significant. The overall comparison of experimental measurements with different theoretical calculations using the leading order demonstrates the importance of our results to improve parameterization of the GPD  $H_T$ . They indicate strong sensitivity to the practically unknown  $-t$  dependencies of the underlying chiral-odd GPDs, which may shed light on the role of higher twist contributions. Combined with the unpolarized structure function measurements and beam spin asymmetry results for  $\pi^0$  production from CLAS [13,11,12], these data provide important constraints for the parameterizations of GPDs  $H_T$  and  $\bar{E}_T$ , giving the first insight to the transverse space distributions of transversely polarized quarks [25,26].

### Acknowledgements

We acknowledge the outstanding efforts of the staff of the Accelerator and Physics Divisions at JLab. This work was supported in part by the U.S. Department of Energy and National Science Foundation, the French Centre National de la Recherche Scientifique and Commissariat à l'Énergie Atomique, the Italian Istituto Nazionale di Fisica Nucleare, the National Research Foundation of Korea and the U.K. Engineering and Physical Science Research Council. Jefferson



**Fig. 4.** The target and double spin asymmetry moments of exclusive  $\pi^0$  electroproduction as a function of  $-t$  for 2 bins in the  $(Q^2, x_B)$  plane (left and right columns). The shaded bands represent the systematic uncertainties (including fit uncertainties), and the curves show the predictions from two GPD formalisms: GK [20] (dashed) and GGL [21] (solid). Note that the  $A_{UL}^{\sin 2\phi}$  moment is zero in the GK model, and therefore is not shown.

Science Associates (JSA) operates the Thomas Jefferson National Accelerator Facility for the United States Department of Energy under contract DE-AC05-06OR23177.

## Appendix A. Supplementary material

Supplementary material related to this article can be found online at <http://dx.doi.org/10.1016/j.physletb.2017.02.032>.

## References

- [1] D. Müller, D. Robaschik, B. Geyer, F.-M. Dittes, J. Hořejší, Wave functions, evolution equations and evolution kernels from light ray operators of QCD, *Fortschr. Phys.* 42 (1994) 101–141.

- [2] X.-D. Ji, Deeply-virtual Compton scattering, *Phys. Rev. D* 55 (1997) 7114–7125.
- [3] A.V. Radyushkin, Scaling limit of deeply virtual Compton scattering, *Phys. Lett. B* 380 (1996) 417–425.
- [4] M. Burkardt, C.A. Miller, W.D. Nowak, Spin-polarized high-energy scattering of charged leptons on nucleons, *Rep. Prog. Phys.* 73 (2010) 016201.
- [5] M. Diehl, Generalized parton distributions, *Phys. Rep.* 388 (2003) 41–277.
- [6] P. Hoodbhoy, X. Ji, Helicity-flip off-forward parton distributions of the nucleon, *Phys. Rev. D* 58 (1998) 054006.
- [7] M. Diehl, Generalized parton distributions with helicity flip, *Eur. Phys. J. C* 19 (3) (2001) 485–492.
- [8] S. Ahmad, G.R. Goldstein, S. Liuti, Nucleon tensor charge from exclusive  $\pi^0$  electroproduction, *Phys. Rev. D* 79 (2009) 054014.
- [9] S.V. Goloskokov, P. Kroll, An attempt to understand exclusive  $\pi^+$  electroproduction, *Eur. Phys. J. C* 65 (2010) 137–151.
- [10] M. Burkardt, Transversity decomposition of quark angular momentum, *Phys. Lett. B* 639 (5) (2006) 462–464.
- [11] I. Bedlinskiy, et al., Measurement of exclusive  $\pi^0$  electroproduction structure functions and their relationship to transverse generalized parton distributions, *Phys. Rev. Lett.* 109 (2012) 112001.
- [12] I. Bedlinskiy, et al., Exclusive  $\pi^0$  electroproduction at  $W > 2$  GeV with CLAS, *Phys. Rev. C* 90 (2014) 025205.
- [13] R. De Masi, et al., Beam spin asymmetry in deep and exclusive  $\pi^0$  electroproduction, *Phys. Rev. C* 77 (2008) 042201.
- [14] E. Fuchey, et al., Exclusive neutral pion electroproduction in the deeply virtual regime, *Phys. Rev. C* 83 (2011) 025201.
- [15] M. Defurne, et al., Rosenbluth separation of the  $\pi^0$  electroproduction cross section, *Phys. Rev. Lett.* 117 (2016) 262001.
- [16] B.A. Mecking, et al., The CEBAF Large Acceptance Spectrometer (CLAS), *Nucl. Instrum. Methods A* 503 (2003) 513–553.
- [17] C.D. Keith, et al., A polarized target for the CLAS detector, *Nucl. Instrum. Methods A* 501 (2–3) (2003) 327–339.
- [18] T.W. Donnelly, A.S. Raskin, Considerations of polarization in inclusive electron scattering from nuclei, *Ann. Phys.* 169 (2) (1986) 247–351.
- [19] C.F. Perdrisat, V. Punjabi, M. Vanderhaeghen, Nucleon electromagnetic form factors, *Prog. Part. Nucl. Phys.* 59 (2) (2007) 694–764.
- [20] S.V. Goloskokov, P. Kroll, Transversity in hard exclusive electroproduction of pseudoscalar mesons, *Eur. Phys. J. A* 47 (2011) 112.
- [21] G.R. Goldstein, J.O.G. Hernandez, S. Liuti, Flexible parametrization of generalized parton distributions: the chiral-odd sector, *Phys. Rev. D* 91 (2015) 114013.
- [22] M. Göckeler, et al., Transverse spin structure of the nucleon from lattice-QCD simulations, *Phys. Rev. Lett.* 98 (2007) 222001.
- [23] M. Anselmino, et al., Update on transversity and Collins functions from SIDIS and data, *Nucl. Phys. B, Proc. Suppl.* 191 (2009) 98–107.
- [24] P. Kroll, Hard exclusive pion leptonproduction, *Few-Body Syst.* 57 (11) (2016) 1041–1050.
- [25] M. Diehl, P. Hagler, Spin densities in the transverse plane and generalized transversity distributions, *Eur. Phys. J. C* 44 (2005) 87–101.
- [26] B. Hannafious, M. Burkardt, The Boer–Mulders function: models and universality, *PoS LC2008* (2008) 032.