Old Dominion University ODU Digital Commons

Chemistry & Biochemistry Faculty Publications

Chemistry & Biochemistry

9-2012

Hot Methane Line Lists for Exoplanet and Brown Dwarf Atmospheres

Robert J. Hargreaves

Christopher A. Beale

Laurent Michaux

Melis Irfan

Peter F. Bernath Old Dominion University, pbernath@odu.edu

Follow this and additional works at: https://digitalcommons.odu.edu/chemistry_fac_pubs Part of the <u>Astrophysics and Astronomy Commons</u>

Repository Citation

Hargreaves, Robert J.; Beale, Christopher A.; Michaux, Laurent; Irfan, Melis; and Bernath, Peter F., "Hot Methane Line Lists for Exoplanet and Brown Dwarf Atmospheres" (2012). *Chemistry & Biochemistry Faculty Publications*. 85. https://digitalcommons.odu.edu/chemistry_fac_pubs/85

Original Publication Citation

Hargreaves, R. J., Beale, C. A., Michaux, L., Irfan, M., & Bernath, P. F. (2012). Hot methane line lists for exoplanet and brown dwarf atmospheres. *Astrophysical Journal*, 757(1), 46 doi:10.1088/0004-637X/757/1/46

This Article is brought to you for free and open access by the Chemistry & Biochemistry at ODU Digital Commons. It has been accepted for inclusion in Chemistry & Biochemistry Faculty Publications by an authorized administrator of ODU Digital Commons. For more information, please contact digitalcommons@odu.edu.

HOT METHANE LINE LISTS FOR EXOPLANET AND BROWN DWARF ATMOSPHERES

ROBERT J. HARGREAVES¹, CHRISTOPHER A. BEALE^{2,4}, LAURENT MICHAUX², MELIS IRFAN^{2,5}, AND PETER F. BERNATH^{1,3}

¹ Department of Chemistry, University of York, Heslington, York YO10 5DD, UK; rjh135@york.ac.uk

² Department of Physics, University of York, Heslington, York YO10 5DD, UK; cbeale@odu.edu, lm595@york.ac.uk, melis.irfan@postgrad.manchester.ac.uk ³ Department of Chemistry & Biochemistry, Old Dominion University, 4541 Hampton Boulevard, Norfolk, VA 23529-0126, USA; pbernath@odu.edu *Received 2012 June 25; accepted 2012 July 26; published 2012 September 4*

ABSTRACT

We present comprehensive experimental line lists of methane (CH₄) at high temperatures obtained by recording Fourier transform infrared emission spectra. Calibrated line lists are presented for the temperatures 300–1400°C at twelve 100°C intervals spanning the 960–5000 cm⁻¹ (2.0–10.4 μ m) region of the infrared. This range encompasses the dyad, pentad, and octad regions, i.e., all fundamental vibrational modes along with a number of combination, overtone and hot bands. Using our CH₄ spectra, we have estimated empirical lower state energies (E_{low} in cm⁻¹) and our values have been incorporated into the line lists along with line positions ($\tilde{\nu}$ in cm⁻¹) and calibrated line intensities (S' in cm molecule⁻¹). We expect our hot CH₄ line lists to find direct application in the modeling of planetary atmospheres and brown dwarfs.

Key words: brown dwarfs – infrared: general – methods: laboratory – molecular data – planetary systems – techniques: spectroscopic

Online-only material: color figures, machine-readable table

1. INTRODUCTION

Methane (CH₄) is the most fundamental alkane molecule and is second only to CO_2 in causing climate change (IPCC 2007). The presence of CH₄ in Earth's atmosphere is a consequence of both anthropogenic (e.g., landfills, rice cultivation, waste water treatment) and natural sources (e.g., methanogens, wetlands, wildfires). CH_4 can also be found throughout the solar system (Karkoschka 1994) including the atmospheres of Jupiter, Saturn, Uranus, Neptune and tentatively in the Martian atmosphere (Mumma et al. 2009). Titan, the largest moon of Saturn, has a dense atmosphere which is mostly made up of nitrogen but contains a relatively large proportion of CH₄ (approximately 1.4% in the stratosphere; Niemann et al. 2005). The concentration of CH₄ is higher near the surface and it is believed that a methane cycle exists which leads to the formation of liquid CH₄ lakes (Stofan et al. 2007). Interestingly, CH₄ has also been observed in the spectral energy distributions (SEDs) of cool substellar objects such as brown dwarfs (Cushing et al. 2006) and more recently in exoplanet atmospheres (Swain et al. 2008; Seager & Deming 2010).

Cool stars and brown dwarfs are classified according to the molecular features in the observed SED (Bernath 2009). The cooler the object, the more molecular signatures can be identified in the SED (Kirkpatrick 2005). L dwarfs display near-infrared electronic transitions of metal hydrides such as FeH (Hargreaves et al. 2010) and weaker TiO and VO bands (Geballe et al. 2002) which are more commonly associated with M dwarfs (Leggett et al. 2001; Bernath 2009). T dwarfs contain significant absorption due to H₂O and CH₄ (Cushing et al. 2006) and have been referred to as "methane dwarfs" (Hauschildt et al. 2009) because methane is the characteristic absorber. The newest category of brown dwarf (Y class) has recently been observed (Cushing et al. 2011) and displays characteristic NH₃ absorptions along with additional CH_4 absorption. Given these band assignments, the SEDs of brown dwarfs are complex and still contain numerous unassigned features.

The study of exoplanets is relatively new, as it was not until the mid-1990s that the first exoplanet was discovered (Mayor & Queloz 1995). In recent years, "hot Jupiters" are being discovered at an increasing rate and Kirkpatrick et al. (2011) were able to identify a further 100 using NASA's Wide-field Infrared Survey Explorer. It is now possible to probe the atmospheres of exoplanets (Charbonneau et al. 2002) by utilizing the natural transit as the planet passes in front of its parent star. The resulting flux undergoes a characteristic "dip" at specific wavelengths during the planetary transit (Charbonneau et al. 2000) and from the extremely small variations in the wavelength dependence of the dips it is possible to determine which molecules are present. By utilizing the transit method, H₂O (Barman 2008; Grillmair et al. 2008), CH₄ (Swain et al. 2008), CO (Swain et al. 2009b) and CO_2 (Swain et al. 2009a; Tinetti et al. 2010) have already been identified. Currently this method has only been successful for large hot Jupiters, but NASA's *Kepler* mission is discovering ever smaller planets which more closely resemble Earth, such as Kepler-22b (Borucki et al. 2012). Exoplanets are now thought to be the rule, rather than the exception, throughout the Milky Way (Cassan et al. 2012) and the ultimate exoplanet goal is to detect the distinctive signatures of extraterrestrial life through observation of the atmospheres of these distant planets.

Infrared spectroscopy of CH₄ is complex due to the high symmetry of the molecule and the polyad structure which arises from close-lying vibrational levels (Boudon et al. 2006). CH₄ is a tetrahedral molecule (T_d symmetry) and is a classic example of a spherical top. There are 3N - 6 = 9 infrared modes with four stretches and five bends. The four C–H stretches correspond to a symmetric a_1 stretch (v_1 at 2916.481 cm⁻¹) and a triply degenerate t_2 antisymmetric stretch (v_3 at 3019.493 cm⁻¹). The five bends correspond to a doubly degenerate e bending mode (v_2 at 1533.333 cm⁻¹) and a triply degenerate t_2 bending mode (v_4 at 1310.761 cm⁻¹). The spectroscopic constants of CH₄ are summarized in Table 1 (Albert et al. 2009; Herzberg 1991) and only

 ⁴ Currently at Department of Chemistry, and Biochemistry, Old Dominion University, 4541 Hampton Boulevard, Norfolk, VA 23529-0126, USA.
 ⁵ Currently at School of Physics and Astronomy, The University of Manchester, Oxford Road, Manchester M13 9PL, UK.

 Table 1

 The Fundamental Infrared Modes of CH4

Mode	Degeneracy	Band Origin ^a (cm ⁻¹)	Туре
$v_1(a_1)$	1	2916.481	Symmetric C-H stretch
$v_2(e)$	2	1533.333	Bend
$v_3(t_2)$	3	3019.493	Antisymmetric C-H stretch
$v_4(t_2)$	3	1310.761	Bend

Note.^a Taken from Albert et al. (2009).

the v_3 and v_4 fundamental modes are nominally infrared active; however, it is possible for other modes to gain intensity through strong interactions within the polyad structure. The interesting polyad structure of the infrared spectrum of CH₄ arises because the bending frequencies are approximately equal ($v_2 \approx v_4$), the stretching frequencies are also approximately equal ($v_1 \approx v_3$) and approximately twice the bending frequencies (Albert et al. 2009). For example, the pentad region near 3 μ m has five vibrational bands since $v_1 \approx v_3 \approx 2v_2 \approx 2v_4 \approx v_2 + v_4$.

CH₄ has also frequently been the target of high-quality theoretical calculations based on the solution of the vibrationrotation Schrödinger equation (Wang & Sibert 2002; Wang & Carrington 2004; Cassam-Chenaï 2003) using ab initio potential energy surfaces (Schwenke & Partridge 2001; Schwenke 2002; Nikitin et al. 2011). The accuracy of vibration-rotational calculations are typically far from experiment; however, recent calculations of nominally forbidden pure rotational lines (Cassam-Chenaï & Liévin 2012) are able to approach experimental accuracy.

Albert et al. (2009) have reported a study of CH₄ in the 0–4800 cm⁻¹ region and these lines form the basis of the HITRAN 2008 database (Rothman et al. 2009). Extensive work has been carried out in Grenoble including the preparation of a cold CH₄ line list (Mondelain et al. 2011; Wang et al. 2011, 2012; Campargue et al. 2012a) which has been successfully applied to the atmosphere of Titan (Campargue et al. 2012b). Recently focus has shifted to the generation of hot line lists for use in brown dwarfs and exoplanet models, either from ab initio calculations (Warmbier et al. 2009) or from experiment (Nassar & Bernath 2003; Perrin & Soufiani 2007; Thiévin et al. 2008).

Radiative transfer models are typically used to identify molecules in the atmospheres of exoplanets (Swain et al. 2008). Adaptations for hot-Jupiter atmospheres include known strongly absorbing molecules (i.e., H₂O and CH₄) and one of the options for CH₄ has been the HITRAN 2008 database (Rothman et al. 2009) combined with experimental data obtained by Nassar & Bernath (2003). HITRAN covers a broad spectral range but is intended for room temperature applications whereas the Nassar & Bernath (2003) data are only recorded at three higher temperatures (800, 1000, and 1273 K). This limits the accuracy of the modeled SEDs and leads to errors in deriving effective temperatures (Hauschildt et al. 2009). We hope to address this issue by providing a series of high-resolution, hot CH₄ line lists at temperatures relevant to exoplanet and brown dwarf atmospheres with the inclusion of empirical lower state energies where possible. This work will cover the dyad (ν_2 , ν_4), pentad $(v_1, v_3, 2v_2, 2v_4, v_2 + v_4)$, and octad $(3v_2, 3v_4, v_1 + v_2, v_1 + v_4)$ $v_3 + v_2$, $v_3 + v_4$, $2v_2 + v_4$, $v_2 + 2v_4$) regions of the infrared including numerous hot bands. Similar work has already been carried out on NH₃ (Hargreaves et al. 2011, 2012). Our data sets, and molecular opacities derived from them, can be used directly in exoplanet and brown dwarf models.

 Table 2

 Experimental Conditions

	Region A	Region B	Region C	Region D
Total file coverage (cm^{-1})	600-1900	1000-3000	1500-5000	2500-8000
Temperature range (°C)	300-1400	300-1400	300-1400	500-1300
Detector	MCT	MCT	InSb	InSb
Beam splitter	KBr	CaF ₂	CaF ₂	CaF ₂
Windows	KRS-5	CaF ₂	CaF ₂	CaF ₂
Lens	ZnSe	CaF ₂	CaF ₂	CaF ₂
Total scans	240	300	300	800
Resolution (cm^{-1})	0.01	0.015	0.015	0.02
Aperture (mm)	2.5	2.5	2.5	1.7
CH ₄ pressure (Torr) ^a	4.0	4.0	4.0	60.0
Zero-fill factor	×16	×16	×16	×16

Note. ^a This is an average pressure to the nearest 0.1 Torr.

2. EXPERIMENTAL METHOD

High-resolution laboratory emission spectra of hot CH_4 were recorded in four separate parts in order to maximize the signalto-noise for each spectral region. The four spectral regions are shown in Figure 1 for 1000°C. A similar setup was used in our previous work on NH₃ (Hargreaves et al. 2011, 2012) and only key experimental features will be summarized here.

The experiment coupled a Fourier transform infrared (FT-IR) spectrometer with a temperature controllable tube furnace capable of maintaining stable temperatures up to 1400°C with an accuracy $\pm 10^{\circ}$ C. The tube furnace surrounds the central portion of an alumina (Al₂O₃) tube which is sealed at both ends by infrared windows. CH₄ gas is allowed to flow through the alumina tube at a stable pressure in order to avoid the build up of impurities. The infrared emission from the alumina tube was focused onto the aperture of the FT-IR spectrometer using a lens. The small region between the spectrometer and tube furnace was purged with dry air in order to minimize H₂O absorption lines in the recorded spectra. Table 2 contains the experimental parameters used for all four regions.

Spectra were recorded at 100°C intervals between 300 and 1400°C for regions A, B, and C and between 500 and 1300°C for region D as this was the temperature coverage of observable CH_4 emission. Each region was limited using an appropriate filter chosen to allow for overlap between neighboring regions. The resolution of each spectral region was based on the Doppler and pressure broadening widths, and was selected to minimize the acquisition time whilst maintaining the maximum line resolution.

The temperature of the tube furnace was measured and maintained at high temperatures using a thermocouple combined with a programmable controller. The ends of the alumina tube required cooling to avoid damaging the rubber o-rings (which maintain the pressure seal) and the consequence of this is a temperature gradient within the tube. The observed infrared emission spectra are dominated by the high temperature central portion of the tube (see Figure 1) and the emission lines are assumed to be formed at the same temperature as the furnace.

The emission lines in each spectrum were picked using the program WSpectra (Carleer 2001) and the line intensity of each line was obtained by fitting a Voigt line profile. The number of CH_4 emission lines recorded for each region is given in Table 3.

A system response function was measured for each region by recording a blackbody spectrum emitted from a graphite rod placed at the center of the alumina tube (i.e., at the center



Figure 1. High-resolution laboratory emission spectra of CH₄ at 1000°C for all four regions (black) with the corresponding normalized instrument response function (blue). The baseline is caused by residual thermal emission from the tube walls and a small amount of absorption due to cold CH₄, CO₂ (ν_3 near 2350 cm⁻¹), and H₂O can be seen.

(A color version of this figure is available in the online journal.)

 Table 3

 Observed Emission Lines of CH4

Temperature	Region A	Region B	Region C	Region D	Total
(°C)					
300	4,365	2,589	4,118		11,072
400	6,656	3,566	7,752		17,974
500	7,950	4,725	9,438	5,242	27,355
600	9,524	6,021	14,109	7,766	37,420
700	10,381	6,344	15,376	10,067	42,168
800	11,189	6,951	19,297	12,696	50,133
900	11,607	7,276	21,148	14,688	54,719
1000	12,005	7,889	21,042	16,563	57,499
1100	12,105	7,667	21,850	16,062	57,684
1200	12,368	7,637	22,511	10,269 ^a	52,785
1300	12,235	8,118	23,332	9,640 ^a	53,325
1400	12,510	6,095 ^b	10,590 ^b		29,195

Notes.

^a These spectra were contaminated by C_2H_2 and C_2H_4 emission lines and have not been included in our analysis as the impurity lines could not be completely removed.

^b These spectra were unusually noisy and as a result the number of lines observed is reduced.

of the furnace). This accounts for the relative contribution of the system (e.g., windows, lens, filters, etc.) to the intensity of the observed lines. The spectrum is compared to a theoretical blackbody at the temperature of the furnace and then normalized. These instrument response functions are shown in Figure 1 along with the CH_4 emission lines at 1000°C.

It was necessary to calibrate both the wavenumber scale (*x*-axis) and the intensity scale (*y*-axis) using the HITRAN line list. The wavenumber scales were calibrated by selecting 20 strong, clear, symmetric lines and averaging the correction factor. Each temperature spectrum had similar calibration factors and typical values (for the 1000°C spectra) were 1.000001351 for region A, 1.000001677 for region B, 1.000001051 for region C, and 1.000000091 for region D. This results in a typical shift of 0.005 cm⁻¹ at 3000 cm⁻¹ for the ν_3 band. We therefore expect the accuracy of each line list to be ± 0.005 cm⁻¹ or better.

The intensity scales have also been calibrated according to HITRAN. First, each emission line list was converted into an arbitrary absorption scale to make them comparable to the intensities of the lines contained in HITRAN. This method has previously been used by Nassar & Bernath (2003) and more recently by Hargreaves et al. (2011, 2012) using the equation:

$$S_a = \frac{S_e}{\nu^3 \exp\left(-\frac{h\nu}{kT}\right)},\tag{1}$$

where S_a/S_e is the intensity of the absorbed/emitted line, ν is the frequency and *T* is the temperature.

Each line list was then put onto the same intensity scale as recorded by region A, and comparisons to HITRAN were THE ASTROPHYSICAL JOURNAL, 757:46 (10pp), 2012 September 20

Table 4Methane Partition Function				
Temperature	Partition Function ^a			
(K)				
296	590.52			
573	1,856.19			
673	2,648.82			
773	3,745.99			
873	5,267.13			
973	7,372.47			
1073	10,276.30			
1173	14,266.42			
1273	19,723.43			
1373	27,161.91			
1473	37,273.63			
1573	50,996.68			
1673	69,611.60			

Note. ^a Calculated using the TIPS_2011.for program (Laraia et al. 2011) based on calculations made by Wenger et al. (2008).

made. The HITRAN lines had to be scaled to the appropriate temperatures using

$$S' = S'_0 \frac{Q_0}{Q} \exp\left(\frac{E_{\text{low}}}{kT_0} - \frac{E_{\text{low}}}{kT}\right) \left[\frac{1 - \exp\left(-\frac{h\nu_{10}}{kT}\right)}{1 - \exp\left(-\frac{h\nu_{10}}{kT_0}\right)}\right], \quad (2)$$

where S' is the intensity, Q is the partition function, T is the temperature, v_{10} is the line frequency, and E_{low} is the empirical lower state energy (Bernath 2005). A zero subscript refers to the same parameters but at the reference temperature. The partition functions can be found in Table 4 and were calculated for each temperature using the TIPS_2011.for program (Laraia et al. 2011) based on calculations made by Wenger et al. (2008).

Wavenumber matches between our data and the temperaturescaled HITRAN lines were made $(\pm 0.005 \text{ cm}^{-1})$ and these intensities were compared. A calibrating factor could then be deduced but as the frequency increased, the intensity calibration factors became larger and more inconsistent. Regions A, B, and C could all be matched without difficulty since these regions cover strong emission bands; however, region D proved to be a problem. In the end, it was decided that an intensity calibration function could be extrapolated into region D based on the matches from regions A, B, and C. The intensity calibration used to convert the observed arbitrary intensity (S'_{arb}) to intensities in HITRAN units (S'_H) is

$$S'_H = [a(\tilde{\nu})^2 + b\tilde{\nu} + c]S'_{\rm arb},\tag{3}$$

where $\tilde{\nu}$ is in cm⁻¹ and S'_H is in cm molecule⁻¹. The calibration function is temperature dependent and was applied to all regions; the constants are given in Table 5.

Each region had to be combined to form a complete line list at each temperature. The optical filters used during the acquisition of each spectrum allowed the regions to overlap; the common lines in the overlapping regions have been used to bring all regions onto the same scale. The number of emission lines in the final line lists were maximized by appropriately choosing to combine the regions at 1500, 1830, and 3350 cm⁻¹. Because emission lines were only recorded above 960 cm⁻¹ and region D was intentionally truncated at 5000 cm⁻¹, the spectral regions are 960–1500 cm⁻¹ (region A), 1500–1830 cm⁻¹ (region B), 1830–3350 cm⁻¹ (region C), and 3350–5000 cm⁻¹ (region D), i.e., 960–5000 cm⁻¹ in total.

The emission lines for each temperature were compared to all other temperatures and all matchable lines were identified ($\pm 0.005 \text{ cm}^{-1}$). It was then possible to calculate the empirical lower state energy (E_{low}) of each line by rearranging Equation (2) so that

$$\ln\left(\frac{SQR_0}{S_0Q_0R}\right) = \frac{E_{\text{low}}}{kT_0} - \frac{E_{\text{low}}}{kT},\tag{4}$$

where

$$R_0 = 1 - \exp\left(-\frac{h\nu_{10}}{kT_0}\right).$$
(5)

The slope of points plotted using Equation (4) for each line yields the E_{low} and an example of this calculation is shown in Figure 2 for five CH₄ emission lines present at all temperatures.

This then forms the basis of the temperature dependent line lists which contain calibrated line positions ($\tilde{\nu}$ in cm⁻¹), calibrated line intensities (S' in cm molecule⁻¹) and empirical lower state energies (E_{low} in cm⁻¹). It was necessary to incorporate missing HITRAN lines into these line lists as explained in the results.

3. RESULTS

Using our method to calculate the E_{low} values, it is possible to identify the dyad, pentad, and octad infrared regions of CH₄ in the plots of E_{low} versus wavenumber (Figure 3).

The two most striking features are the v_4 region (dyad) around 1300 cm⁻¹ and the v_3 region (pentad) around 3020 cm⁻¹. The v_4 region contains lines of the fundamental v_4 bend and distinct $2v_4-v_4$ and $3v_4-2v_4$ hot bands at a E_{low} of 1300 and 2600 cm⁻¹, respectively. The v_3 region contains a strong *P*-, *Q*-, and *R*-branch of the $v_3 + v_4 - v_4$ hot band above a E_{low} of 1300 cm⁻¹ and parts of the fundamental *P*-, and *R*-branch of the v_3 mode. Another hot band, possibly the $2v_3 - v_3$, can be seen above a E_{low} of ~3000 cm⁻¹. Less obvious but still present are tentative observations of the *Q*-branch of the "forbidden" v_2 fundamental bend at approximately 1530 cm⁻¹. Also present are features of the $v_3 + v_4$ combination region (octad) around 4300 cm⁻¹, although the features in this region are less distinct as it is congested by many other bands. Comparisons to HITRAN for this region shows a complex band structure as expected due to the high density of transitions.

There is reasonable agreement between the empirical E_{low} values and the HITRAN E_{low} values (Figure 3) within the ν_3 and ν_4 region. A hot band at a lower state energy of approximately 3000 cm⁻¹ in the pentad region (tentatively assigned the $2\nu_3 - \nu_3$ hot band) is absent in HITRAN and is not very well defined in our data. We have observed emission lines with higher rotational levels than are present in HITRAN which is most apparent in the vicinity of the ν_3 region. The HITRAN database clearly contains many CH₄ lines, but the majority of E_{low} values have been obtained from calculations; these E_{low} values are visible in Figure 3 as horizontal "stripes" which correspond to the distinct rotational levels.

Our method is also able to distinguish rotational structure of the branches of the v_3 fundamental mode as shown in Figure 4. The rotational levels of a rigid spherical top are given by BJ(J + 1) but there is a $(2J + 1)^2$ -fold degeneracy for each J (i.e., (2J + 1) from the M quantum number and (2J + 1) from K). As the molecule rotates and distorts, the K degeneracy is partly lifted and the lines show a "cluster-splitting." Figure 4 highlights the observed structure which can be seen in the calculated E_{low} values of the v_3 region of CH₄.



Figure 2. Natural logarithm of the intensity ratios of five lines present in all 12 spectra are plotted against 1/kT (in cm), where *T* is the temperature. The slope of each line directly yields the empirical lower state energy (in cm⁻¹) and for the lines plotted: $E_{low}[1] = 1364.00 \text{ cm}^{-1}$ (red), $E_{low}[2] = 1590.79 \text{ cm}^{-1}$ (orange), $E_{low}[3] = 1644.41 \text{ cm}^{-1}$ (green), $E_{low}[4] = 1879.95 \text{ cm}^{-1}$ (blue), and $E_{low}[5] = 2036.83 \text{ cm}^{-1}$ (purple). (A color version of this figure is available in the online journal.)

	Table 5	
Wavenumber	Calibration	Function

Temperature	а	b	с
(°C)	$(/10^{12} \text{ cm}^3 \text{ molecule}^{-1})$	$(/10^{15} \text{ cm}^2 \text{ molecule}^{-1})$	$(/10^{18} \text{ cm molecule}^{-1})$
300	2.92	-9.75	12.00
400	1.48	-2.15	4.98
500	2.82	-6.67	9.03
600	3.62	-9.87	12.30
700	4.04	-10.70	13.30
800	2.56	-3.35	6.02
900	3.24	-6.17	8.63
1000	4.87	-12.40	14.20
1100	4.00	-7.77	9.33
1200	4.07	-7.18	7.95
1300	5.06	-12.00	12.60
1400	4.68	-7.48	7.29

It is worth noting that points below an E_{low} of approximately 1000 cm⁻¹ are absent from Figure 3 due to the temperature gradient within the alumina tube. The CH₄ in the cooler ends of the tube absorbs the low-*J* emission lines of the fundamental bands and accounts for the fact that only high-*J* levels are seen in the branches of the v_3 mode as shown in Figures 3 and 4.

Self-absorption appears in our emission spectra as either complete absorption resulting in a missing line, or as partial absorption which gives a distinctive double peak (Hargreaves et al. 2011). In order to provide more complete line lists we have added the missing lines from HITRAN to our line lists. We therefore incorporate HITRAN lines (± 0.0025 cm⁻¹) with an intensity greater than that of a threshold value at each temperature. Our sensitivity decreases toward higher wavenumbers (regions A, C, and D from Figure 1) and as a result the cutoff threshold is not a constant and changes over each region in order to match the sensitivity of the observations. The effect of adding HITRAN lines to our line lists is shown in Table 6 which lists the added HITRAN lines and the observed emission lines. In order

	Table 6	
ne	List Summary	1

Li

Temperature (°C)	Observed Lines	Added HITRAN Lines	Total Lines
300	6,297	3,269	9,566
400	6,970	12,793	19,763
500	12,621	20,288	32,909
600	19,255	29,040	48,295
700	23,438	31,868	55,306
800	30,090	39,361	69,451
900	34,513	40,571	75,084
1000	37,214	41,322	78,536
1100	38,923	42,912	81,835
1200	28,118	32,788	60,906
1300	29,717	30,746	60,463
1400	19,462	19,096	38,558



Figure 3. Comparison of the empirical lower state energies (top panel) with the HITRAN lower state energies (bottom panel). The dyad (ν_4 region), pentad (ν_3 region), and octad ($\nu_3 + \nu_4$) regions are clearly visible.

	1			
Line ^a	Wavenumber (cm ⁻¹)	Calculated E_{low}^{b} (cm ⁻¹)	HITRAN E_{low} (cm ⁻¹)	Difference %
ṽ[1]	1313.30	1364.00	1336.88	2.0
ν̃[2]	3072.57	1590.79	1494.72	6.4
ν̃[3]	1323.51	1644.41	1639.18	0.3
ν̃[4]	1304.20	1879.95	1780.71	5.6
ĩ[5]	1301.59	2036.83	1976.72	3.0

 Table 7

 Empirical Lower State Energy Errors

Notes.

^a These lines are displayed in Figure 2.

^b As shown in the slope of the lines in Figure 2.

to avoid double counting of lines, all lines in the emission line list within ± 0.0025 cm⁻¹ of a HITRAN line (with an intensity above the selection threshold) were deleted and replaced by the HITRAN lines.

Comparisons made with Nassar & Bernath (2003) at 1000°C ($\pm 0.0025 \text{ cm}^{-1}$) suggest our intensities are accurate to within a factor of two (Figure 5). To estimate the error in the empirical E_{low} values they were compared to those found in HITRAN for a number of lines which occur in both data sets. Table 7 shows that the percentage difference between the empirical E_{low} calculation and HITRAN for the lines plotted in Figure 2 is less than 10%. The examples shown are lines which appear in all spectra and as a result are the most reliable. These particular lines have also been removed from our final line lists because they coincide with

HITRAN lines. As the number of points used to calculate the lower state energy decreases, the error increases. It is therefore difficult to quote an overall error for our E_{low} values but we aim to represent the error by the inclusion of a quality factor within the line lists. "H" signifies that the line position, intensity and lower state energy has been inserted from the HITRAN database. "1" means that the lower state energy has been calculated from the intensities of a line that appears in 10 or more spectra. "2" refers to any line which appears in 6-9 spectra, and "3" denotes that the lower state energy has been determined from 3 to 5 spectra. Finally, "0" indicates that the lower state energy could not be determined from the spectra available; in this case it means that the line occurs in less than three spectra in total. It should be noted that for region D a maximum quality factor of only "2" is achievable. An extract of the hot CH₄ line list at 1000°C is displayed in Table 8; the complete line lists are available online in an electronic format.

4. DISCUSSION

We have produced CH₄ line lists which can be used at temperatures from 300°C to 1400°C. The calibration of the wavenumber scale for each spectrum was done by making wavenumber comparisons of 20 lines to the HITRAN database and this gives an overall accuracy of better than 0.005 cm⁻¹. The wavenumber calibration is straightforward but intensity calibration is more difficult. Line intensities are notoriously hard to calibrate and our calibration was made by comparing as many lines as possible to the HITRAN database. In the end our overall





Figure 4. Cluster splitting of the *P*-branch lines of the ν_3 mode of CH₄. Panel (a) contains the HITRAN E_{low} values plotted with the scaled HITRAN lines at 1000°C, and panel (b) contains the empirical E_{low} values plotted with the calibrated lines at 1000°C. The gray scale of the E_{low} value is taken from the intensity of the line at 1000°C.

line intensity accuracy is approximately a factor of two. Intensity matches for lower wavenumbers (regions A and B) were good and allowed a consistent calibration function to be determined. Each region overlapped the neighboring region with enough emission lines to place all regions on the same scale. For region D the intensity matches with HITRAN were effectively absent so a calibration factor could not be determined from this region alone. Since all regions are on the same scale, the calibration function was extrapolated into region D. In order to validate our extrapolation we were able to compare our 1000°C spectrum with the equivalent 1273 K spectrum from Nassar & Bernath

(2003). The matches ($\pm 0.0025 \text{ cm}^{-1}$) can be seen in Figure 5 and show a clear intensity correlation to within a factor of two which gives confidence in the calibration applied. The slope is present because the Nassar & Bernath (2003) calibrations were performed using a single factor whereas the calibration used here is a parabolic function.

It is difficult to estimate an error for the empirical E_{low} values presented here because self-absorption causes inaccurate intensities resulting in inaccurate empirical E_{low} values. This means comparisons can only be made between emission lines removed from the line lists and lines inserted from the HITRAN database.



Figure 5. Comparison of line intensities at 1000°C with the values for corresponding lines (within ± 0.0025 cm⁻¹) recorded by Nassar & Bernath (2003) at 1273 K. (A color version of this figure is available in the online journal.)



Figure 6. Histogram showing the ratios of the HITRAN E_{low} values to the empirical E_{low} values. The vertical bars of the histogram have been shaded to indicate the number of points with the quality factors (QF) 1, 2, and 3.

(A color version of this figure is available in the online journal.)

Comparison of the E_{low} values of the five lines in Figure 2 can give a false impression of the accuracy, as in Table 7. A better estimate can be obtained by plotting the ratio of E_{low} from HI-TRAN compared to the empirical E_{low} values of the removed lines (Figure 6 at 1000°C). It clearly shows that the most common value of the HITRAN E_{low} /empirical E_{low} ratio is approximately one and that the majority of empirical E_{low} values are accurate to within 20%. It can be seen that the most accurate lines (with a quality factor of "1") also display the smallest spread as opposed to the least accurate lines (quality factor of "3"). Figure 6 also shows why it was necessary to remove these lines because there is a tendency toward low ratios (<1) due to inaccurate line intensities caused by self-absorption. At high temperatures, thermal decomposition of methane is expected (Thiévin et al. 2008) and lower intensities are indeed observed for our high temperature spectra. The highest temperature spectra recorded in region D ($1200^{\circ}C$ and $1300^{\circ}C$) were contaminated by the absorption and emission of the products of thermal decomposition, namely C₂H₂ and C₂H₄. Attempts have been made to remove these species by using recent line lists (Moudens et al. 2011; Rothman et al. 2009); however, the emission features could not be completely removed. As this is the case, the hot CH₄ emission lines could not be distinguished from the products of thermal decomposition and the 1200°C and 1300°C files of region D have been excluded from our analysis.

 Table 8

 Extract of the 1000°C Hot CH4 Line List

Temperature (°C)	Wavenumber (cm ⁻¹)	Intensity $(\text{cm molecule}^{-1})$	$E_{\rm low}$ (cm ⁻¹)	Quality Factor
1000	1343.952393	4.21E-22	3.29E+03	1
1000	1343.962722	1.57E-23	1.82E+03	Н
1000	1343.974972	2.30E-23	2.17E+03	Н
1000	1343.980835	8.99E-23	3.08E+03	2
1000	1343.982910	8.99E-23	2.78E+03	1
1000	1343.992990	3.38E-23	1.93E+03	Н
1000	1344.015503	3.96E-23	5.20E+03	3
1000	1344.033569	1.19E-22	3.18E+03	1

(This table is available in its entirety in a machine-readable form in the online journal. A portion is shown here for guidance regarding its form and content.)

Empirical E_{low} values have been determined previously using a similar method by our group for the hot line lists of NH₃ (Hargreaves et al. 2011, 2012). It has also been used in the submillimeter regime (Fortman et al. 2010) and on "cold" CH₄ in the infrared (Sciamma-O'Brien et al. 2009; Lyulin et al. 2010; Wang et al. 2012). Fortman et al. (2010) obtained E_{low} values of an astrophysical "weed" (ethyl cyanide) using numerous cool (-40-140°C) spectra. Sciamma-O'Brien et al. (2009), Lyulin et al. (2010), and Wang et al. (2012) have used a similar approach to study CH₄ based on a small number of room temperature and cold (\sim 80 K) spectra. The method presented here uses a dozen "hot" (300-1400°C) spectra which were recorded under conditions that are similar to those found in the atmospheres of brown dwarfs and hot Jupiters. The use of a large number of spectra improves the accuracy of the E_{low} values. However a single spectrum takes us many hours to record (including heating and cooling of the system) so it is impractical to record many additional spectra. Nevertheless, combining both hot and cold CH₄ line lists would help to complete the knowledge of CH₄ in the infrared.

Our method produces a relatively small number of lines with high line position accuracy. An alternative approach is to use ab initio calculations which provide significantly more lines but the line positions have much lower accuracy. The two methods are complementary and they can be combined to produce a merged line list. Zobov et al. (2011) have used the ab initio approach and were able to assign quantum numbers to some of our new NH₃ data; similar calculations are currently underway for CH₄.

The line lists are available as electronic supplements and spectra can be provided upon request. The line lists should be used at the specified temperature; if a temperature is needed between those provided, we recommend scaling the lines using Equation (2).

5. CONCLUSION

Experimental line lists of hot CH₄ have been provided which contain calibrated wavenumbers (in cm⁻¹), calibrated intensities (in cm molecule⁻¹) and empirical lower state energies (in cm⁻¹). The line lists were based on 12 emission spectra recorded at $300-1400^{\circ}$ C in 100° C intervals using an FT-IR spectrometer and cover the 960–5000 cm⁻¹ wavenumber range. This range contains the dyad, pentad, and octad infrared regions of CH₄. The line lists can be used directly to model CH₄ in brown dwarfs and exoplanets. They also provide new experimental data for comparison with theoretical predictions.

Support for this work was provided by a Research Project Grant from the Leverhulme Trust and a Department of Chemistry (University of York) studentship. L.M. was supported by a Nuffield Foundation Undergraduate Research Bursary.

REFERENCES

- Albert, S., Bauerecker, S., Boudon, V., et al. 2009, Chem. Phys., 356, 131
- Barman, T. S. 2008, ApJ, 676, L61
- Bernath, P. F. 2005, Spectra of Atoms and Molecules (2nd ed.; Oxford: Oxford Univ. Press)
- Bernath, P. F. 2009, Int. Rev. Phys. Chem., 28, 681
- Borucki, W. J., Koch, D. G., Batalha, N., et al. 2012, ApJ, 745, 120
- Boudon, V., Rey, M., & Loëte, M. 2006, J. Quant. Spectrosc. Radiat. Transfer, 98, 394
- Campargue, A., Leshchishina, O., Wang, L., et al. 2012a, J. Quant. Spectrosc. Radiat. Transfer, 113, 1855
- Campargue, A., Wang, L., Mondelain, D., et al. 2012b, Icarus, 219, 110 Carleer, M. R. 2001, Proc. SPIE, 4168, 337
- Cassan, A., Kubas, D., Beaulieu, J.-P., et al. 2012, Nature, 481, 167
- Cassam-Chenaï, P., & Liévin, J. 2012, J. Chem. Phys., 136, 174309
- Cassam-Chenaï, P. 2003, J. Quant. Spectrosc. Radiat. Transfer, 82, 251
- Charbonneau, D., Brown, T. M., Latham, D. W., & Mayor, M. 2000, ApJ, 529, L45
- Charbonneau, D., Brown, T. M., Noyes, R. W., & Gilliland, R. L. 2002, ApJ, 568, 377
- Cushing, M. C., Kirkpatrick, J. D., Gelino, C. R., et al. 2011, ApJ, 743, 50
- Cushing, M. C., Roellig, T. L., Marley, M. S., et al. 2006, ApJ, 648, 614
- Fortman, S. M., Medvedev, I. R., Neese, C. F., & De Lucia, F. C. 2010, ApJ, 714, 476
- Geballe, T. R., Knapp, G. R., Leggett, S. K., et al. 2002, ApJ, 564, 466
- Grillmair, C. J., Burrows, A., Charbonneau, D., et al. 2008, Nature, 456, 767
- Hargreaves, R. J., Hinkle, K. H., Bauschlicher, C. W., et al. 2010, AJ, 140, 919
- Hargreaves, R. J., Li, G., & Bernath, P. F. 2011, ApJ, 735, 111
- Hargreaves, R. J., Li, G., & Bernath, P. F. 2012, J. Quant. Spectrosc. Radiat. Transfer, 113, 670
- Hauschildt, P. H., Warmbier, R., Schneider, R., & Barman, T. 2009, A&A, 504, 225
- Herzberg, G. 1991, Molecular Spectra and Molecular Structure: Volume III – Electronic Spectra and Electronic Structure of Polyatomic Molecules (Malabar, FL: Krieger)
- Intergovernmental Panel on Climate Change 2007, Fourth Assessment Report: Climate Change 2007: The AR4 Synthesis Report (Geneva: IPCC)
- Karkoschka, E. 1994, Icarus, 111, 174
- Kirkpatrick, J. D. 2005, ARA&A, 43, 195
- Kirkpatrick, J. D., Cushing, M. C., Gelino, C. R., et al. 2011, ApJS, 197, 19
- Laraia, A. L., Gamache, R. R., Lamouroux, J., Gordon, I. E., & Rothman, L. S. 2011, Icarus, 215, 391
- Leggett, S. K., Allard, F., Geballe, T. R., Hauschildt, P. H., & Schweitzer, A. 2001, ApJ, 548, 908
- Lyulin, O. M., Kassi, S., Sung, K., Brown, L. R., & Campargue, A. 2010, J. Mol. Spectrosc., 261, 91
- Mayor, M., & Queloz, D. 1995, Nature, 378, 355
- Mondelain, D., Kassi, S., Wang, L., & Campargue, A. 2011, Phys. Chem. Chem. Phys., 13, 7985
- Moudens, A., Georges, R., Benidar, A., et al. 2011, J. Quant. Spectrosc. Radiat. Transfer, 112, 540
- Mumma, M. J., Villanueva, G. L., Novak, R. E., et al. 2009, Science, 323, 1041
- Nassar, R., & Bernath, P. 2003, J. Quant. Spectrosc. Radiat. Transfer, 82, 279
- Niemann, H. B., Atreya, S. K., Bauer, S. J., et al. 2005, Nature, 438, 779
- Nikitin, A. V., Rey, M., & Tyuterev, V. G. 2011, Chem. Phys. Lett., 501, 179
- Perrin, M.-Y., & Soufiani, A. 2007, J. Quant. Spectrosc. Radiat. Transfer, 103,
- Rothman, L. S., Gordon, I. E., Barbe, A., et al. 2009, J. Quant. Spectrosc. Radiat. Transfer, 110, 533
- Schwenke, D. W. 2002, Spectrochim. Acta A, 58, 849
- Schwenke, D. W., & Partridge, H. 2001, Spectrochim. Acta A, 57, 887
- Sciamma-O'Brien, E., Kassi, S., Gao, B., & Campargue, A. 2009, J. Quant. Spectrosc. Radiat. Transfer, 110, 951
- Seager, S., & Deming, D. 2010, ARA&A, 48, 631
- Stofan, E. R., Elachi, C., Lunine, J. I., et al. 2007, Nature, 445, 61
- Swain, M. R., Tinetti, G., Vasisht, G., et al. 2009a, ApJ, 704, 1616
- Swain, M. R., Vasisht, G., & Tinetti, G. 2008, Nature, 452, 329
- Swain, M. R., Vasisht, G., Tinetti, G., et al. 2009b, ApJ, 690, L114

- Thiévin, J., Georges, R., Carles, S., et al. 2008, J. Quant. Spectrosc. Radiat. Transfer, 109, 2027
- Tinetti, G., Deroo, P., Swain, M. R., et al. 2010, ApJ, 712, L139 Wang, L., Kassi, S., Liu, A. W., Hu, S. M., & Campargue, A. 2011, J. Quant. Spectrosc. Radiat. Transfer, 112, 937
- Wang, L., Mondelain, D., Kassi, S., & Campargue, A. 2012, J. Quant. Spectrosc. Radiat. Transfer, 113, 47
- Wang, X.-G., & Carrington, T., Jr. 2004, J. Chem. Phys., 121, 2937
- Wang, X.-G., & Sibert, E. L., III 2002, Spectrochim. Acta A, 58, 863
- Warmbier, R., Schneider, R., Sharma, A. R., et al. 2009, A&A, 495, 655
- Wenger, Ch., Champion, J. P., & Boudon, V. 2008, J. Quant. Spectrosc. Radiat. Transfer, 109, 2976 Zobov, N. F., Shirin, S. V., Ovsyannikov, R. I., et al. 2011, J. Mol. Spectrosc.,
- 269, 104

ERRATUM: "HOT METHANE LINE LISTS FOR EXOPLANET AND BROWN DWARF ATMOSPHERES" (2012, ApJ, 757, 46)

 ROBERT J. HARGREAVES¹, CHRISTOPHER A. BEALE^{2,4}, LAURENT MICHAUX², MELIS IRFAN^{2,5}, AND PETER F. BERNATH^{1,3} ¹ Department of Chemistry, University of York, Heslington, York YO10 5DD, UK; rjh135@york.ac.uk, pbernath@odu.edu
 ² Department of Physics, University of York, Heslington, York YO10 5DD, UK; cbeale@odu.edu, Im595@york.ac.uk, melis.irfan@postgrad.manchester.ac.uk
 ³ Department of Chemistry & Biochemistry, Old Dominion University, 4541 Hampton Boulevard, Norfolk, VA 23529-0126, USA *Received 2013 July 8; published 2013 August 19*

Online-only material: machine-readable table

1. DETAILS OF CORRECTIONS

It has been brought to our attention that an erroneous supplementary line list was included in our published article. In this article, we presented comprehensive experimental line lists of CH_4 at high temperatures. This error was caused by the addition of temperature-adjusted HITRAN lines into our purely experimental line lists. These added HITRAN lines had the wrong temperature.

This problem has been addressed and, in correction, we have included a new line list summary which lists the added HITRAN lines and the observed emission lines (Table 6). Furthermore, the complete corrected line list is available online in electronic format and Table 8 is an extract of the hot CH_4 line list at 1000°C. We advise that the previous supplementary material not be used and this corrected version be used in its place.

We thank Michael Rey (Université de Reims) who brought this problem to our attention.

⁴ Currently at Department of Chemistry, and Biochemistry, Old Dominion University, 4541 Hampton Boulevard, Norfolk, VA 23529-0126, USA

⁵ Currently at School of Physics and Astronomy, The University of Manchester, Oxford Road, Manchester M13 9PL, UK

Temperature	Total Lines		
(°C)			
300	3,255	9,120	12,375
400	6,542	14,803	21,345
500	12,437	21,518	33,955
600	19,247	29,471	48,718
700	23,644	31,249	54,893
800	30,368	37,708	68,076
900	34,929	38,218	73,147
1000	37,706	38,559	76,265
1100	39,485	39,613	79,098
1200	28,532	30,419	58,951
1300	30,189	28,218	58,407
1400	19,788	17,067	36,855

Table 6

 Table 8

 Extract of the 1000°C Hot CH4 Line List

Temperature	Wavenumber (cm^{-1})	Intensity $(cm molecule^{-1})$	E_{low}	Quality Eactor
()	(cm)	(em molecule)	(cm)	Pactor
1000	1343.952393	4.21E-22	3.29E+03	1
1000	1343.962722	1.31E-23	1.82E+03	Н
1000	1343.974972	1.98E-23	2.17E+03	Н
1000	1343.980835	8.99E-23	3.08E+03	2
1000	1343.982910	8.99E-23	2.78E+03	1
1000	1343.992990	2.85E-23	1.93E+03	Н
1000	1344.015503	3.96E-23	5.20E+03	3
1000	1344.033569	1.19E-22	3.18E+03	1

(This table is available in its entirety in a machine-readable form in the online journal. A portion is shown here for guidance regarding its form and content.)