12-2017

Molecular Geometries and Other Properties of $\text{H}_2\text{O}\cdots\text{AgI}$ and $\text{H}_3\text{N}\cdots\text{AgI}$ as Characterised by Rotational Spectroscopy and *ab initio* Calculations

Chris Medcraft

Eva Gougoula

Dror M. Bittner
Old Dominion University

John C. Mullaney

Susana Blanco

See next page for additional authors

Follow this and additional works at: https://digitalcommons.odu.edu/chemistry_fac_pubs

Part of the [Biochemistry Commons](#), and the [Chemistry Commons](#)

Repository Citation

Medcraft, Chris; Gougoula, Eva; Bittner, Dror M.; Mullaney, John C.; Blanco, Susana; Tew, David P.; Walker, Nicholas R.; and Legon, Anthony C., "Molecular Geometries and Other Properties of $\text{H}_2\text{O}\cdots\text{AgI}$ and $\text{H}_3\text{N}\cdots\text{AgI}$ as Characterised by Rotational Spectroscopy and *ab initio* Calculations" (2017). *Chemistry & Biochemistry Faculty Publications*. 98.
https://digitalcommons.odu.edu/chemistry_fac_pubs/98

Original Publication Citation

Medcraft, C., Gougoula, E., Bittner, D. M., Mullaney, J. C., Blanco, S., Tew, D. P., . . . Legon, A. C. (2017). Molecular geometries and other properties of $\text{H}_2\text{O}\cdots\text{AgI}$ and $\text{H}_3\text{N}\cdots\text{AgI}$ as characterised by rotational spectroscopy and *ab initio* calculations. *The Journal of Chemical Physics*, 147(23) 234308. doi:<http://dx.doi.org/10.1063/1.5008744>

Authors

Chris Medcraft, Eva Gougoula, Dror M. Bittner, John C. Mullaney, Susana Blanco, David P. Tew, Nicholas R. Walker, and Anthony C. Legon

Molecular geometries and other properties of $\text{H}_2\text{O}\#\text{AgI}$ and $\text{H}_3\text{N}\#\text{AgI}$ as characterised by rotational spectroscopy and ab initio calculations

Chris Medcraft, Eva Gougoula, Dror M. Bittner, John C. Mullaney, Susana Blanco, David P. Tew, Nicholas R. Walker, and Anthony C. Legon

Citation: *The Journal of Chemical Physics* **147**, 234308 (2017);

View online: <https://doi.org/10.1063/1.5008744>

View Table of Contents: <http://aip.scitation.org/toc/jcp/147/23>

Published by the [American Institute of Physics](#)

Articles you may be interested in

[Structure, spectroscopy, and dynamics of the phenol-\(water\)₂ cluster at low and high temperatures](#)

The Journal of Chemical Physics **147**, 234307 (2017); 10.1063/1.5006055

[Halogen bonding properties of 4-iodopyrazole and 4-bromopyrazole explored by rotational spectroscopy and ab initio calculations](#)

The Journal of Chemical Physics **147**, 214303 (2017); 10.1063/1.5002662

[Towards a quantum chemical protocol for the prediction of rovibrational spectroscopic data for transition metal molecules: Exploration of CuCN, CuOH, and CuCCH](#)

The Journal of Chemical Physics **147**, 234303 (2017); 10.1063/1.5006931

[Channel branching ratios in \$\text{CH}_2\text{CN}^-\$ photodetachment: Rotational structure and vibrational energy redistribution in autodetachment](#)

The Journal of Chemical Physics **147**, 234309 (2017); 10.1063/1.5001475

[The reactivity of the Criegee intermediate \$\text{CH}_3\text{CHOO}\$ with water probed by FTMW spectroscopy](#)

The Journal of Chemical Physics **148**, 014308 (2018); 10.1063/1.5009033

[Photodissociation of van der Waals complexes of iodine \$\text{X}-\text{I}_2\$ \(\$\text{X} = \text{I}_2, \text{C}_2\text{H}_4\$ \) via charge-transfer state: A velocity map imaging investigation](#)

The Journal of Chemical Physics **147**, 234304 (2017); 10.1063/1.5001104

Molecular geometries and other properties of $\text{H}_2\text{O} \cdots \text{AgI}$ and $\text{H}_3\text{N} \cdots \text{AgI}$ as characterised by rotational spectroscopy and *ab initio* calculations

Chris Medcraft,¹ Eva Gougoula,¹ Dror M. Bittner,² John C. Mullaney,¹ Susana Blanco,³ David P. Tew,⁴ Nicholas R. Walker,^{1,a)} and Anthony C. Legon^{4,a)}

¹Chemistry-School of Natural and Environmental Sciences, Newcastle University, Bedson Building, Newcastle-upon-Tyne NE1 7RU, United Kingdom

²Department of Chemistry and Biochemistry, Old Dominion University, 4541 Hampton Boulevard, Norfolk, Virginia 23529-0126, USA

³Departamento de Química Física y Química Inorgánica, Facultad de Ciencias, Universidad de Valladolid, 47011 Valladolid, Spain

⁴School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 ITS, United Kingdom

(Received 9 October 2017; accepted 27 November 2017; published online 20 December 2017)

The rotational spectra of $\text{H}_3\text{N} \cdots \text{AgI}$ and $\text{H}_2\text{O} \cdots \text{AgI}$ have been recorded between 6.5 and 18.5 GHz by chirped-pulse Fourier-transform microwave spectroscopy. The complexes were generated through laser vaporisation of a solid target of silver or silver iodide in the presence of an argon gas pulse containing a low concentration of the Lewis base. The gaseous sample subsequently undergoes supersonic expansion which results in cooling of rotational and vibrational motions such that weakly bound complexes can form within the expanding gas jet. Spectroscopic parameters have been determined for eight isotopologues of $\text{H}_3\text{N} \cdots \text{AgI}$ and six isotopologues of $\text{H}_2\text{O} \cdots \text{AgI}$. Rotational constants, B_0 ; centrifugal distortion constants, D_J , D_{JK} or Δ_J , Δ_{JK} ; and the nuclear quadrupole coupling constants, $\chi_{aa}(\text{I})$ and $\chi_{bb}(\text{I}) - \chi_{cc}(\text{I})$ are reported. $\text{H}_3\text{N} \cdots \text{AgI}$ is shown to adopt a geometry that has C_{3v} symmetry. The geometry of $\text{H}_2\text{O} \cdots \text{AgI}$ is C_s at equilibrium but with a low barrier to inversion such that the vibrational wavefunction for the $\nu = 0$ state has C_{2v} symmetry. Trends in the nuclear quadrupole coupling constant of the iodine nucleus, $\chi_{aa}(\text{I})$, of $\text{L} \cdots \text{AgI}$ complexes are examined, where L is varied across the series (L = Ar, H_3N , H_2O , H_2S , H_3P , or CO). The results of experiments are reported alongside those of *ab initio* calculations at the CCSD(T)(F12*)/AVXZ level (X = T, Q). Published by AIP Publishing. <https://doi.org/10.1063/1.5008744>

I. INTRODUCTION

Microwave spectroscopy has recently been applied to explore the molecular structures of complexes in which a Lewis base is attached to either CuX , AgX , or AuX (where X is a halogen atom) to form $\text{L} \cdots \text{MX}$,^{1–11} where, for example, L = H_2O ,^{1,10} H_2S ,^{3–5} NH_3 ,¹² C_2H_2 ,^{6,8,11} or CO.^{13,14} The same technique has also been applied to study complexes that are similar in composition and structure but distinct in that L is not a typical Lewis base. For example, $\text{H}_2 \cdots \text{MX}$,^{15,16} $\text{Ar} \cdots \text{MX}$,^{17–19} $\text{Kr} \cdots \text{MX}$,^{18,20} and $\text{Xe} \cdots \text{MX}$ ²¹ have each been characterised. Significant structural changes were identified in each of H_2 ,^{15,16} C_2H_2 ,^{6,8,11} and C_2H_4 ⁶ on their attachment to a coinage metal atom. Both $\text{H}_2\text{O} \cdots \text{MX}$ and $\text{H}_2\text{S} \cdots \text{MX}$ adopt pyramidal geometries of C_s symmetry at equilibrium. $\text{H}_2\text{O} \cdots \text{MX}$ complexes were observed to undergo rapid inversion between two equivalent C_s geometries on the time scale of a molecular rotation such that the geometry of $\text{H}_2\text{O} \cdots \text{MX}$ is effectively C_{2v} (planar) in the $\nu = 0$ state.^{1,3–5,10,22} Rotational

spectroscopy has proven to be a powerful tool for investigation of the structure and dynamics in such complexes. Previous studies have made comparisons with hydrogen- and halogen-bonded complexes of similar geometrical structure.

Examination of the trend in the nuclear quadrupole coupling constant of chlorine, $\chi_{aa}(\text{Cl})$, across the $\text{L} \cdots \text{AgCl}$ ^{4,8,10,15,20,21} series revealed that the attachment of Ar to AgCl ¹⁹ induces only a very small change in the quantity relative to the value of the parameter in diatomic AgCl . The described change varies with L in the order $\text{CO} > \text{H}_2\text{S} > \text{H}_3\text{N} > \text{H}_2\text{O}$. A very similar trend was observed in the equivalent $\text{L} \cdots \text{CuCl}$ series.^{4,6,10,12} The variation in $\chi_{aa}(\text{Cl})$ provides an indication of the extent of electron redistribution accompanying the formation of each of these complexes. Evidently, stabilisation of $\text{OC} \cdots \text{MCl}$ through back-donation of electrons from d orbitals of the metal onto π^* orbitals of CO leads to a stronger interaction than that existing within $\text{Ar} \cdots \text{MCl}$. The present work aims to determine rotational constants, centrifugal distortion constants, and nuclear quadrupole coupling constants of the $\text{L} \cdots \text{AgI}$ complexes for which L = H_2O or H_3N and use these quantities to draw conclusions about the geometries, interaction strength, and electric charge redistribution associated with the complexes. The results will

^{a)}Authors to whom correspondence should be addressed: Nick.Walker@newcastle.ac.uk and a.c.legon@bristol.ac.uk

be compared with those from recent studies of other $L \cdots AgI$ to allow trends to be established across a wide range of $L \cdots AgI$ for the first time.

II. METHODS

A. Experimental

Broadband microwave spectra were measured using a chirped-pulse Fourier-transform microwave (CP-FTMW) spectrometer fitted with a laser ablation source. Detailed descriptions of the spectrometer and laser ablation source are provided in Refs. 9 and 23. The method employed to generate $H_3N \cdots AgI$ is identical to that used previously during a study of $H_2S \cdots AgI$.^{3,5} A gas sample containing a low concentration of precursors is diluted in argon and prepared at a total pressure of 6 bars. The sample is prepared to contain $\sim 1.5\%$ CF_3I and $\sim 4.0\%$ NH_3 . The mixed sample is introduced into the spectrometer via a pulsed nozzle where it passes over the surface of a silver rod from which material is ablated by the 2nd harmonic of an Nd:YAG laser pulse ($\lambda = 532$ nm, pulse duration of 10 ns, pulse energy of 20 mJ). The rotational and vibrational motions of molecules are subsequently cooled through supersonic expansion of the gas sample. The target rod is continually translated and rotated to expose a fresh surface to each laser pulse and ensure shot-to-shot reproducibility of signal intensities. Pulses are introduced with a repetition rate of approximately 2 Hz. Observations of rotational transition intensities suggest that the rotational temperature of probed species is of the order of 2 K. The generation of $H_2O \cdots AgI$ was achieved while using a solid target rod composed of silver iodide combined with a bespoke sample holder containing water and placed immediately in front of the nozzle.²⁴ Attempts to generate $H_2O \cdots AgI$ through vaporisation of a silver rod in the presence of gaseous H_2O and CF_3I precursors were unsuccessful. Isotopically enriched samples of $^{15}NH_3$ (Sigma-Aldrich, 98% ^{15}N), ND_3 (Sigma-Aldrich, 99% D), D_2O , and $H_2^{18}O$ were used to generate and record the spectra of isotopologues as appropriate.

The sequence employed to record broadband microwave spectra involves (i) polarization of the sample by a microwave chirp that sweeps from 6.5 to 18.5 GHz within 1 μs and then (ii) recording of the free induction decay (FID) of the molecular emission over a period of 20 μs . The sequence of (i) and (ii) is repeated eight times following each gas sample introduction pulse. The free induction decay (FID) of the polarization is digitized at 25 GS/s using a digital oscilloscope after down-mixing against a 19 GHz local oscillator. Frequency-domain spectra are obtained through Fourier transformation of the time-domain data to give a full width at half maximum (FWHM) of 100 kHz. The spectra of $H_3N \cdots AgI$ and $H_2O \cdots AgI$ were averaged for 600 k and 570 k free induction decays (FIDs), respectively, prior to being Fourier transformed.

B. *Ab initio* calculations

Geometry optimisations and frequency calculations were performed using the MOLPRO²⁵ and TURBOMOLE²⁶ packages at the CCSD(T)(F12*)²⁷ level of theory. The latter is

a coupled-cluster method that includes single and double excitations, explicit correlation,²⁸ and a perturbative treatment of triple excitations.²⁹ Basis set combinations consisting of aug-cc-pVXZ ($X = T, Q$) on C, H, and N and aug-cc-pVXZ-PP ($X = T, Q$) on Ag and I with relativistic effective core potentials MDF-28^{30,31} were used and will be referred to as AVXZ ($X = T, Q$). For the density fitting approximation which was used to accelerate the CCSD(T)(F12*) calculation, the respective def2-QZVPP basis sets were employed for the MP2 and Fock³² terms. For the complementary auxiliary basis required for the F12 treatment,³³ the def2-TZVPP MP2 density fitting basis sets³² were used. Only valence electrons were included in the correlation treatment.

III. RESULTS

A. Measurement of transition frequencies

1. $H_3N \cdots AgI$

As described in Sec. II A, the generation of $H_3N \cdots AgI$ was achieved by using CF_3I and NH_3 diluted in argon carrier gas interacting with the plasma produced by ablating silver. Transitions of CF_3I molecules were observed with high intensity, as were those of AgI . Transitions of $CF_3I \cdots NH_3$ ³⁴ were also observed but more weakly. Spectra containing AgI are distinctive by virtue of the characteristic isotopic abundance ratio of the two isotopes of silver and the hyperfine splitting introduced by the iodine nucleus ($I = 5/2$). This distinctiveness greatly assisted the initial assignments of the spectra of $H_3N \cdots AgI$ as shown in Fig. 1. Subsequent experiments employed synthetically enriched isotopic samples to confirm that these complexes are the carriers of the observed spectra and allowed measurements of shifts in the frequencies of transitions following isotopic substitution(s).

Spectra of the partially deuterated complexes, $HD_2N \cdots AgI$, were recorded in addition to that of $D_3N \cdots AgI$ during experiments that employed ND_3 . It is assumed that HD_2N was generated through reaction of a fraction of the introduced ND_3 sample with H_2O or NH_3 (perhaps present as a contaminant) within the sample vessel.

FIG. 1. Expanded section of the spectrum recorded in the 6.5–18.5 GHz frequency range for $H_3N \cdots ^{109}AgI$ (left) and $H_3N \cdots ^{107}AgI$ (right) showing the $(J + 1) \leftarrow J = 10 \leftarrow 9$ transition. Blue and red downward pointing lines indicate the simulated spectra of isotopologues containing ^{109}Ag and ^{107}Ag , respectively.

FIG. 2. Expanded section of the spectrum obtained in the 6.5–18.5 GHz frequency range for $\text{H}_2\text{O} \cdots \text{AgI}$, $\text{HDO} \cdots \text{AgI}$, and $\text{H}_2^{18}\text{O} \cdots \text{AgI}$ (from left to right) in the regions of the $(J+1) \leftarrow J = 10 \leftarrow 9$ transition (first two panels) and the $(J+1) \leftarrow J = 11 \leftarrow 10$ transition (far right panel). Blue and red downward pointing lines indicate the simulated spectra of isotopologues containing ^{109}Ag and ^{107}Ag , respectively.

2. $\text{H}_2\text{O} \cdots \text{AgI}$

Laser vaporisation of silver iodide in the presence of argon and a low concentration of H_2O allowed the observation of intense features that assign to the AgI diatomic molecule.³⁵ Transitions of the $\text{Ar} \cdots \text{AgI}$ complex characterised previously¹⁷ and a distinctive transition of the water dimer near 12 321 MHz³⁶ were also observed. Series of other transitions share some qualitative features of the spectrum of $\text{Ar} \cdots \text{AgI}$ but were clearly associated with a different molecular carrier. The frequency increment between $(J+1) \leftarrow J$ transitions was consistent with a lighter molecule than $\text{Ar} \cdots \text{AgI}$ suggesting $\text{H}_2\text{O} \cdots \text{AgI}$ as a possible carrier (Fig. 2). Subsequent experiments to probe $\text{HDO} \cdots \text{AgI}$ and $\text{H}_2^{18}\text{O} \cdots \text{AgI}$ isotopologues revealed isotopic shifts consistent with the assignment to $\text{H}_2\text{O} \cdots \text{AgI}$.

B. Spectroscopic analysis

Parameters of the Hamiltonian shown in Eq. (1) were fitted to measured transition frequencies using Western's program PGOPHER³⁷ for each molecule,

$$H = H_R - \frac{1}{6} \mathbf{Q}(\text{I}) : \nabla \mathbf{E}(\text{I}), \quad (1)$$

where H_R is Watson's A-reduced Hamiltonian³⁸ in the Γ representation for a semi-rigid symmetric top in the case of $\text{H}_3\text{N} \cdots \text{AgI}$. For such symmetric-top species, ground-state rotational constants are denoted by A_0 , B_0 , and C_0 and the determinable quartic centrifugal distortion constants are

denoted by D_J and D_{JK} which appear in H_R . Fits of the transition frequencies of $\text{H}_2\text{O} \cdots \text{AgI}$, and also of the $\text{D}_2\text{HN} \cdots \text{AgI}$ isotopologue of $\text{H}_3\text{N} \cdots \text{AgI}$, employed the version of H_R appropriate to a semi-rigid, nearly prolate asymmetric top. The only centrifugal distortion constants of $\text{H}_2\text{O} \cdots \text{AgI}$ that contributed significantly to transition frequencies reported here were Δ_J and Δ_{JK} .

The second term of Eq. (1) describes coupling of the electric quadrupole moment of the I nucleus with the electric field gradient at that nucleus, where $\mathbf{Q}(\text{I})$ is the iodine nuclear quadrupole moment dyadic and $\nabla \mathbf{E}(\text{I})$ is the dyadic of the electric field gradient at the iodine nucleus. The nuclear quadrupole coupling constants of iodine are denoted by $\chi_{aa}(\text{I})$ and $\chi_{bb}(\text{I}) - \chi_{cc}(\text{I})$, with the latter required only when fitting the spectra of $\text{H}_2\text{O} \cdots \text{AgI}$. It was unnecessary to include the corresponding term that describes nuclear quadrupole coupling associated with ^{14}N ($I = 1$) in the Hamiltonian of Eq. (1) because hyperfine structure arising from this nucleus was not apparent in the observed spectra of $\text{H}_3\text{N} \cdots \text{AgI}$ at the resolution of the present experiments. The naturally occurring isotopes of silver, ^{107}Ag and ^{109}Ag , each have $I = 1/2$ and consequently nuclear electric quadrupole moments of zero. Magnetic nuclear spin-rotation and spin-spin interactions were not observed at the resolution of the present experiments, consistent with previous observations performed on similar molecules. A Kaiser-Bessel window function was used to perform the Fourier transform of data for both $\text{H}_3\text{N} \cdots \text{AgI}$ and $\text{H}_2\text{O} \cdots \text{AgI}$.

The spectroscopic constants evaluated for the various isotopologues of $\text{H}_3\text{N} \cdots \text{AgI}$ and $\text{H}_2\text{O} \cdots \text{AgI}$ are provided

TABLE I. Experimentally determined spectroscopic constants of isotopologues of $\text{H}_3\text{N} \cdots \text{AgI}$.^a

	$\text{H}_3\text{N} \cdots ^{107}\text{AgI}$	$\text{H}_3\text{N} \cdots ^{109}\text{AgI}$	$\text{D}_3\text{N} \cdots ^{107}\text{AgI}$	$\text{D}_3\text{N} \cdots ^{109}\text{AgI}$
B_0 (MHz)	866.07484(54)	862.33436(78)	808.39412(51)	805.38125(72)
D_J (kHz)	0.0867(33)	0.0817(44)	0.0628(22)	0.0602(32)
D_{JK} (kHz)	8.65(30)	8.46(27)	7.00(24)	6.97(15)
$\chi_{aa}(\text{I})$ (MHz)	−795.42(54)	−795.31(69)	−794.61(72)	−793.40(76)
N^b	18	24	19	19
$\sigma_{r.m.s.}^c$ (kHz)	9.9	11.3	10.0	6.9

^aNumbers in parentheses are the standard deviation in units of the last significant figure.

^bNumber of nuclear quadrupole hyperfine components included in the fit.

^cRoot mean square deviation of the fit.

TABLE II. Experimentally determined spectroscopic constants of isotopologues of $\text{H}_3\text{N} \cdots \text{AgI}$ containing ^{15}N or D.

	$\text{H}_3\text{ }^{15}\text{N} \cdots ^{107}\text{AgI}$	$\text{H}_3\text{ }^{15}\text{N} \cdots ^{109}\text{AgI}$	$\text{HD}_2\text{N} \cdots ^{107}\text{AgI}$	$\text{HD}_2\text{N} \cdots ^{109}\text{AgI}$
$(B_0 + C_0)/2$ (MHz)	850.20711(23) ^{a,b}	846.68607(30) ^b	1653.1886(38)	1646.7204(34)
D_J (kHz)	0.0645(12)	0.0662(14)	[0.0628] ^c	0.0611(75) ^c
D_{JK} (kHz)	8.62(13)	8.55(17)	...	...
$\chi_{aa}(\text{I})$ (MHz)	-796.90(26)	-796.60(31)	-794.3(19)	-795.9(30)
N^d	41	28	6	11
$\sigma_{r.m.s.}^e$ (kHz)	7.3	8.1	9.7	14.0

^aNumbers in parentheses are the standard deviation in units of the last significant figure.^bResults for $\text{H}_3\text{ }^{15}\text{N} \cdots ^{107}\text{AgI}$ and $\text{H}_3\text{ }^{15}\text{N} \cdots ^{109}\text{AgI}$ are included in this table for concise presentation. Note that $B_0 = C_0 = (B_0 + C_0)/2$ for this symmetric rotor.^cWatson's A reduction was employed.^dNumber of nuclear quadrupole hyperfine components included in the fit.^eRoot mean square deviation of the fit.TABLE III. Experimentally determined spectroscopic constants of isotopologues of $\text{H}_2\text{O} \cdots \text{AgI}$.

	$\text{H}_2\text{O} \cdots ^{107}\text{AgI}$	$\text{H}_2\text{O} \cdots ^{109}\text{AgI}$	$\text{H}_2\text{ }^{18}\text{O} \cdots ^{107}\text{AgI}$	$\text{H}_2\text{ }^{18}\text{O} \cdots ^{109}\text{AgI}$
$(B_0 + C_0)$ (MHz)	1690.1972(28) ^a	1683.2961(28)	1629.6036(72)	1623.486(13)
$(B_0 - C_0)$ (MHz)	1.64674(20)	1.63346(20)	1.52831(40)	1.51669(46)
Δ_J (kHz)	0.07072(79)	0.07111(77)	0.0649(17)	0.0593(28)
Δ_{JK} (kHz)	8.642(91)	8.626(87)	8.80(17)	8.69(19)
$\chi_{aa}(\text{I})$ (MHz)	-885.54(11)	-885.33(10)	-884.24(49)	-885.10(71)
$(\chi_{bb}(\text{I}) - \chi_{cc}(\text{I}))$ (MHz)	-12.73(28)	-12.79(25)	-13.6(13)	-13.2(17)
N^b	75	68	40	27
$\sigma_{r.m.s.}^c$ (kHz)	5.5	5.1	9.6	9.3

^aNumbers in parentheses are the standard deviation in units of the last significant figures.^bNumber of nuclear quadrupole hyperfine components included in the fit.^cRoot mean square deviation of the fit.TABLE IV. Experimentally determined spectroscopic constants of isotopologues of $\text{HDO} \cdots \text{AgI}$.

	$\text{HDO} \cdots ^{107}\text{AgI}$	$\text{HDO} \cdots ^{109}\text{AgI}$
$(B_0 + C_0)$ (MHz)	1648.9189(38) ^a	1642.5292(60)
Δ_J (kHz)	0.0679(10)	0.07090(16)
$\chi_{aa}(\text{I})$ (MHz)	-885.08(24)	-884.42(41)
$(\chi_{bb}(\text{I}) - \chi_{cc}(\text{I}))$ (MHz)	...	...
N^b	28	26
$\sigma_{r.m.s.}^c$ (kHz)	5.3	7.5

^aNumbers in parentheses are the standard deviation in units of the last significant figures.^bNumber of nuclear quadrupole hyperfine components included in the fit.^cRoot mean square deviation of the fit.

in Tables I–IV. Only R-branch, a -type transitions³⁹ having $K_{-1} = 0$ or 1 were observed for $\text{H}_3\text{ }^{14/15}\text{N} \cdots ^{107/109}\text{AgI}$, $\text{D}_3\text{ }^{14}\text{N} \cdots ^{107/109}\text{AgI}$, and $\text{H}_2\text{ }^{16/18}\text{O} \cdots ^{107/109}\text{AgI}$, as expected for molecules having a large rotational constant A_0 , which ensures rotational energy levels having $K_{-1} \geq 2$ are too high in energy to be detectably populated at the temperature of the expanded gas pulse. The rotational constant, A_0 , could not be determined from the available data for either the symmetric-top species, $\text{H}_3\text{N} \cdots \text{AgI}$, or for the nearly prolate asymmetric rotor, $\text{H}_2\text{O} \cdots \text{AgI}$. This parameter was therefore fixed equal to the appropriate calculated values 180 GHz and 60 GHz, when fitting data and simulating spectra for $\text{H}_3\text{N} \cdots \text{AgI}$ and $\text{H}_2\text{O} \cdots \text{AgI}$, respectively. For $\text{H}_2\text{ }^{16}\text{O} \cdots \text{AgI}$ and $\text{H}_2\text{ }^{18}\text{O} \cdots \text{AgI}$, transitions having $K_{-1} = 0$ or 1 were

observed, which allowed determination of $(B_0 + C_0)$ and $(B_0 - C_0)$. Transitions having $K_{-1} = 1$ were not observed for $\text{HD}_2\text{N} \cdots ^{107/109}\text{AgI}$ or $\text{HDO} \cdots ^{107/109}\text{AgI}$, so $B_0 - C_0$ was fixed to zero when fitting $K_{-1} = 0$ transitions for these species. All other isotopologues for which data were recorded are symmetric rotors. The output files of spectroscopic fits performed in PGOPHER are provided as the [supplementary material](#).

For the $\text{H}_2\text{O} \cdots \text{AgI}$, $\text{HDO} \cdots \text{AgI}$, and $\text{H}_2\text{ }^{18}\text{O} \cdots \text{AgI}$ isotopologues, rotational transitions of the type $(J + 1) \leftarrow J$ were observed and were regularly separated in energy by a factor of $B_0 + C_0$. As indicated, determination of the $B_0 - C_0$ value for $\text{H}_2\text{O} \cdots \text{AgI}$ and $\text{H}_2\text{ }^{18}\text{O} \cdots \text{AgI}$ was possible due to the detection of $K_{-1} = 1$ transitions. For $\text{HDO} \cdots \text{AgI}$, only $K_{-1} = 0$ transitions were observed. The ratio of relative intensities of the $K_{-1} = 1$ and $K_{-1} = 0$ transitions (for a given $(J + 1) \leftarrow J$) of $\text{H}_2\text{O} \cdots \text{AgI}$ and $\text{H}_2\text{ }^{18}\text{O} \cdots \text{AgI}$ was approximately 3:1 in favour of the former. This strongly implies that $\text{H}_2\text{O} \cdots \text{AgI}$ adopts either a geometry that is planar and of C_{2v} symmetry at equilibrium or an alternative arrangement that has C_s symmetry at equilibrium, a pyramidal configuration at O, but with a sufficiently low potential energy barrier to the planar C_{2v} form that the wavefunction of the ground vibrational state has C_{2v} symmetry. The orientation of the lone pair on the oxygen atom in the conventional model of the water molecule supports the second of these possibilities, leading to the equilibrium geometry of the general type shown in Fig. 3. This conclusion is corroborated by the *ab initio* calculations at the CCSD(T)(F12)/AVTZ level of theory which find a

FIG. 3. The experimentally determined r_0 geometry of $\text{H}_2\text{O} \cdots \text{AgI}$ drawn to scale (see Table V).

barrier to inversion of only 125 cm^{-1} . As a consequence of the described inversion in the vibrational ground state, each of $\text{H}_2\text{O} \cdots \text{AgI}$ and $\text{H}_2^{18}\text{O} \cdots \text{AgI}$ contain a pair of protons that are exchanged by a C_2^a rotation and the usual arguments predict a 3:1 nuclear spin statistical weight ratio for $K_{-1} = 1$ to $K_{-1} = 0$ states. Cooling by collisional transfer⁴⁰ of population from $K_{-1} = 1$ (triplet) levels to $K_{-1} = 0$ (singlet) levels is a spin-forbidden process such that population of the $K_{-1} = 1$ levels remains close to the room temperature value. Consequently, transitions involving levels with $K_{-1} = 1$ are readily observable in our experiment. Conversely, in $\text{HDO} \cdots \text{AgI}$, the H and D nuclei are not equivalent particles and collisional transfer from $K_{-1} = 1$ to $K_{-1} = 0$ levels is no longer spin-forbidden. Cooling of $K_{-1} = 1$ states then occurs and $K_{-1} = 1$ transitions are not observed.⁴⁰ Transitions having $K_{-1} = 1$ are also observed for $\text{H}_3\text{N} \cdots \text{AgI}$, consistent with r_0 geometries for each complex that are of C_{3v} symmetry and have $\angle(\text{N}-\text{Ag}-\text{I})$ of 180° . The model geometry of $\text{H}_3\text{N} \cdots \text{AgI}$ is shown in Fig. 4. Similar arguments to those presented above explain the absence of $K_{-1} = 1$ transitions for the partially deuterated isotopologues of $\text{NH}_3 \cdots \text{AgI}$, an observation consistent with previous results obtained through studies of $\text{H}_3\text{N} \cdots \text{MX}$ complexes.^{12,13}

C. Molecular geometry

Figures 3 and 4, respectively, show the model geometries of the $\text{H}_2\text{O} \cdots \text{AgI}$ and $\text{H}_3\text{N} \cdots \text{AgI}$ complexes. Structural parameters in these models were directly fitted to moments of inertia using Kisiel's STRFIT program⁴¹ to yield r_0 results. For each of the above complexes, isotopic substitutions were available at all atoms except iodine, which permitted measurement of the spectra of six isotopologues of $\text{H}_2\text{O} \cdots \text{AgI}$ and eight isotopologues of $\text{H}_3\text{N} \cdots \text{AgI}$. It was assumed that the r_0 geometries of the isolated H_2O and NH_3 monomers do not change on formation of each complex when determining the remaining r_0 structural parameters. The rationale for the described assumption has been presented previously.¹⁰ Partial substitution structures (r_s) were also determined for $\text{H}_2\text{O} \cdots \text{AgI}$ and $\text{H}_3\text{N} \cdots \text{AgI}$ using Kraitchman's equations.⁴² The available isotopic data allow determination of $r_s(\text{Ag}-\text{O}) = 2.232(2) \text{ Å}$ for $\text{H}_2\text{O} \cdots \text{AgI}$ (Table V) and $r_s(\text{Ag}-\text{N})$ for

FIG. 4. The experimentally determined geometry of $\text{H}_3\text{N} \cdots \text{AgI}$ drawn to scale (see Table VI).

TABLE V. r_0 and r_e geometries of $\text{H}_2\text{O} \cdots \text{AgI}$.

	Expt. (r_0)	Calc. (r_e) ^a
$r(\text{Ag}-\text{I})$ (Å)	2.536(3)	2.5374
$r(\text{Ag}-\text{O})$ (Å)	2.227(7) ^b	2.2329
$r(\text{H}-\text{O})$ (Å)	[0.95785]	0.9605
φ (deg)	36.3(12)	40.74

^aCalculated *ab initio* at the CCSD(T) (F12*)/AVTZ level.

^bThe coordinates determined for the silver and oxygen atoms imply a r_s result of 2.232(2) Å for this parameter.

$\text{H}_3\text{N} \cdots \text{AgI} = 2.182(1) \text{ Å}$ (Table VI). Under the assumption that the geometry of the NH_3 sub-unit is unchanged from that in the r_0 geometry of H_3N , it is further possible to determine $\angle(\text{Ag}-\text{N}-\text{H}) = 110.93(3)^\circ$ for $\text{H}_3\text{N} \cdots \text{AgI}$. The *ab initio* calculated geometries of $\text{H}_2\text{O} \cdots \text{AgI}$ and $\text{H}_3\text{N} \cdots \text{AgI}$ are also included in Tables V and VI, respectively. We note that the agreement between the observed and calculated angle $\angle\text{Ag} \cdots \text{X}-\text{H}$ is excellent for $\text{X} = \text{N}$ but for $\text{X} = \text{O}$ the experimental value of ϕ is smaller by 4.4° . The poorer agreement in the case of $\text{X} = \text{O}$ probably arises because the potential energy barrier to the inversion of configuration at O in $\text{H}_2\text{O} \cdots \text{AgI}$ is low (as it is in $\text{H}_2\text{O} \cdots \text{AgCl}$ ¹⁰ and $\text{H}_2\text{O} \cdots \text{AgF}$ ²²) and therefore this motion makes a large amplitude contribution to the zero-point motion. The *ab initio* version of the angle is an equilibrium quantity and does not reflect such motion. On the other hand, the inversion is quenched in the case of $\text{X} = \text{N}$ and there is no corresponding large amplitude contribution to the zero-point motion. We note that the experimental value of ϕ is smaller than its *ab initio* counterpart in each of $\text{H}_2\text{O} \cdots \text{AgCl}$,¹⁰ $\text{H}_2\text{O} \cdots \text{CuCl}$,¹⁰ and $\text{H}_2\text{O} \cdots \text{AgF}$.²²

D. Intermolecular stretching force constants

We recently proposed a model⁴³ that allows in principle some of the quadratic force constants of $\text{L} \cdots \text{M}-\text{X}$ complexes to be established from centrifugal distortion constants determined experimentally. This model goes beyond that of Millen⁴⁴ by describing the centrifugal distortion constant D_J (for symmetric top or linear complexes) or Δ_J (for asymmetric-top complexes) in terms of three quadratic bond-stretching force constants F_{11} , F_{22} , and F_{12} , where the subscripts 1 and 2, respectively, refer to the $\text{M}-\text{X}$ and $\text{L} \cdots \text{M}$ bonds (such that $F_{11} = F_{\text{M}-\text{X}}$ and $F_{22} = F_{\text{L} \cdots \text{M}}$) with F_{12} being the off-diagonal coupling term. The forms of the equations relating D_J or Δ_J to F_{11} , F_{22} , and F_{12} are set out in detail in Refs. 7 and 43. If it is assumed that F_{12} is zero, these equations reduce to Eq. (2) when applied to a molecule like $\text{H}_2\text{O} \cdots \text{AgI}$ having effectively C_{2v} symmetry,

TABLE VI. r_0 , r_s , and r_e geometries of $\text{H}_3\text{N} \cdots \text{AgI}$.

	Expt. (r_0)	Expt. (r_s)	Calc. (r_e) ^a
$r(\text{Ag}-\text{I})$ (Å)	2.5375(3)	...	2.5457
$r(\text{Ag}-\text{N})$ (Å)	2.180(1)	2.182(1)	2.1767
$r(\text{H}-\text{N})$ (Å)	[1.01699]	[1.01619]	1.0128
$\angle(\text{Ag}-\text{N}-\text{H})$ (deg)	110.86(5)	110.93(3)	111.64

^aCalculated *ab initio* at the CCSD(T) (F12*)/AVQZ level.

$$h\Delta_J = \frac{\hbar^4}{4} \left\{ \frac{1}{(I_{bb}^e)^4} + \frac{1}{(I_{cc}^e)^4} \right\} \left\{ (m_I a_I)^2 / F_{11} + (m_I a_I + m_{Ag} a_{Ag})^2 / F_{22} \right\}, \quad (2)$$

where m_{Ag} and m_I are the appropriate nuclide masses, a_{Ag} and a_I are the principal axis coordinates of the Ag and I atoms and I_{bb}^e and I_{cc}^e are the equilibrium principal moments of inertia of the given isotopologue of $H_2O \cdots AgI$. Here it is assumed that zero-point quantities may be used in Eq. (2) without serious error. The corresponding form of Eq. (2) for D_J of the symmetric-top molecule $H_3N \cdots AgI$ is obtained by setting $I_{bb}^e = I_{cc}^e$.

By assuming that F_{11} is equal to the force constant of the isolated Ag–I diatomic molecule (145.8 N m^{-1}),⁷ F_{22} could be evaluated for $H_2O \cdots AgI$ and $H_3N \cdots AgI$ by fitting the appropriate centrifugal distortion constants. Data for the ^{15}N isotopologues were used in the case of $H_3N \cdots AgI$ because the absence of broadening through unresolved ^{14}N nuclear quadrupole hyperfine structure leads to more accurate values of D_J . The resulting values of F_{22} are compared with the same quantity for several other molecular complexes of silver iodide in Table VII. Also included in Table VII are the corresponding quantities that arise from the use of the Millen model (these are referred to as k_σ to use Millen's nomenclature). It is clear that the Millen model severely underestimates this force constant, even for cases where the Lewis base is bound relatively weakly to the silver atom, as discussed in Refs. 7 and 43.

To inspect the sensitivity of F_{22} to the assumption of unchanged F_{11} , F_{22} was further evaluated over a range of F_{11} values around that of free AgI. A plot showing the variation of F_{22} with the force constant F_{11} is displayed in Fig. 5. The horizontal asymptote (where $F_{11} \rightarrow \infty$) represents the case in which AgI can be regarded as rigid, and thus the value then obtained for the force constant F_{22} is equal to that evaluated under the Millen model. The vertical asymptote ($F_{22} \rightarrow \infty$) represents the value of the force constant F_{11} below which a negative (and therefore unphysical) value of F_{11} is required to fit the centrifugal distortion constant. It is worth noting that

TABLE VII. Experimentally determined quadratic force constants, F_{22} , for $L \cdots AgI$ complexes ($L = Ar, H_3N, H_2O, H_2S, H_3P$, or CO).

	$F_{22} (\text{N m}^{-1})^a$	$k_\sigma (\text{N m}^{-1})^b$
$Ar \cdots AgI^c$	20.2(8)	14.8
$H_2O \cdots AgI^d$	83(3)	13.9
$H_2S \cdots AgI^e$	92.0(17)	29.0
$H_3P \cdots AgI^f$	122(5)	31.7
$OC \cdots AgI^g$	156(26)	47
$H_3N \cdots AgI^d$	192(38)	15.1

^aCalculated by using the two-force constant model to fit D_J values, assuming $F_{11} = 145.8 \text{ N m}^{-1}$, the value for free AgI in its vibrational ground state. Numbers in parentheses are one standard deviation of the fit in units of the last significant figures.

^b k_σ values were determined from D_J values by means of the model described by Millen (Ref. 44).

^cReference 17.

^dThis work.

^eReference 3.

^fReference 7.

^gReference 9. F_{22} recalculated from D_J values reported in this reference.

FIG. 5. Dependence of F_{22} on the assumed value of F_{11} for various $L \cdots AgI$ complexes. The vertical dotted line indicates the value of the force constant of free AgI.

the ordering of F_{22} for these complexes is different depending on whether it is assumed that F_{11} is equal to 145.8 N m^{-1} (the value appropriate to the AgI diatomic molecule) or infinity (as would be implicit if $M-X$ were approximated as a rigid diatomic molecule). By examining the slope of the curve near $F_{11} = 145.8 \text{ N m}^{-1}$, it can be seen that the sensitivity of F_{22} to the value of F_{11} changes greatly over the series of Lewis bases shown, with the highest sensitivity being $F_{11} \approx F_{22}$.

E. Electric charge redistribution across the Ag–I bond upon formation of $H_2O \cdots AgI$

The ionicity, i_c , or fractional ionic character of the iodine nucleus can be estimated from the expression

$$i_c = 1 + \frac{\chi_{aa}(I)}{eQq_{(5,1,0)}(I)}, \quad (3)$$

where $eQq_{(5,1,0)}(I)$ is the contribution to the iodine nuclear quadrupole coupling constant along the a inertial axis that arises from a single electron in an iodine $5p_a$ orbital and is equal to 2292.71 MHz .⁴² Table VIII compares the ionicities of the complexes studied here with those of other Lewis base-silver iodide complexes similarly obtained. For all these complexes, the ionicity of the Ag–X bond increases upon attachment of the Lewis base. The largest increases are observed for the H_3P , OC , NH_3 and H_2S complexes, all lying between $i_c = 0.64$ and 0.68 ,

TABLE VIII. Ionicity of Ag–I in $L \cdots AgI$ complexes (where $L = Ar, H_3N, H_2O, H_2S, H_3P$, or CO).

Molecule	$\chi_{aa}(I) (\text{MHz})$	i_c^a
Ag–I	$-1062.5299(15)^b$	0.537
$Ar \cdots AgI^c$	$-985.411(54)$	0.570
$H_2O \cdots AgI$	$-885.54(11)$	0.613
$H_2S \cdots AgI^d$	$-807.36(4)$	0.648
$H_3N \cdots AgI$	$-795.42(54)$	0.65
$OC \cdots AgI^e$	$-769.84(22)$	0.66
$H_3P \cdots AgI^f$	$-733.83(34)$	0.68

^aCalculated using Eq. (3) and $eQq_{(5,1,0)}(I)$ values cited in Ref. 43.

^bReference 35.

^cReference 17.

^dReference 3.

^eReference 9.

^fReference 7.

whereas i_c for isolated AgI is 0.537. The complex with H₂O causes a smaller increase in the ionicity of the Ag–X subunit. This is different from the trend observed when these Lewis bases bind to silver chloride.^{4,10,17} The ordering of the i_c values displayed in Table VIII is also different from the ordering of complexes according to the value of the force constant, F_{22} . It might be expected that the change in charge distribution across the Ag–I bond upon attachment of a Lewis base, as measured by the change in nuclear quadrupole coupling constant, will be approximately proportional to the strength of the interaction of the Lewis base with the silver atom. Whilst this is generally true for the complexes listed in Tables VII and VIII, there are some notable exceptions. The force constant (F_{22}) of the bond formed between water and silver iodide is similar in magnitude to that for the intermolecular bond in H₂S···AgI but the values of i_c for the corresponding complexes span a relatively wide range from 0.61 to 0.65. Likewise NH₃ is calculated to bind more strongly to the silver atom in AgI ($F_{22} = 192(38) \text{ N m}^{-1}$) than does H₂S to Ag in H₂S···AgI ($F_{22} = 92.0(1.7) \text{ N m}^{-1}$) but has a very similar value of i_c .

IV. CONCLUSIONS

Two Lewis base-silver iodide complexes, L···AgI (L = H₃N and H₂O), have been generated through laser vaporisation of a solid target in the presence of a gas sample undergoing supersonic expansion and characterized by pure rotational spectroscopy. The basic geometries of the complexes studied are qualitatively the same as those established previously for several other members of the L···AgCl series. The values of $r(\text{Ag–O})$ and φ for H₂O···AgCl are 2.229(35) Å and 42(5)°, respectively, whereas those for H₂O···AgI are 2.227(7) Å and 36.3(12)°, respectively. Comparing results for H₃N···AgCl and H₃N···AgI, the values of $r(\text{Ag–N})$ are 2.154 44(6) Å and 2.180(1) Å, respectively. Evidently, the identity of the halogen atom has only a small effect on structural parameters associated with interactions between L and AgX sub-units. It has previously been noted that φ for H₂O···AgCl is very similar in magnitude to an equivalent parameter in the geometry of H₂O···HCl and H₂O···CuCl. These observations are all therefore consistent with a model in which the geometry of each complex is principally determined by the orientation of the lone pair on the oxygen atom which aligns with the axis of the bond in the electrophile acting as the Lewis acid.

SUPPLEMENTARY MATERIAL

See [supplementary material](#) for the final cycle of the fit to the ground-state rotational spectrum of each reported isotopologue of H₃N···AgI and H₂O···AgI using PGOPHER and the spectroscopic constants thereby determined.

ACKNOWLEDGMENTS

The authors thank the European Research Council for a postdoctoral fellowship awarded to C.M., a Ph.D. studentship awarded to J.C.M., and for project funding (Grant No. CPFTMW-307000); Newcastle University for Ph.D. studentships awarded to D.M.B. and E.G. The University of Bristol is thanked for a Senior Research Fellowship awarded

to A.C.L. S.B. thanks the Ministerio de Educación, Cultura y Deporte of Spain, “Salvador de Madariaga” program, for financial support.

- ¹S. J. Harris, A. C. Legon, N. R. Walker, and D. E. Wheatley, *Angew. Chem., Int. Ed.* **49**(1), 181–183 (2010).
- ²G. S. Grubbs II, D. A. Obenchain, H. M. Pickett, and S. E. Novick, *J. Chem. Phys.* **143**(2), 029901 (2015).
- ³C. Medcraft, D. M. Bittner, D. P. Tew, N. R. Walker, and A. C. Legon, *J. Chem. Phys.* **145**(19), 194306 (2016).
- ⁴N. R. Walker, D. P. Tew, S. J. Harris, D. E. Wheatley, and A. C. Legon, *J. Chem. Phys.* **135**(1), 014307 (2011).
- ⁵S. Z. Riaz, S. L. Stephens, W. Mizukami, D. P. Tew, N. R. Walker, and A. C. Legon, *Chem. Phys. Lett.* **531**, 1–5 (2012).
- ⁶S. L. Stephens, D. M. Bittner, V. A. Mikhailov, W. Mizukami, D. P. Tew, N. R. Walker, and A. C. Legon, *Inorg. Chem.* **53**(19), 10722–10730 (2014).
- ⁷S. L. Stephens, D. P. Tew, N. R. Walker, and A. C. Legon, *Phys. Chem. Chem. Phys.* **18**(28), 18971–18977 (2016).
- ⁸S. L. Stephens, W. Mizukami, D. P. Tew, N. R. Walker, and A. C. Legon, *J. Chem. Phys.* **137**(17), 174302 (2012).
- ⁹S. L. Stephens, W. Mizukami, D. P. Tew, N. R. Walker, and A. C. Legon, *J. Chem. Phys.* **136**(6), 064306 (2012).
- ¹⁰V. A. Mikhailov, F. J. Roberts, S. L. Stephens, S. J. Harris, D. P. Tew, J. N. Harvey, N. R. Walker, and A. C. Legon, *J. Chem. Phys.* **134**(13), 134305 (2011).
- ¹¹D. P. Zaleski, S. L. Stephens, D. P. Tew, D. M. Bittner, N. R. Walker, and A. C. Legon, *Phys. Chem. Chem. Phys.* **17**(29), 19230–19237 (2015).
- ¹²D. M. Bittner, D. P. Zaleski, S. L. Stephens, D. P. Tew, N. R. Walker, and A. C. Legon, *J. Chem. Phys.* **142**(14), 144302 (2015).
- ¹³D. M. Bittner, S. L. Stephens, D. P. Zaleski, D. P. Tew, N. R. Walker, and A. C. Legon, *Phys. Chem. Chem. Phys.* **18**(19), 13638–13645 (2016).
- ¹⁴C. J. Evans, L. M. Reynard, and M. C. L. Gerry, *Inorg. Chem.* **40**(24), 6123–6131 (2001).
- ¹⁵G. S. Grubbs II, D. A. Obenchain, H. M. Pickett, and S. E. Novick, *J. Chem. Phys.* **141**(11), 114306 (2014).
- ¹⁶D. A. Obenchain, D. S. Frank, G. S. Grubbs II, H. M. Pickett, and S. E. Novick, *J. Chem. Phys.* **146**(20), 204302 (2017).
- ¹⁷C. Medcraft, J. C. Mullaney, N. R. Walker, and A. C. Legon, *J. Mol. Spectrosc.* **335**, 61–67 (2017).
- ¹⁸C. J. Evans, A. Lesarri, and M. C. L. Gerry, *J. Am. Chem. Soc.* **122**(25), 6100–6105 (2000).
- ¹⁹C. J. Evans and M. C. L. Gerry, *J. Chem. Phys.* **112**(3), 1321–1329 (2000).
- ²⁰L. M. Reynard, C. J. Evans, and M. C. L. Gerry, *J. Mol. Spectrosc.* **206**(1), 33–40 (2001).
- ²¹S. A. Cooke and M. C. L. Gerry, *Phys. Chem. Chem. Phys.* **6**(13), 3248–3256 (2004).
- ²²S. L. Stephens, D. P. Tew, N. R. Walker, and A. C. Legon, *J. Mol. Spectrosc.* **267**(1–2), 163–168 (2011).
- ²³S. L. Stephens and N. R. Walker, *J. Mol. Spectrosc.* **263**(1), 27–33 (2010).
- ²⁴D. Loru, M. A. Bermúdez, and M. E. Sanz, *J. Chem. Phys.* **145**(7), 074311 (2016).
- ²⁵H.-J. Werner, P. J. Knowles, G. Knizia, F. R. Manby, and M. Schütz, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.* **2**(2), 242–253 (2012).
- ²⁶TURBOMOLE V7.1, a Development of the University of Karlsruhe, 1989–2007 and TURBOMOLE GmbH, since 2007, available at <http://www.turbomole.com>.
- ²⁷C. Hättig, D. P. Tew, and A. Köhn, *J. Chem. Phys.* **132**(23), 231102 (2010).
- ²⁸C. Hättig, W. Klopper, A. Köhn, and D. P. Tew, *Chem. Rev.* **112**(1), 4–74 (2012).
- ²⁹K. Raghavachari, G. W. Trucks, J. A. Pople, and M. Head-Gordon, *Chem. Phys. Lett.* **157**(6), 479–483 (1989).
- ³⁰K. A. Peterson, D. Figgen, E. Goll, H. Stoll, and M. Doig, *J. Chem. Phys.* **119**(21), 11113–11123 (2003).
- ³¹I. S. Lim, P. Schwerdtfeger, B. Metz, and H. Soll, *J. Chem. Phys.* **122**(10), 104103 (2005).
- ³²A. Hellweg, C. Hättig, S. Höfener, and W. Klopper, *Theor. Chem. Acc.* **117**(4), 587–597 (2007).
- ³³E. F. Valeev, *Chem. Phys. Lett.* **395**(4), 190–195 (2004).
- ³⁴S. L. Stephens, N. R. Walker, and A. C. Legon, *Phys. Chem. Chem. Phys.* **13**(46), 20736–20744 (2011).
- ³⁵S. G. Batten, A. G. Ward, and A. C. Legon, *J. Mol. Struct.* **780–781**, 300–305 (2006).

- ³⁶R. D. Thomas and J. S. Muentner, *J. Chem. Phys.* **60**(7), 2929–2930 (1974).
- ³⁷C. M. Western, *J. Quant. Spectrosc. Radiat. Transfer* **186**, 221–242 (2017).
- ³⁸J. K. G. Watson, *J. Chem. Phys.* **46**(5), 1935–1949 (1967).
- ³⁹S. A. Cooke and P. Ohring, *J. Spectrosc.* **2013**, 1–10.
- ⁴⁰E. Knozinger, P. Suppan, and A. C. Legon, *Faraday Discuss. Chem. Soc.* **86**, 269–270 (1988).
- ⁴¹Z. Kisiel, *J. Mol. Spectrosc.* **218**, 58–67 (2003).
- ⁴²R. L. Cook and W. Gordy, *Microwave Molecular Spectra*, 1st ed. (Interscience Publishers, Durham, North Carolina, 1970).
- ⁴³D. M. Bittner, N. R. Walker, and A. C. Legon, *J. Chem. Phys.* **144**(7), 074308 (2016).
- ⁴⁴D. J. Millen, *Can. J. Chem.* **63**(7), 1477–1479 (1985).