Old Dominion University ODU Digital Commons

Physics Faculty Publications

Physics

2001

Effective Field Theory for the Small-x Evolution

I. Balitsky Old Dominion University, ibalitsk@odu.edu

Follow this and additional works at: https://digitalcommons.odu.edu/physics_fac_pubs Part of the Astrophysics and Astronomy Commons, Elementary Particles and Fields and String Theory Commons, Nuclear Commons, and the Quantum Physics Commons

Repository Citation

Balitsky, I., "Effective Field Theory for the Small-x Evolution" (2001). *Physics Faculty Publications*. 105. https://digitalcommons.odu.edu/physics_fac_pubs/105

Original Publication Citation

Balitsky, I. (2001). Effective field theory for the small-*x* evolution. *Physics Letters B*, 518(3–4), 235-242. doi:https://doi.org/10.1016/S0370-2693(01)01041-3

This Article is brought to you for free and open access by the Physics at ODU Digital Commons. It has been accepted for inclusion in Physics Faculty Publications by an authorized administrator of ODU Digital Commons. For more information, please contact digitalcommons@odu.edu.



18 October 2001

PHYSICS LETTERS B

Physics Letters B 518 (2001) 235-242

www.elsevier.com/locate/npe

Effective field theory for the small-x evolution

I. Balitsky^{a,b}

^a Physics Department, Old Dominion University, Norfolk, VA 23529, USA
 ^b Theory Group, Jefferson Lab, Newport News, VA 23606, USA

Received 31 July 2001; accepted 16 August 2001 Editor: H. Georgi

Abstract

The small-*x* behavior of structure functions in the saturation region is determined by the non-linear generalization of the BFKL equation. I suggest the effective field theory for the small-*x* evolution which solves formally this equation. The result is the 2 + 1 functional integral for the structure functions at small *x*.

© 2001 Published by Elsevier Science B.V. Open access under CC BY license.

PACS: 12.38.Bx; 11.10.Jj; 11.55.Jy

The great success of pQCD in describing the Q^2 behavior of structure functions of deep inelastic scattering (DIS) can be traced back to the fact that the Q^2 -dependence is governed by DGLAP evolution equations which have two remarkable properties: they are linear equations, and the evolution at high Q^2 is purely perturbative (the non-perturbative physics enters the game only when we lower the normalization point μ^2 down to the typical hadronic scale ~ 1 GeV). The higher-order terms of perturbative expansion for both the coefficient functions and the anomalous dimensions of the light-cone operators lie in the same framework of linear evolution and lead to the corrections $\sim \alpha_s$, α_s^2 , etc.

The situation for the small-*x* DIS is more complicated. In the leading logarithmic approximation (LLA) the small-*x* asymptotics is described by the BFKL pomeron [1]. It is possible to reformulate the BFKL equation as an evolution equation where the relevant operators are Wilson lines — infinite gauge links [2]. ¹ The evolution of the two-Wilson-line operator ("color dipole") with respect to the slope of Wilson lines reproduces the BFKL equation.

Unfortunately, the theoretical status of the BFKL evolution is not as clear as the DGLAP one (for the review, see Refs. [3]). The biggest problem is the lack of unitarity: the power behavior of the BFKL cross section violates the Froissart bound and therefore, in order get the true asymptotics at small x, we must go beyond the LLA. At this step, we face a new problem. In the DGLAP case, the sub-leading logarithms follow the same general pattern of linear DGLAP equation and the problem is purely technical: calculating the loop corrections to the kernels. In the case of small-x evolution there are also α_s corrections to the BFKL kernel [5], but, on the top of that, there are the unitarity corrections which lie outside the framework of the BFKL equation. At small α_s and x, these corrections seem to dominate over the NLO BFKL ones [6].

¹ At high energies the particles move so fast that their trajectories can be approximated by straight lines collinear to their velocities. The proper degrees of freedom for the fast particles moving along the straight lines are the (infinite) gauge factors ordered along the straight line [4].

Another problem with the BFKL evolution is infrared instability. We can safely apply pQCD to the small-*x* DIS if the characteristic transverse momenta of the gluons k_{\perp} in the gluon ladder are large. For the the first few diagrams, one can check by explicit calculation that the characteristic k_{\perp}^2 are $\sim Q^2$. However, as *x* decreases, it turns out that the characteristic transverse momenta in the middle of the gluon ladder drift to $\Lambda_{\rm QCD}$ making the application of pQCD questionable.

Recently, an idea has emerged that these two difficulties may cancel each other out. Consider the DIS from the heavy nuclei where the large density sets the saturation scale Q_s [7–10] which effectively cuts the integration over k_{\perp} even at relatively low energy. As we shall see below, the small-*x* evolution in this case is non-linear which leads to the growth of the saturation scale with energy, see the discussion in Refs. [7–12]. It is natural to assume that even for the DIS from the nucleon where there is no saturation at low energies, the saturation scale at sufficiently small *x* may be generated by the non-linear evolution itself. Indeed, the parton recombination described by the non-linear evolution must balance at some point the effects of parton splitting so the partons will reach the state of the saturation. In this high-density regime the coupling constant is small but the characteristic fields are large, making a perfect case for the application of the semiclassical QCD methods [9,13,14]. The high-density regime of QCD can serve as a bridge between the domain of pQCD and the "real" non-perturbative QCD regime governed by the physics of confinement.

In this Letter I suggest the effective field theory which describes the small-*x* evolution in the saturation region. First, let me remind the OPE for high-energy amplitudes derived in [2]. Consider the amplitude of forward $\gamma^*\gamma^*$ -scattering at small $x_B = Q^2/s$. In the target frame, the virtual photon splits into $q\bar{q}$ pair which approaches the nucleon at high speed. Due to the high speed the classical trajectories of the quarks are straight lines collinear to the momentum of the incoming photon q. The corresponding operator expansion switched between nucleon states has the form [2]:

$$\int d^4x \, e^{iq \cdot x} \langle p | T\{j_\mu(x) j_\nu(0)\} | p \rangle = \int d^2x_\perp I_{\mu\nu}(x_\perp) \langle p | \operatorname{Tr}\{\widehat{U}(x_\perp)\widehat{U}^\dagger(0)\} | p \rangle, \tag{1}$$

where $I_{\mu\nu}(x_{\perp})$ is a certain numerical function of the transverse separation of quarks x_{\perp} and virtuality of the photon $Q^2 = -q^2$. The relevant operators $U(U^{\dagger})$ are gauge factors ordered along the classical trajectories which are almost light-like lines stretching from minus to plus infinity:

$$U(z_{\perp}) = P \exp\left(i \int_{-\infty}^{\infty} du \, e^{\mu} A_{\mu}(ue + z_{\perp})\right),\tag{2}$$

where e is collinear to q and z_{\perp} is the transverse position of the Wilson line.

It turns out that the small-*x* behavior of structure functions is governed by the evolution of these operators with respect to the deviation of the Wilson lines from the light cone; this deviation serves as a kind of "renormalization point" for these operators. At infinite energy, the vector *e* is light-like and the corresponding matrix elements of the operators (2) have a logarithmic divergence in longitudinal momenta. To regularize it, we consider operators corresponding to large but finite velocity and take $e_{\zeta} = e_1 + \zeta e_2$, where $e_1 = (q - \frac{q^2}{2pq}p)$ and $e_2 = p$ are the lightlike vectors close to the directions of the colliding particles. Now, instead of studying the energy-dependence of the physical amplitude we must investigate the dependence of the operators (2) on the slope ζ . Large energies mean small ζ and we can sum up logarithms of ζ instead of logarithms of *s* (at present, we can do it only in the leading logarithmic approximation (LLA) $\alpha_s \ll 1$, $\alpha_s \ln \frac{s}{m^2} \sim 1$). The equation governing the dependence of *U*



Fig. 1. BFKL evolution in terms of Wilson-line operators (denoted by dotted lines).

on ζ has the form $[2]^2$

$$\zeta \frac{d}{d\zeta} \mathcal{U}(x_{\perp}, y_{\perp}) = \frac{\alpha_s N_c}{2\pi^2} \int dz_{\perp} \frac{(x_{\perp} - y_{\perp})^2}{(x_{\perp} - z_{\perp})^2 (z_{\perp} - y_{\perp})^2} \times \left\{ \mathcal{U}(x_{\perp}, z_{\perp}) + \mathcal{U}(z_{\perp}, y_{\perp}) - \mathcal{U}(x_{\perp}, y_{\perp}) + \mathcal{U}(x_{\perp}, z_{\perp})\mathcal{U}(z_{\perp}, y_{\perp}) \right\},$$
(3)

where $\mathcal{U}(x_{\perp}, y_{\perp}) \equiv \frac{1}{N_c} \operatorname{Tr}\{U(x_{\perp})U^{\dagger}(y_{\perp})\} - 1$. The first three linear terms in braces in the r.h.s. of Eq. (3) reproduces the BFKL pomeron [1] while the quadratic term will give us the three-pomeron vertex [18]. The solution of the linearized evolution equation is especially simple in the case of zero momentum transfer (e.g., for the total cross section of small-*x* DIS):

$$\langle p | \mathcal{U}^{\zeta = x_B}(x_{\perp}, 0) | p \rangle = \int \frac{d\nu}{2\pi^2} (x_{\perp}^2)^{-\frac{1}{2} + i\nu} \left(\frac{s}{m^2}\right)^{\omega(\nu)} \int dz_{\perp}(z_{\perp}^2)^{-\frac{1}{2} - i\nu} \langle p | \mathcal{U}^{\zeta_0}(z_{\perp}, 0) | p \rangle, \tag{4}$$

where $\omega(v) = 2N_c \frac{\alpha_s}{\pi} [-\operatorname{Re} \psi(\frac{1}{2} + iv) - C]$ and m^2 is either Q^2 or m_N^2 (in LLA, we cannot distinguish between $\alpha_s \ln \frac{s}{Q^2}$ and $\alpha_s \ln \frac{s}{m_N^2}$). The sketch of linear evolution is presented in Fig. 1. The starting point of the evolution is the slope collinear to the momentum of the incoming photon q ($\zeta = x_B$) and it is convenient to stop the evolution at a certain intermediate point $\zeta_0 = Q^2/s_0$, where $s_0 \gg m_N^2$, $\frac{\alpha_s}{\pi} \ln \frac{s_0}{m_N^2} \ll 1$. The first of these conditions means that s_0 is still high from the viewpoint of low-energy nucleon physics while the second condition means that s_0 is sufficiently small from the viewpoint of high-energy physics (so one can neglect the BFKL logs). The matrix element of the double-Wilson-line operator at this slope is a phenomenological input for the BFKL evolution (just as the structure function at low Q^2 serves as the input for ordinary DGLAP evolution). At large *s* the integral over v is dominated by the vicinity of v = 0 which gives the familiar BFKL asymptotics $\sigma^{\text{tot}} \simeq x_B^{-12\frac{\alpha_s}{\pi} \ln 2}$.

Unlike the linear evolution, the general picture is very complicated since the number of operators U and U^{\dagger} increases after each evolution. At the time being, it is not known how to solve the non-linear evolution equation in an explicit form. It is possible, however, to write down the solution of the non-linear equation (3) in the form of a functional integral over the double set of the variables, $\varsigma_{i=1,2}(z_{\perp}, \eta) = t^a \varsigma_{i=1,2}^a(z_{\perp}, \eta)$ belonging to the Lie algebra of the SU(3) color group and $\Omega_{i=1,2}(z_{\perp}, \eta)$ belonging to the group itself:

$$U^{\eta_A}(x_{\perp}) \otimes U^{\dagger \eta_A}(y_{\perp}) = \int_{\Omega_{1,2}(\eta_0)=1}^{\pi_{1,2}(\eta_A)=0} D_{\varsigma_1}(z,\eta) D_{\varsigma_2}(z,\eta) D\Omega_1(z,\eta) D\Omega_2(z,\eta)$$

² The first non-linear equation for parton densities is known since 1983 as the GLR equation (it was conjectured in Ref. [7] and proved in the double-log limit in Ref. [8]). The full LLA x result was first derived in Ref. [2] by the above method. After that, it was reobtained in Ref. [11] in the framework of the dipole model [15,16], in Ref. [17] by direct summation of relevant Feynman diagrams, and in Refs. [19,20] by the semiclassical methods.

$$\times \Omega_{1}^{\dagger}(x_{\perp},\eta_{A})U_{x}^{\eta_{0}}\Omega_{2}(x_{\perp},\eta_{A}) \otimes \Omega_{2}^{\dagger}(y_{\perp},\eta_{A})U_{y}^{\dagger\eta_{0}}\Omega_{1}(y_{\perp},\eta_{A})$$

$$\times \exp\left\{\int_{\eta_{0}}^{\eta_{A}}d\eta \int d^{2}z \left[\frac{1}{g}\sum_{i=1,2}\varsigma_{i}^{a}(z,\eta)\vec{\partial}^{2}\left(\Omega_{i}^{\dagger}(z,\eta)\dot{\Omega}_{i}(z,\eta)\right)^{a} -\frac{1}{4\pi}\varsigma_{1}^{a}(z,\eta)\varsigma_{2}^{b}(z,\eta)\vec{\partial}^{2}\left(\Omega_{1}^{\dagger}(z,\eta)U_{z}^{\eta_{0}}\Omega_{2}(z,\eta)\right)^{ab}\right]\right\},$$

$$(5)$$

where $\dot{\Omega} \equiv \partial \Omega / \partial \eta$ and $(\Omega^{\dagger} \dot{\Omega})^a \equiv 2 \operatorname{Tr} \{ t^a \Omega^{\dagger} \dot{\Omega} \}$. Going to the variables $\pi = \vec{\partial}_{\perp}^2 \varsigma$ we see that Eq. (5) is a phase-space functional integral for the non-local Hamiltonian

$$\widehat{H}(\pi_1, \pi_2, \Omega_1, \Omega_2) = \int dx_{\perp} dy_{\perp} \pi_1^a(x_{\perp}) ((x_{\perp} | \frac{1}{\vec{p}_{\perp}^2} [\vec{\partial}_{\perp}^2 (\Omega_1^{\dagger} \Omega_2)^{ab}] \frac{1}{\vec{p}_{\perp}^2} |y_{\perp})) \pi_2^b(y_{\perp}),$$
(6)

where $|x\rangle$ is an eigenstate of the coordinate operator normalized according to $((x|y)) = \delta^{(2)}(x - y)$, see, e.g, Ref. [21]. The rapidity η serves as a Euclidean "time" for this evolution.

We shall demonstrate that the perturbative expansion of the functional integral (5) reproduces the evolution of the color dipole $U(x_{\perp}) \otimes U^{\dagger}(y_{\perp})$ in the LLA. To get the perturbative series, we substitute $\Omega(x_{\perp}, \eta) = e^{-ig\phi(x_{\perp}, \eta)}$:

$$U_{x}^{\eta_{A}} \otimes U_{y}^{\dagger\eta_{A}} = \int_{\phi_{1,2}(\eta_{0})=0}^{\pi_{1,2}(\eta_{A})=0} \prod_{i=1,2} D_{\varsigma_{i}}(z,\eta) D\phi_{i}(z,\eta) \\ \times e^{ig\phi_{1}(x_{\perp},\eta_{A})} U_{x}^{\eta_{0}} e^{-ig\phi_{2}(x_{\perp},\eta_{A})} \otimes e^{ig\phi_{2}(y_{\perp},\eta_{A})} U_{y}^{\dagger\eta_{0}} e^{-ig\phi_{1}(y_{\perp},\eta_{A})} \\ \times \exp\Biggl\{ \int_{\eta_{0}}^{\eta_{A}} d\eta \int d^{2}z \Biggl[\frac{1}{g} \sum_{i=1,2} \varsigma_{i}^{a}(z,\eta) \vec{\partial}^{2} \Biggl(e^{ig\phi_{i}(z,\eta)} \frac{\partial}{\partial \eta} e^{-ig\phi_{i}(z,\eta)} \Biggr)^{a} \\ - \frac{1}{4\pi} \varsigma_{1}^{a}(z,\eta) \varsigma_{2}^{b}(z,\eta) \vec{\partial}^{2} \Bigl(e^{ig\phi_{1}(z,\eta)} U_{z}^{\eta_{0}} e^{-ig\phi_{2}(z,\eta)} \Biggr)^{ab} \Biggr] \Biggr\}.$$
(7)

Next, we can represent the r.h.s. of Eq. (7) in the form $(\dot{\phi} \equiv \partial \phi / \partial \eta)$

$$\int_{\phi_{1,2}(\eta_{0})=0}^{\pi_{1,2}(\eta_{A})=0} \prod_{i=1,2} D_{\varsigma_{i}}(z,\eta) D\phi_{i}(z,\eta) [\eta_{A},\eta_{0}]_{x} U_{x}^{\eta_{0}}\{\eta_{0},\eta_{A}\}_{x} \otimes \{\eta_{A},\eta_{0}\}_{y} U_{y}^{\dagger\eta_{0}}[\eta_{0},\eta_{A}]_{y} \\ \times \exp\left\{\int_{\eta_{0}}^{\eta_{A}} d\eta \int d^{2}z \left[-i \sum_{i=1,2} \varsigma_{i}^{a}(z,\eta) \vec{\partial}^{2} \dot{\phi}_{i}(z,\eta) - \frac{1}{4\pi} \varsigma_{1}^{a}(z,\eta) \vec{\partial}^{2} \left([\eta_{A},\eta_{0}]_{z} U_{z}^{\eta_{0}}\{\eta_{0},\eta_{A}\}_{z}\right)^{ab} \varsigma_{2}^{b}(z,\eta)\right]\right\},$$

$$(8)$$

where we introduced the notations

$$[\eta_1, \eta_2]_x \equiv \mathrm{T}e^{ig\int_{\eta_2}^{\eta_1} \dot{\phi}_1(x_\perp, \eta)}, \qquad \{\eta_1, \eta_2\}_x \equiv \mathrm{T}e^{ig\int_{\eta_2}^{\eta_1} \dot{\phi}_2(x_\perp, \eta)}.$$
(9)

Let us now expand the r.h.s. of Eq. (8) in powers of g. The first nontrivial term in this expansion is

$$\begin{aligned} U_{x}^{\eta_{A}} \otimes U_{y}^{\dagger\eta_{A}} \\ &= \alpha_{s} \int_{\phi_{1,2}(\eta_{0})=0}^{\pi_{1,2}(\eta_{A})=0} \prod_{i=1,2} D_{\varsigma_{i}}(z,\eta) D\phi_{i}(z,\eta) \\ &\times \left[\left(\phi_{1}(x_{\perp},\eta_{A}) U_{x}^{\eta_{0}} - U_{x}^{\eta_{0}} \phi_{2}(x_{\perp},\eta_{A}) \right) \otimes \left(\phi_{2}(y_{\perp},\eta_{A}) U_{y}^{\dagger\eta_{0}} - U_{y}^{\dagger\eta_{0}} \phi_{1}(y_{\perp},\eta_{A}) \right) \\ &- \left(\phi_{1}(x_{\perp},\eta_{0}) U_{x}^{\eta_{0}} \phi_{2}(x_{\perp},\eta_{A}) \right) \otimes U_{y}^{\dagger\eta_{0}} + U_{x}^{\eta_{0}} \otimes \left(\phi_{2}(y_{\perp},\eta_{A}) U_{y}^{\dagger\eta_{0}} \phi_{1}(y_{\perp},\eta_{A}) \right) \right] \\ &\times \int_{\eta_{0}}^{\eta_{A}} d\eta \int d^{2}z \, \varsigma_{1}^{a}(z,\eta) \, \varsigma_{2}^{b}(z,\eta) \, \vec{\partial}^{2} \left(U^{\eta_{0}}(z_{\perp}) \right)^{ab} \exp \left\{ -i \int_{\eta_{0}}^{\eta_{A}} d\eta \int d^{2}z \, \sum_{i=1,2}^{z} \varsigma_{i}^{a}(z,\eta) \, \vec{\partial}^{2} \dot{\phi}_{i}^{a}(z,\eta) \right\}. \end{aligned}$$
(10)

The propagators for this phase-space functional integral are

$$\left\langle \phi_i^a(x,\eta)\varsigma_j^b(y,\eta') \right\rangle = i\delta_{ij}\delta^{ab}((x|\frac{1}{\vec{p}^2}|y))\theta(\eta-\eta'), \left\langle \phi_i^a(x,\eta)\phi_j^b(y,\eta') \right\rangle = 0, \qquad \left\langle \varsigma_i^a(x,\eta)\varsigma_j^b(y,\eta') \right\rangle = 0.$$

$$(11)$$

With these propagators, the r.h.s of Eq. (10) reduces to

$$-\alpha_{s}(\eta_{A}-\eta_{0})\bigg[\Big(t^{a}U_{x}^{\eta_{0}}\otimes t^{b}U_{y}^{\dagger\eta_{0}}+U_{x}^{\eta_{0}}t^{b}\otimes U_{y}^{\dagger\eta_{0}}t^{a}\Big)((x|\frac{1}{\vec{p}^{2}}\vec{\partial}^{2}U^{\eta_{0}}\frac{1}{\vec{p}^{2}}|y))^{ab} \\ -t^{a}U_{x}^{\eta_{0}}t^{b}\otimes U_{y}^{\dagger\eta_{0}}((x|\frac{1}{\vec{p}^{2}}\big(\vec{\partial}^{2}U^{\eta_{0}}\big)\frac{1}{\vec{p}^{2}}|x))^{ab} - U_{x}^{\eta_{0}}\otimes t^{b}U_{y}^{\dagger\eta_{0}}t^{a}((y|\frac{1}{\vec{p}^{2}}\big(\vec{\partial}^{2}U^{\eta_{0}}\big)\frac{1}{\vec{p}^{2}}|y))^{ab}\bigg],$$

$$(12)$$

which coincides with the Eq. (B17) from Ref. [2]. Taking trace over the color dipole indices one reproduces the Eq. (3). Similarly, it can be demonstrated that further terms of the expansion of Eq. (8) in powers of g reproduce the subsequent iterations of the non-linear equation (3).

The integral over π variables can be easily performed resulting in:

$$U^{\eta_{A}}(x_{\perp}) \otimes U^{\dagger\eta_{A}}(y_{\perp})$$

$$= \int_{\Omega_{1,2}(\eta_{0})=1}^{\dot{\Omega}_{1,2}(\eta_{A})=0} D\Omega_{1}(z,\eta) D\Omega_{2}(z,\eta) \Omega_{1}^{\dagger}(x_{\perp},\eta_{A}) U^{\eta_{0}}(x_{\perp}) \Omega_{2}(x_{\perp},\eta_{A}) \otimes \Omega_{2}^{\dagger}(y_{\perp},\eta_{A}) U^{\dagger\eta_{0}}(y_{\perp}) \Omega_{1}(y_{\perp},\eta_{A})$$

$$\times \exp\left\{-\frac{1}{\alpha_{s}} \int_{\eta_{0}}^{\eta_{A}} d\eta \int d^{2}z \left[\vec{\partial}^{2} \left(\Omega_{2}^{\dagger}(z,\eta) U_{z}^{\dagger\eta_{0}} \Omega_{1}(z,\eta)\right)\right]_{ab}^{-1} \\ \times \vec{\partial}^{2} \left(i \Omega_{1}^{\dagger}(z,\eta) \dot{\Omega}_{1}(z,\eta)\right)^{a} \vec{\partial}^{2} \left(i \Omega_{2}^{\dagger}(z,\eta) \dot{\Omega}_{2}(z,\eta)\right)^{b}\right\}.$$
(13)

Note that the action of this effective field theory is local. This functional integral for the small-x evolution of the Wilson-line operators is the main result of the Letter.

In the case of large nuclei it is possible to write initial conditions for the small-*x* evolution using the McLerran– Venugolalan model. The nuclear matrix element of the two-Wilson-line operator ("color dipole") is given by the Glauber formula [22–24], see Fig. 2.

$$\int d^2 z_{\perp} \langle A | \operatorname{Tr} U(x_{\perp} + z_{\perp}) U(z_{\perp}) | A \rangle = N_c \int d^2 b \left[1 - e^{-g^2 c_F \mathcal{G}(x_{\perp}^2) L_b} \right].$$
⁽¹⁴⁾



Fig. 2. Propagation of the color dipole through the nucleus.

Here $L_b \equiv 2\sqrt{R^2 - b^2}$ is the propagation length of the dipole (located at the impact parameter *b*) through the nucleus, $\rho = \frac{A}{4/3\pi R^3}$ is the nuclear density, and

$$\mathcal{G}(x_{\perp}^{2}) \equiv \frac{\pi x_{\perp}^{2}}{4(N_{c}^{2}-1)} \rho \sigma_{0} G\left(\sigma_{0}, \mu^{2} = \frac{1}{x_{\perp}^{2}}\right).$$
(15)

The Eq. (15) is derived under the assumption that the characteristic size of the dipole (the "saturation scale") is smaller than the size of the nucleon.³ In this case, the quarks propagating along the straight light-like lines⁴ interact by the instantaneous (in the light-cone time x_+) potential

$$\rho g^{2} t^{a} \otimes t^{a} \int \frac{d^{2} p_{\perp}}{(2\pi)^{2}} \frac{g^{2}}{2p_{\perp}^{4}} \left(e^{i(p,x-y)_{\perp}} - 1 \right) = g^{2} t^{a} \otimes t^{a} \rho \frac{\alpha_{s}}{8} (x-y)_{\perp}^{2} \ln(x-y)_{\perp}^{2} m_{0}^{2}, \tag{16}$$

where $m_0 \simeq m_N/2$ is the IR cutoff [23]. It is worth noting that the factor -1 in the parenthesis in the l.h.s. comes from the diagrams with the two gluons attached to the same nucleon and the same Wilson line. Taking into account the color factors, one obtains the Eq. (14) with $x_B G(x_B, \mu^2 = x_{\perp}^{-2}) = \frac{4\alpha_s}{\pi} \ln x_{\perp}^{-2}/m_0^2$, see Ref. [23]. Similarly to Eq. (13), it is possible to represent this result as a functional integral over a variable $\Lambda(x_{\perp}, l) \in$

SU(3):

$$\int d^{2}z \langle A | U_{x+z}^{\eta_{0}} U_{z}^{\dagger \eta_{0}} | A \rangle$$

$$= \int d^{2}b \int_{\Lambda(0,y)=1}^{\Lambda'(L_{b},y)=0} D\Lambda(y,l) \Lambda(x+z,L_{b}) \Lambda^{\dagger}(z,L_{b})$$

$$\times \exp\left\{\frac{1}{2g^{2}\rho} \int_{0}^{L_{b}} dl \int d^{2}y \left(\Lambda(l,y)\Lambda'(l,y)\right)^{a} \left(-\vec{\partial}^{2} + m_{0}^{2}\right)^{2} \left(\Lambda(l,y)\Lambda'(l,y)\right)^{a}\right\},$$
(17)

where $\Lambda' \equiv \partial \Lambda / \partial l$. Extra $U^{\eta_0}(x)$ $(U^{\dagger \eta_0}(x))$ lead to extra $\Lambda(x, L_b)$ $(\Lambda^{\dagger}(x, L_b))$ in the pre-exponent.

³ This assumption is certainly true at $A \to \infty$. For real nuclei, one should find the saturation scale Q_{δ} from the final result for the matrix element of the color dipole between the nuclear states, and verify that $Q_s \gg 1$ GeV.

⁴ As we mentioned above, the energy $s\sigma_0$ should be high enough so we can replace the slope $p_1 + \zeta_0 p_2$ by p_1 in the non-logarithmical expressions.

The final formula for the matrix element of the color dipole operator at small x_B is obtained by combining the functional integrals (13) and (17):

$$\begin{cases}
d^{2}z \langle A | U^{\eta_{A}}(x+z) \otimes U^{\dagger\eta_{A}}(z) | A \rangle \\
= \int d^{2}b \int_{\Lambda(0,y_{\perp})=1}^{\Lambda'(L_{b},y_{\perp})=0} \int_{\Omega_{1,2}(\eta_{0},y)=1}^{\Omega_{1,2}(\eta_{A},y)=0} D\Lambda(l,y) D\Omega_{1}(y,\eta) D\Omega_{2}(y,\eta) \Omega_{1}^{\dagger}(x+z,\eta_{A}) \Lambda(L_{b},x+z) \\
\times \otimes \Omega_{2}(x+z,\eta_{A}) \Omega_{2}^{\dagger}(z,\eta_{A}) \Lambda^{\dagger}(L_{b},z) \Omega_{1}(z,\eta_{A}) \\
\times \exp\left\{-\frac{1}{2g^{2}\rho} \int_{0}^{L_{b}} dl \int d^{2}y \left(i\Lambda(l,y)\Lambda'(l,y)\right)^{a} \left(m_{0}^{2}-\vec{\partial}^{2}\right)^{2} \left(i\Lambda(l,y_{\perp})\Lambda'(l,y)\right)^{a} \\
-\frac{1}{\alpha_{s}} \int_{\eta_{0}}^{\eta_{A}} d\eta \int d^{2}y \vec{\partial}^{2} \left(i\Omega_{1}^{\dagger}(y,\eta)\dot{\Omega}_{1}(y,\eta)\right)^{a} \left[\vec{\partial}^{2} \left(\Omega_{2}^{\dagger}(y,\eta)\Lambda^{\dagger}(L_{b},y)\Omega_{1}(y,\eta)\right)\right]_{ab}^{-1} \\
\times \vec{\partial}^{2} \left(i\Omega_{2}^{\dagger}(y,\eta)\dot{\Omega}_{2}(y,\eta)\right)^{b} \right\}.$$
(18)

The gluon structure function in the LLA is proportional to the matrix element of the dipole operator $x_B G(x_B, \mu^2 = x_{\perp}^{-2}) = -\frac{2\pi}{s} \langle A | \operatorname{Tr} U_i^{\eta_A}(x_{\perp}) U_i^{\eta_A}(0) | A \rangle$, so the numerical calculation of the functional integral (18) should give the nuclear structure functions at small *x*. This would be complementary to the approximate solutions of Refs. [11,12, 17,19,20,26] since it could give the structure functions not only in the asymptotic black-body limit, but also in the intermediate region defining the saturation scale Q_s .

It should be mentioned that our formula (13) gives the evolution of the color dipole only in the LLA. In the case of large nucleus we have an additional parameter $A \gg 1$ so our LLA approximation based on the non-linear equation (3) has a window $\alpha_s^2 A^{1/3} \sim 1$, $\alpha_s \ln x_B \sim 1$ where it is justified even at moderately small x_B . In the case of nucleon, our $\alpha_s(Q_s) \ll 1$, $\alpha_s(Q_s) \ln x_B \sim 1$ approximation should be justified *a posteriori* after checking that the saturation does occur at sufficiently small x_B . If the saturation takes place at such low x that $\alpha_s(Q_s) \ln x_B \gg 1$, our LLA breaks down and we need to take into account the non-fan diagrams such as t-channel loops formed by BFKL pomerons. However, the non-linear equation (3) leads to the result for the structure function which does not violate unitarity (see the discussion in Refs. [11–13,17,25]) and therefore we should not expect the large discrepancy between the unitary LLA result and the exact amplitude at present energies.

Acknowledgements

The author is grateful to Y.V. Kovchegov, E.M. Levin and L. McLerran for valuable discussions. This work was supported by the US Department of Energy under contract DE-AC05-84ER40150.

References

- [1] V.S. Fadin, E.A. Kuraev, L.N. Lipatov, Phys. Lett. B 60 (1975) 50;
- I. Balitsky, L.N. Lipatov, Sov. J. Nucl. Phys. 28 (1978) 822.
- [2] I. Balitsky, Nucl. Phys. B 463 (1996) 99.
- [3] L.N. Lipatov, Phys. Rep. 286 (1997) 131;

I. Balitsky, High-energy QCD and Wilson lines, in: M. Shifman (Ed.), Boris Ioffe Festschrift "Handbook on QCD", World Scientific, Singapore, 2001, hep-ph/0101042.

- [4] O. Nachtmann, Ann. Phys. 209 (1991) 436;
- J.C. Collins, R.K. Ellis, Nucl. Phys. B 360 (1991) 3. [5] V.S. Fadin, L.N. Lipatov, Phys. Lett. B 429 (1998) 127;
- G. Carnici, M. Ciafaloni, Phys. Lett. B 430 (1998) 349.
- [6] Y.V. Kovchegov, A.H. Mueller, Phys. Lett. B 439 (1998) 428.
- [7] L.V. Gribov, E.M. Levin, M.G. Ryskin, Phys. Rep. 100 (1983) 1.
- [8] A.H. Mueller, J.W. Qiu, Nucl. Phys. B 268 (1986) 427.
- [9] L. McLerran, R. Venugopalan, Phys. Rev. D 49 (1994) 2233;
 L. McLerran, R. Venugopalan, Phys. Rev. D 49 (1994) 3352;
 L. McLerran, R. Venugopalan, Phys. Rev. D 50 (1994) 2225.
- [10] A.H. Mueller, Nucl. Phys. B 558 (1999) 285.
- [11] Yu.V. Kovchegov, Phys. Rev. D 60 (1999) 034008;
 Yu.V. Kovchegov, Phys. Rev. D 61 (2000) 074018.
- [12] E. Levin, K. Tuchin, Nucl. Phys. B 573 (2000) 833.
- J. Jalilian-Marian, A. Kovner, L. McLerran, H. Weigert, Nucl. Phys. B 504 (1997) 415;
 J. Jalilian-Marian, A. Kovner, L. McLerran, H. Weigert, Phys. Rev. D 59 (1999) 014014;
 J. Jalilian-Marian, A. Kovner, L. McLerran, H. Weigert, Phys. Rev. D 59 (1999) 034007.
- [14] I. Balitsky, Phys. Rev. Lett. 81 (1998) 2024, hep-ph/9808215;
 I. Balitsky, Phys. Rev. D 60 (1999) 014020.
- [15] A.H. Mueller, Nucl. Phys. B 415 (1994) 373;
 A.H. Mueller, B. Patel, Nucl. Phys. B 425 (1994) 471.
- [16] N.N. Nikolaev, B.G. Zakharov, Phys. Lett. B 332 (1994) 184;
 N.N. Nikolaev, B.G. Zakharov, Z. Phys. C 64 (1994) 631.
- [17] M.A. Braun, Eur. Phys. J. C 16 (2000) 337.
- [18] J. Bartels, M. Wusthoff, Z. Phys. C 66 (1995) 157.
- [19] E. Iancu, L. McLerran, Preprint SACLAY-T01-026, hep-ph/0103032.
- [20] H. Weigert, Preprint NORDITA-2000-34-HE, hep-ph/0004044.
- [21] I. Balitsky, V.M. Braun, Nucl. Phys. B 311 (1989) 541.
- [22] B.Z. Kopeliovich, I.L. Lapidus, Al.B. Zamolodchikov, JETP Lett. 33 (1981) 612.
- [23] A.H. Mueller, Nucl. Phys. B 335 (1990) 115.
- [24] Yu.V. Kovchegov, A.H. Mueller, Nucl. Phys. B 529 (1998) 451.
- [25] A.H. Mueller, Nucl. Phys. B 437 (1995) 107.
- [26] E. Levin, K. Tuchin, Preprint TAUP-2659-2000, hep-ph/0012167;
 E. Levin, K. Tuchin, Preprint TAUP-2664-2001, hep-ph/0101275.