

Old Dominion University

ODU Digital Commons

Community & Environmental Health Faculty
Publications

Community & Environmental Health

2011

Self-Reported Poor Oral Hygiene Among In-School Adolescents in Zambia

Seter Siziya

Adamson S. Muula

Emmanuel Rudatsikira
Old Dominion University

Follow this and additional works at: https://digitalcommons.odu.edu/commhealth_fac_pubs



Part of the [Dental Public Health and Education Commons](#), and the [International Public Health Commons](#)

Original Publication Citation

Siziya, S., Muula, A. S., & Rudatsikira, E. (2011). Self-reported poor oral hygiene among in-school adolescents in Zambia. *BMC Research Notes*, 4(1), 1-4, Article 255. <https://doi.org/10.1186/1756-0500-4-255>

This Article is brought to you for free and open access by the Community & Environmental Health at ODU Digital Commons. It has been accepted for inclusion in Community & Environmental Health Faculty Publications by an authorized administrator of ODU Digital Commons. For more information, please contact digitalcommons@odu.edu.

SHORT REPORT

Open Access

Self-reported poor oral hygiene among in-school adolescents in Zambia

Seter Siziya¹, Adamson S Muula^{2*} and Emmanuel Rudatsikira³

Abstract

Background: Dental health is a neglected aspect of adolescent health globally but more so in low-income countries. Secondary analysis using the 2004 Zambia Global School-Based Health Survey (GSHS) was conducted in which we estimated frequencies of relevant socio-demographic variables and explored associations between selected explanatory variables and self-reported poor oral hygiene (not cleaning or brushing teeth) within the last 30 days of the completion of questionnaire.

Findings: Most of the 2257 respondents were males (53.9%) and went hungry (82.5%). More than 4 in 10 respondents drank alcohol (42.2%) while 37.2% smoked cannabis. Overall 10.0% of the respondents reported to have poor oral hygiene. Male respondents were 7% less likely to report to have poor oral hygiene compared to females. Compared to respondents who never drank alcohol, those who drank alcohol were 27% more likely to report to have poor oral hygiene. Respondents who smoked cannabis were 4% more likely to report to have poor oral hygiene compared to those who did not smoke cannabis. Finally, respondents who went hungry were 35% more likely to report to have poor oral hygiene compared to those who did not go hungry.

Conclusions: Results from this study indicate that female gender, alcohol drinking, cannabis smoking, and going hungry were associated with self-reported poor oral hygiene. The identification of these factors should guide the design and implementation of programs aimed to improve oral health among adolescents.

Background

Dental health has received limited attention in many countries in sub-Saharan Africa. Despite the paucity of data of dental health in most African countries, some reports are worth highlighting. Folayan et al have reported that DMFT (decayed, missing and filled teeth) was negatively associated with age but positively associated with sugar intake in a sample of Nigerian children [1]. In a study of South African children, 52% of the children with early childhood caries (ECC) were reported not to have been supervised during tooth brushing [2]. Masanja and Mumghamba have reported that, in Tanzania, secondary school adolescents aged 13 to 17 years had partial knowledge about gingivitis but a good knowledge of the basic oral hygiene measures necessary to maintain proper oral health [3]. About 99% of the study participants perceived that it was necessary to brush their teeth on a daily basis.

There are limited published reports on oral health in Zambia. Noar and Portnoy have suggested that the changing social and economic situation for most families, resulting in high sugar intake in Zambia were influencing the epidemiology of dental conditions in the country [4]. Meanwhile, Dickson highlighted the reduced prevalence of dental caries in western province of Zambia partly due to almost no availability of refined sugar and use of wood ash to clean teeth [5]. Baboo et al have corroborated this finding and reported that urbanization and confectionary eating were contributing to the increased dental pathology among children in Zambia [6]. Tooth brushing is the primary recommended practice for maintaining dental hygiene [7]. It is, therefore, important that persons with poor oral hygiene are identified and interventions instituted to improve their oral health.

We used data from the Zambia Global School-Based Health Survey (GSHS) conducted in 2004 to estimate the prevalence of self-reported poor oral hygiene and assess the socio-demographic factors that were associated with the practice.

* Correspondence: amuula@medcol.mw

²Department of Community Health, University of Malawi, College of Medicine, Blantyre, Malawi

Full list of author information is available at the end of the article

Methods

Details of the Zambia GSHS of 2004, especially with regard to the sampling strategy, have been published elsewhere [8,9]. A two-stage sampling technique was used to recruit students mainly aged 13 to 15 years in schools. In the first stage of sampling, schools with grades having 13-15 years olds were included in a sampling frame. Each school's chance of being selected was proportional to its enrollment size. In the second stage of sampling, a systematic sample of classes within each school was obtained. The survey achieved a response rate of 75%.

Both the Ministries of Health and Education provided ethical oversight of the Zambia GSHS. Although eligible students were encouraged to participate, they were also informed that they were free not to participate or decide not to answer any of the questions they felt uncomfortable with. Questionnaires, designed in a multiple choice fashion, were self-completed in class by the students within one school period i.e. about 40-50 minutes. The questionnaire has been used in over 50 countries, and as such it has been validated. No follow-up sessions were conducted for students who were absent from school on the day the questionnaire was administered.

The main question was: "During the past 30 days, how many times per day did you usually clean or brush your teeth?" Students were given the option of replying: I did not clean or brush my teeth during the past 30 days; 1 time per day; 2 times per day; 3 times per day; 4 or more times per day. All respondents who reported "I did not clean or brush my teeth during the past 30 days" were coded "1" i.e. did not brush teeth, while those who reported at least 1 time were coded as "0" i.e. positive history of reporting ever brushed teeth in the past 30 days. The category of no cleaning or brushing teeth was termed as self-reported poor oral hygiene. No clinical parameters or indices of plaque or gingival status were collected.

The Statistical Package for the Social Sciences (SPSS) version 11.5 (Chicago Illinois, United States) was used for the analysis. We obtained frequencies and proportions of the outcome variable (self-reported poor oral hygiene) and some key variables. The Pearson's Chi-square test was used to determine associations between the explanatory variables and the outcome variable during bivariate analyses. The cut off point for statistical significance was set at the 5% level. A Backward logistic regression analysis was conducted to estimate the magnitudes of associations (Adjusted Odds Ratios [AOR] with 95% Confidence Intervals [CI]) between the outcome variable and the following explanatory variables that were identified in the literature to be significantly associated with poor oral hygiene and collected in the survey: sex, age, drank alcohol, smoked cannabis, and went hungry. The questions were: During

the past 30 days, on how many days did you have at least one drink containing alcohol?; During your life, how many times have you used drugs cannabis?; During the past 30 days, how often did you go hungry because there was not enough food in your home? The last question was used as an indirect measure of poverty or household socio-economic status. No other socio-economic variables such as parental income, number of siblings and housing status were collected.

Description of the sample for the 2004 Zambia sample

A total of 2257 respondents took part in the study. Most of the respondents were male (53.9%), drank alcohol (42.2%), ever smoked cannabis (37.2%), and went hungry (82.5%). Most of the male respondents were aged 16 years or more (36.5%), whilst most female respondents were 13 years olds or younger (30.9%). Overall 10.0% of the respondents reported to have poor oral hygiene. The description of the sample is shown in Table 1.

Correlates for self-reported poor oral hygiene among Zambian school-going adolescents

Table 2 shows results from bivariate analyses. All the factors (age, sex, drank alcohol, smoked cannabis and went hungry) were significantly associated with self-

Table 1 Description of the sample for the 2004 Zambian Global School Health Survey

Factor	Total n* (%)**	Male n* (%)**	Female n* (%)**
Age			
< 14	463 (27.5)	156 (21.5)	263 (30.9)
14	386 (19.1)	156 (17.4)	219 (21.6)
15	513 (22.8)	260 (24.5)	238 (21.6)
16+	708 (30.6)	394 (36.5)	306 (25.9)
Sex			
Male	994 (53.9)	-	-
Female	1039 (46.1)	-	-
Drank alcohol			
Yes	528 (42.2)	217 (38.5)	283 (45.1)
No	805 (57.8)	388 (61.50)	388 (54.9)
Smoked cannabis			
Yes	695 (37.2)	300 (34.5)	357 (39.4)
No	1301 (62.8)	634 (65.5)	614 (60.6)
Went hungry			
Yes	1691 (82.5)	787 (82.3)	834 (83.4)
No	366 (17.5)	173 (17.7)	170 (16.6)
Self reported poor oral hygiene			
Yes	186 (10.0)	81 (10.3)	96 (9.8)
No	1891 (90.0)	884 (89.7)	920 (90.2)

* unweighted frequency

** weighted percent

Numbers may not add up due to missing information.

Table 2 Factors associated with self-reported poor oral hygiene in bivariate analyses

Factor	Self-reported poor oral hygiene		p value
	No* (%)**	Yes* (%)**	
Age			
< 14	58 (37.2)	392 (26.5)	< 0.001
14	34 (18.8)	341 (19.2)	
15	30 (13.8)	470 (23.9)	
16+	61 (30.2)	627 (30.4)	
Sex			
Male	81 (54.8)	884 (53.6)	< 0.001
Female	96 (45.2)	920 (46.4)	
Drank Alcohol			
Yes	55 (59.0)	459 (40.7)	< 0.001
No	48 (41.0)	739 (59.3)	
Smoked cannabis			
Yes	82 (49.4)	590 (35.5)	< 0.001
No	93 (50.6)	1179 (64.5)	
Went hungry			
Yes	154 (87.6)	1485 (81.7)	< 0.001
No	23 (12.4)	339 (18.3)	

* unweighted frequency

** weighted percent

Numbers may not add up due to missing information.

reported poor oral hygiene. Respondents reporting poor oral hygiene tended to be younger than 14 years, males, drank alcohol, smoked cannabis and went hungry.

The results of multivariate analysis are shown in Table 3. Although age remained significantly associated with self-reported poor oral hygiene during the multivariate analysis, it was not consistently associated with the outcome. While respondents who were aged less than 14 years (AOR = 0.74, 95%CI [0.72, 0.77]) and those of age 15 years (AOR = 0.74, 95%CI [0.71, 0.76]) were less likely to report to have poor oral hygiene, those of age 14 years (AOR = 1.51, 95% CI [1.47, 1.56]) were more likely to report to have poor oral hygiene compared to respondents aged 16 years or older. Male respondents were 7% (AOR = 0.93, 95%CI [0.91, 0.94]) less likely to report to have poor oral hygiene compared to female respondents. Compared to respondents who never drank alcohol, those who drank alcohol were 27% (AOR = 1.27, 95%CI [1.25, 1.30]) more likely to report to have poor oral hygiene. Respondents who smoked cannabis were 4% (AOR = 1.04, 95%CI [1.02, 1.07]) more likely to report to have poor oral hygiene. Finally, adolescents who went hungry were 35% (AOR = 1.35, 95%CI [1.31, 1.39]) more likely to report to have poor oral hygiene compared to the respondents who did not go hungry. These results are shown in Table 3.

Discussion

In a survey of in-school adolescents in Zambia in 2004, 10.0% reported to have poor oral hygiene in the 30 days

Table 3 Correlates for self reported poor oral hygiene among Zambian school-going adolescents

Factor	AOR* (95%CI**)
Age	
< 14	0.74 (0.72, 0.77)
14	1.51 (1.47, 1.56)
15	0.74 (0.71, 0.76)
16+	1
Sex	
Male	0.93 (0.91, 0.94)
Female	1
Drank alcohol	
Yes	1.27 (1.25, 1.30)
No	1
Smoked cannabis	
Yes	1.04 (1.02, 1.07)
No	1
Went hungry	
Yes	1.35 (1.31, 1.39)
No	1

AOR* Adjusted Odds Ratio

CI** Confidence Interval.

preceding the survey. In multivariable logistic regression analysis, age, sex, substance use (alcohol, cannabis) and going hungry were significantly associated with self-reported poor oral hygiene.

Although age was significantly associated with self-reported poor oral hygiene, the results from this study do not suggest a specific pattern as only the 14 years olds group had higher odds of not brushing while the other age groups had lower odds compared to 16 years of age or older. We offer no explanation for this finding.

Although, girls reported significantly more frequent optimal oral hygiene practice than boys in Australia [10] and Kenya [11], on the contrary, female respondents in our study were more likely to report to have poor oral hygiene. However, unhealthy lifestyles such as substance use was associated with self-reported poor dental hygiene, highlighting that multiple unhealthy practices may converge within the same individual [9]. Finally, if we assumed that having gone hungry because of lack of food in the household is some measure of poverty, it would suggest, from our results that poverty was positively associated with self-reported poor oral hygiene. This finding i.e. that adolescents from poor households are less likely to attend to their dental hygiene or have poor dental health has also been reported by other authors [12,13].

This study has a number of limitations. Being a cross sectional design, we may not be able to assign causation to any of the factors. To the extent that survey participants mis-reported, our results may be biased. We have no specific information to suggest that survey participants

may have mis-reported or not on any of the variables. There were some missing data, and these were declared missing in the analysis. The relatively low response rate may have biased our results. However, we do not have data on the non-respondents that we may use to compute the magnitude and direction of the bias. Furthermore, only adolescents in school were eligible for recruitment. The findings may therefore be applicable to adolescents who are in school but not drop-outs and those who have never been in school. Finally, no clinical parameters or indices of plaque or gingival status were collected.

Conclusion

Results from this study indicate that female gender, alcohol drinking, cannabis smoking, and going hungry were positively associated with self-reported poor oral hygiene. The identification of these factors should guide the design and implementation of school and community-based programs aimed to improve oral health among adolescents. Some of the programs may include educational and awareness outreach through media and youth social networks.

Acknowledgements

We thank the World Health Organization for making these data available to us for analysis. The WHO however had no part in deciding which analyses to run and the decision to publish the findings. Grateful thanks to the students and the survey teams who made the Zambia 2004 Global School Based Survey possible.

Author details

¹Department of Community Medicine, University of Zambia, Medical School, Lusaka, Zambia. ²Department of Community Health, University of Malawi, College of Medicine, Blantyre, Malawi. ³School of Community and Environmental Health, Old Dominion University, Norfolk, Virginia, USA.

Authors' contributions

SS led data analysis and interpretation, ER contributed to the interpretation of the data and ASM led the drafting of the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 25 January 2011 Accepted: 22 July 2011

Published: 22 July 2011

References

1. Folayan M, Sowole A, Kola-Jebutu A: Risk factors for caries in children from south-western Nigeria. *J Clin Pediatr Dent* 2008, **32**:171-5.
2. Mohamed N, Barnes J: Characteristics of children under 6 years of age treated for early childhood caries in South Africa. *J Clin Pediatr Dent* 2008, **32**:247-52.
3. Masanja IM, Mumghamba EG: Knowledge on gingivitis and oral hygiene practices among secondary school adolescents in rural and urban Morogoro, Tanzania. *Int J Dent Hyg* 2004, **2**:172-8.
4. Noar J, Portnoy S: Dental status of children in a primary and secondary school in rural Zambia. *Int Dent J* 1991, **41**:142-8.
5. Dickson ID: Preventing dental caries. *Br Med J* 1968, **1**:579.
6. Baboo KS, Bhatia IK, Nkowane BS: A study of dental caries in rural and an urban primary school of Zambia. *Med J Zambia* 1981, **15**:56-9.
7. Claydon NC: Current concepts in toothbrushing and interdental cleaning. *Periodontol 2000* 2008, **48**:10-22.

8. Muula AS, Kazembe LN, Rudatsikira E, Siziya S: Suicidal ideation and associated factors among in-school adolescents in Zambia. *Tanzan Health Res Bull* 2007, **9**:202-6.
9. Siziya S, Muula AS, Kazembe LN, Rudatsikira E: Harmful lifestyles' clustering among sexually active in-school adolescents in Zambia. *BMC Pediatr* 2008, **8**:6.
10. Tran D, Phongsavan P, Bauman AE, Havea D, Galea G: Hygiene behaviour of adolescents in the Pacific: associations with socio-demographic, health behaviour and school environment. *Asia Pac J Public Health* 2006, **18**:3-11.
11. Okemwa KA, Gatongi PM, Rotich JK: The oral health knowledge and oral hygiene practices among primary school children age 5-17 years in a rural area of Uasin Gishu district, Kenya. *East Afr J Public Health* 2010, **7**:187-90.
12. Cinar AB, Kosku N, Sandalli N, Murtomaa H: Individual and maternal determinants of self-reported dental health among Turkish school children aged 10-12 years. *Community Dent Health* 2008, **25**(2):84-8.
13. Livny A, Assali R, Sgan-Cohen HD: Early Childhood Caries among a Bedouin community residing in the eastern outskirts of Jerusalem. *BMC Public Health* 2007, **7**:167.

doi:10.1186/1756-0500-4-255

Cite this article as: Siziya et al.: Self-reported poor oral hygiene among in-school adolescents in Zambia. *BMC Research Notes* 2011 **4**:255.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

