

2004

Increased Risk for Aplastic Anemia and Myelodysplastic Syndrome in Individuals Lacking Glutathione S-Transferase Genes

Joanne F. Sutton

Michael Stacey

Old Dominion University, mstacey@odu.edu


William G. Kearns

Thomas S. Roeg

Neal S. Young

See next page for additional authors

Follow this and additional works at: https://digitalcommons.odu.edu/bioelectrics_pubs

 Part of the [Hematology Commons](#), [Oncology Commons](#), and the [Pediatrics Commons](#)

Repository Citation

Sutton, Joanne F.; Stacey, Michael; Kearns, William G.; Roeg, Thomas S.; Young, Neal S.; and Liu, Johnson M., "Increased Risk for Aplastic Anemia and Myelodysplastic Syndrome in Individuals Lacking Glutathione S-Transferase Genes" (2004). *Bioelectrics Publications*. 212.

https://digitalcommons.odu.edu/bioelectrics_pubs/212

Original Publication Citation

Sutton, J. F., Stacey, M., Kearns, W. G., Rieg, T. S., Young, N. S., & Liu, J. M. (2004). Increased risk for aplastic anemia and myelodysplastic syndrome in individuals lacking glutathione s-transferase genes. *Pediatric Blood Cancer*, 42(2), 122-126. doi:10.1002/pbc.10479

Authors

Joanne F. Sutton, Michael Stacey, William G. Kearns, Thomas S. Roeg, Neal S. Young, and Johnson M. Liu

Increased Risk for Aplastic Anemia and Myelodysplastic Syndrome in Individuals Lacking Glutathione S-Transferase Genes

Joanne F. Sutton, MD,^{1,2,3} Michael Stacey, PhD,⁴ William G. Kearns, PhD,¹ Thomas S. Rieg, PhD,³ Neal S. Young, MD,¹ and Johnson M. Liu, MD^{1*}

Background. Aplastic anemia (AA) and myelodysplastic syndrome (MDS) are marrow failure states that may be associated with chromosomal instability. An absence of the glutathione S-transferase (GST) enzyme may genetically predispose individuals to AA or MDS. **Procedure and Results.** To test this hypothesis, we determined the *GSTM1* and *GSTT1* genotypes in a total of 196 patients using multiplex PCR. The *GSTT1* null genotype was found to be overrepresented in Caucasian, Asian, and Hispanic patients with either AA or MDS. We confirmed a difference in the expected frequency of the *GSTM1* null genotype in

Caucasian MDS patients. The double null *GSTM1/GSTT1* genotype was also overrepresented in Caucasian AA and MDS patients. In our population, 26% of AA patients and 40% of MDS patients had a chromosomal abnormality identified by karyotype or FISH analyses for chromosomes 7 and 8. Patients with AA and the *GSTT1* null genotype had an increased frequency of chromosomal abnormalities ($P=0.003$). **Conclusion.** There seems to be an increased risk for AA and MDS in individuals lacking *GSTT1* or both *GSTM1/GSTT1*. *Pediatr Blood Cancer* 2004;42:122–126. Published 2003 Wiley-Liss, Inc.[†]

Key words: aplastic anemia; chromosomal abnormality; genotype; glutathione S-transferase; myelodysplastic syndrome

BACKGROUND

Acquired aplastic anemia (AA) and myelodysplastic syndrome (MDS) are distinct but related bone marrow failure syndromes, diagnosed by marrow morphology. Approximately 20% of MDS cases present with a hypocellular marrow and, in turn, AA may evolve to MDS or develop cytogenetic abnormalities [1]. For each disorder, the etiology is unknown, however, there may be gene-environment interactions that predispose to disease [1,2]. For example, an increased incidence of AA has been observed in Asia [3]. Glutathione S-transferase (GST) is a key biometabolic enzyme for a number of chemical toxins, including the organophosphates, alkylating agents, epoxides, and polycyclic aromatic hydrocarbons [4]. Low GST activity may genetically predispose some individuals to marrow failure, perhaps as a result of increased susceptibility to endogenous or environmental toxins.

Glutathione S-transferase M1 (*GSTM1*) and T1 (*GSTT1*) genes are polymorphic in humans. It has been well established that the frequencies of *GSTT1* null and *GSTM1* null genotypes in healthy human subjects vary greatly among different ethnic populations and can reach frequencies as high as 0.58 (*GSTT1* null) and 0.67 (*GSTM1* null) in Asians [5–10]. Racial or ethnic variation in bone marrow failure risk may reflect innate biological susceptibility as well as differences in environmental exposures or socioeconomic and demographic factors [1,11].

Recently, increased frequencies of *GSTM1* and *GSTT1* gene deletions were reported in 57 Korean patients with AA [12]. Similarly, a higher risk for MDS was noted in individuals with *GSTT1* gene deletions, although subsequent studies of patients with different ethnicities yielded conflicting results [13–19]. Here, we have determined the frequency of homozygous *GSTM1* and *GSTT1* gene deletions in an ethnically heterogeneous cohort of AA and MDS patients referred to the National Institutes of Health. We also expanded our study to determine if there

¹Hematology Branch, NHLBI, Bethesda, Maryland

²Department of Pediatrics, Uniformed Services University of the Health Sciences, Bethesda, Maryland

³Naval Medical Center Portsmouth, Portsmouth, Virginia

⁴Center for Pediatric Research, Eastern Virginia Medical School, Norfolk, Virginia

Joanne F. Sutton and Michael Stacey contributed equally to the work.

Grant sponsor: Division of Intramural Research, NHLBI, NIH.

*Correspondence to: Johnson M. Liu, Division of Hematology/Oncology, Mount Sinai School of Medicine, One Gustave L. Levy Place, Box 1079, New York, NY 10029.

E-mail: Johnson.Liu@mssm.edu

Received 25 June 2003; Accepted 13 November 2003

Published 2003 Wiley-Liss, Inc.

[†]This article is a US government work and, as such, is in the public domain of the United States of America.

DOI 10.1002/pbc.10479

Published online 15 December 2003 in Wiley InterScience (www.interscience.wiley.com)

was a difference in the frequency of chromosomal abnormalities in patients with *GSTM1* or *GSTT1* null genotypes compared to those with *GSTM1* or *GSTT1* present.

STUDY DESIGN AND METHODS

The study population included 196 patients, 108 diagnosed with either moderate or severe AA and 88 with MDS, referred to the National Heart Lung and Blood Institute, Hematology Branch between 1997 and 2000 (age range, 3–85 years; median age, 41 years; 112 males and 84 females). Standard diagnostic criteria were utilized, and no patients with congenital bone marrow failure (Fanconi anemia) were included. Race was self-reported by each patient (142 Caucasian, 28 Hispanic, 16 Black, and 10 Asian). To assure patient anonymity, no other patient identifiers were used. Cytogenetic data by conventional karyotype of the bone marrow were available for 185 (94%), and a bone marrow pellet was available for 113 (57%). Control values were based on selecting the median value of previously published gene frequencies of healthy human subjects sorted by race [5–10].

DNA was isolated from bone marrow samples using Qiagen tissue prep kit according to the manufacturer's guidelines (Qiagen, Valencia, CA). Analysis for the *GSTM1* and *GSTT1* genes was performed by multiplex PCR, as previously described [5]. In brief, 30 ng of isolated DNA was analyzed in a single assay for *GSTM1*, *GSTT1*, and *CYP1A1*. All primers were custom synthesized at Life Technologies (Invitrogen, Corp, Rockville, MD). Primers were: *GSTM1* (5'-GAACTCCCTGAAAAGCTAAA-GC-3') and (5'-GTTGGGCTCAAATATACGGTGG-3'); *GSTT1* (5'-TTCCTTACTGGTCCTCACATCTC-3') and (5'-TCACCGGATCATGGCCAGCA-3'); *CYP1A1* (5'-GAACTGCCACTTCAGCTGTCT-3') and (5'-CAGCTGCATTTGGAAGTGCTC-3'). A 50 μ l reaction mixture containing 30 pmol of each primer, 200 μ mol dNTPs, 5 μ l of 10 \times PCR buffer, 2 U Taq DNA polymerase (Qiagen PCR Core Kit) was amplified in a PTC-200 Peltier Thermal Cycler. PCR conditions were as follows: initial melting 94°C \times 5 min, followed by 35 cycles of melting 94°C \times 2 min, annealing 59°C \times 1 min, extension 72°C \times 1 min, and a final extension at 72°C \times 10 min. The PCR products were analyzed on an ethidium bromide stained 2% agarose gel. The presence or absence of *GSTT1* and *GSTM1* genes was determined by examination for characteristic bands at 480 bp (*GSTT1*) and 215 bp (*GSTM1*). A band at 312 bp corresponding to the internal control (*CYP1A1*) was used to confirm successful PCR amplification.

Fluorescent in situ hybridization (FISH) was performed on 113 patients with available bone marrow pellets and 14 normal bone marrow controls, using alpha satellite DNA probes for chromosomes 7 and 8 (VYSIS, Inc., Downer's Grove, IL). A probe mixture was applied to the

target area and codenatured at 75°C for 3 min followed by hybridization at 42°C. Approximately 300 interphase nuclei per slide were scored for monosomy 7 and trisomy 8 as per criteria established by Vysis. Aneuploidy was established for monosomy 7 as $>2.91\%$ and for trisomy 8 as $>2.8\%$ of cells, based on results obtained in the normal samples.

Descriptive analyses included gene frequencies and contingency tables. Associations between variables were tested with chi-square and Fisher exact test where appropriate. Standard odds ratio and power were computed. Significance was obtained for $P < 0.05$.

RESULTS AND DISCUSSION

Frequency of *GSTM1* and *GSTT1* Null Genotypes in Bone Marrow Failure Patients Stratified by Race

Control values were selected (see ref. [8], expected frequencies, Table I) for each genotype based on the median value of previously published gene frequencies of healthy human subjects sorted by race. A recent study reported metabolic gene polymorphism frequencies for over 15,000 control subjects. Significant differences were observed between races, as expected, but much less heterogeneity was observed between White populations from different countries and no differences were seen by age, sex, or type of controls (hospital patients vs. population controls).

The *GSTT1* null genotype was overrepresented in Caucasian, Hispanic, and Asian patients with either AA or MDS (Table I). In contrast, the *GSTM1* null genotype was significantly overrepresented only in Caucasian MDS patients. No significant association was found for the null genotypes in Black patients. The double null genotype was significantly overrepresented only in Caucasians with either AA or MDS.

Data on the *GSTM1* and *GSTT1* null genotypes in marrow failure patients had previously been reported in relatively homogeneous populations, many of which did not include Black or Hispanic patients. For example, an increased frequency of the *GSTT1* null genotype was reported in White American and Japanese MDS patients [13,19]. Subsequent studies were unable to confirm this result in British, French, and Japanese populations [14–17]. Overrepresentation of the *GSTM1* null genotype was also reported in Greek MDS patients [18]. Acquired AA has not been as extensively examined for GST gene deletions. A study of 57 Korean AA patients reported a significant increase in frequency of the *GSTM1* or *GSTT1* null genotypes [12], however, a Brazilian study of 37 AA patients found no difference [20]. Ethnic variation in frequency of GST polymorphisms and different environmental exposures may account for the lack of concordance in the literature. Our referral population was ethnically and geographically diverse, and we have

TABLE I. Frequency of *GSTM1* and *GSTT1* Null Genotype Stratified by Race and Diagnosis

	<i>GSTM1</i> null							<i>GSTT1</i> null							<i>GSTM1/GSTT1</i> null						
	Total	No.	Odds ratio	Exp freq	Freq	P-value	Power	No.	Odds ratio	Exp freq	Freq	P-value	Power	No.	Odds ratio	Exp freq	Freq	P-value	Power		
Caucasian																					
Total	142	91	1.78	0.52	0.64	<0.003	0.095	47	0.49	0.15	0.33	<0.001	0.789	34	0.32	0.07	0.24	<0.001	0.998		
AA	69	40	1.38	0.52	0.58	0.28		21	0.44	0.15	0.30	<0.001	0.265	15	0.28	0.07	0.22	<0.001	0.724		
MDS	73	51	2.32	0.52	0.70	0.002	0.163	26	0.55	0.15	0.36	<0.001	0.723	19	0.35	0.07	0.26	<0.001	0.995		
Black																					
Total	16	4	0.33	0.32	0.25	0.58		6	0.60	0.24	0.38	0.21		2	0.14	0.04	0.13				
AA	13	2	0.18	0.32	0.15	0.21		5	0.63	0.24	0.39	0.22		1	0.08	0.04	0.076				
MDS	3 ^a	2	2.00	0.32	0.66	0.19		1	0.50	0.24	0.33	0.051		1	0.50	0.04	0.33	0.008	0.957		
Hispanic																					
Total	28	13	0.87	0.51	0.46	0.63		8	0.40	0.1	0.29	<0.001	0.473	4 ^a	0.17	NA	0.14				
AA	18	8	0.80	0.51	0.44	0.57		5	0.38	0.1	0.28	0.01	0.274	3 ^a	0.20	NA	0.17				
MDS	10	5	1.00	0.51	0.50	0.95		3	0.43	0.1	0.30	0.03	0.246	1 ^a	NA	NA	0.1				
Asian																					
Total	10	6	1.50	0.55	0.60	0.75		8	4.00	0.48	0.80	0.043	0.195	4	0.67	0.25	0.4	0.27			
AA	8	5	1.67	0.55	0.63	0.67		6	3.00	0.48	0.75	0.13		3	0.60	0.25	0.38	0.41			
MDS	2 ^a	1	1.00	0.55	0.50	NA		2		0.48	1.0	NA		1		0.25	0.5				

AA, aplastic anemia; MDS, myelodysplastic syndrome; Exp freq, expected frequency; NA, not available.

^aNumber of patients too low to determine statistical significance.

confirmed a difference in expected frequency of the *GSTT1* null genotype and AA or MDS in White and Hispanic patients. However, unlike the Korean study, we found no significant difference in frequency of the *GSTM1* null genotype and AA.

A striking difference in frequency of the double null *GSTM1/GSTT1* genotype in Caucasian AA and MDS patients was also observed. The double null genotype is an important variable to investigate because nullizyosity for *GSTM1* and *GSTT1* has been linked independently to various cancers [4], and deletion of both genes may have different significance from that of either gene alone.

High Frequency of Chromosomal Abnormalities Seen in AA Patients With *GSTT1* Null Genotype

Of the 187 patients with cytogenetics available, 26 of 102 (26%) patients with AA and 34 of 85 (40%) patients with MDS had a chromosomal abnormality identified by either conventional karyotype or FISH (Table II). Of those patients with bone marrow pellets available, FISH analyses identified monosomy 7 or trisomy 8 in 26 (44%) AA patients, of whom 18 had no chromosomal abnormalities identified by conventional karyotype. FISH analyses identified 27 (50%) MDS patients with either monosomy 7 or trisomy 8, of whom 8 had a normal conventional karyotype. The karyotype of each patient was not known at the time that FISH was performed, in order to prevent bias in scoring.

FISH analysis is known to be more sensitive than conventional cytogenetics. In this setting, FISH was limited to determining aneuploidy of chromosomes 7 and 8, which are two frequent chromosomal abnormalities identified in MDS and AA [1,2]. Among the MDS patients, there was no significant correlation between the *GSTM1* null and *GSTT1* null genotypes and chromosomal abnormalities. However, a significant difference in frequency of the *GSTT1* null genotype was observed in

AA patients with chromosomal abnormalities, consistent with findings from a Korean study [12]. The etiology of chromosomal aberrations in MDS cannot be correlated to a deletion of either the *GSTT1* or *GSTM1* gene. However, it is possible that the *GSTT1* null genotype decreases an individual's ability to biometabolize toxins, which may lead to an increased risk for chromosomal aberrations in AA.

ACKNOWLEDGMENT

We thank Kathleen L. McManaman (United States Department of Veterans Affairs, Washington, DC) and Dean Follmann (National Institutes of Health, Bethesda, MD) for their assistance in the statistical analyses. The Chief, Navy Bureau of Medicine and Surgery, Washington, DC, Clinical Investigation Program sponsored LCDR Joanne F. Sutton, MC, USNR. The views expressed in this study are those of the authors and do not reflect the official policy or position of the Department of the Navy, Department of Defense, or the United States Government.

REFERENCES

1. Young NS. Acquired aplastic anemia. *Ann Intern Med* 2002; 136:534–546.
2. Barrett J, Sauntharajah Y, Molldrem J. Myelodysplastic syndrome and aplastic anemia: Distinct entities or diseases linked by a common pathophysiology? *Semin Hematol* 2000;37:15–29.
3. Young NS. Acquired aplastic anemia. *JAMA* 1999;282: 271–278.
4. Hayes JD, Pulford DJ. The glutathione S-transferase supergene family: Regulation of GST and the contribution of the isoenzymes to cancer chemoprotection and drug resistance. *Crit Rev Biochem Mol Biol* 1995;30:445–600.
5. Abdel-Rahman SZ, El Zein RA, Anwar WA, et al. A multiplex PCR procedure for polymorphic analysis of *GSTM1* and *GSTT1* genes in population studies. *Cancer Lett* 1996;107: 229–233.
6. Au WW, Oh HY, Grady J, et al. Usefulness of genetic susceptibility and biomarkers for evaluation of environmental health risk. *Environ Mol Mutagen* 2001;37:215–225.
7. Chen CL, Liu Q, Relling MV. Simultaneous characterization of glutathione S-transferase M1 and T1 polymorphisms by polymerase chain reaction in American whites and blacks. *Pharmacogenetics* 1996;6:187–191.
8. Garte S, Gaspari L, Alexandrie AK, et al. Metabolic gene polymorphism frequencies in control populations. *Cancer Epidemiol Biomarkers Prev* 2001;10:1239–1248.
9. Nelson HH, Wiencke JK, Christiani DC, et al. Ethnic differences in the prevalence of the homozygous deleted genotype of glutathione S-transferase theta. *Carcinogenesis* 1995;16:1243–1245.
10. Rebbeck TR. Molecular epidemiology of the human glutathione S-transferase genotypes *GSTM1* and *GSTT1* in cancer susceptibility. *Cancer Epidemiol Biomarkers Prev* 1997;6:733–743.
11. Perera FP. Environment and cancer: Who are susceptible? *Science* 1997;278:1068–1073.
12. Lee KA, Kim SH, Woo HY, et al. Increased frequencies of glutathione S-transferase (*GSTM1* and *GSTT1*) gene deletions in Korean patients with acquired aplastic anemia. *Blood* 2001;98: 3483–3485.

TABLE II. Patient Karyotype Stratified by Genotype and Diagnosis

	<i>GSTM1</i> null	<i>GSTT1</i> null	<i>GSTM1</i> / <i>T1</i> null	<i>GSTM1</i> / <i>T1</i> present
AA				
NL karyotype ^a	38	22	14	32
Abn karyotype ^b	13	13	7	7
P-value (power)	0.160	0.003 (0.243)	0.066	
MDS				
NL karyotype ^a	36	17	11	12
Abn karyotype ^b	23	15	11	7
P-value	0.733	0.239	0.200	

AA, aplastic anemia; MDS, myelodysplastic syndrome; NL, normal karyotype; Abn, abnormal karyotype.

^aKaryotype from conventional and FISH analysis on bone marrow cells.

^bIncludes aneuploidy and structural defects.

13. Chen H, Sandler DP, Taylor JA, et al. Increased risk for myelodysplastic syndromes in individuals with glutathione transferase theta 1 (GSTT1) gene defect. *Lancet* 1996;347:295–297.
14. Basu T, Gale RE, Langabeer S, et al. Glutathione S-transferase theta 1 (GSTT1) gene defect in myelodysplasia and acute myeloid leukaemia. *Lancet* 1997;349:1450.
15. Atoyebi W, Kusec R, Fidler C, et al. Glutathione S-transferase gene deletions in myelodysplasia. *Lancet* 1997;349:1450–1451.
16. Preudhomme C, Nisse C, Hebbar M, et al. Glutathione S transferase theta 1 gene defects in myelodysplastic syndromes and their correlation with karyotype and exposure to potential carcinogens. *Leukemia* 1997;11:1580–1582.
17. Okada M, Okamoto T, Wada H, et al. Glutathione S-transferase theta 1 gene (GSTT1) defect in Japanese patients with myelodysplastic syndromes. *Int J Hematol* 1997;66:393–394.
18. Tsabouri SE, Georgiou I, Alamanos I, et al. Increased prevalence of GSTM(1) null genotype in patients with myelodysplastic syndrome: A case-control study. *Acta Haematol* 2000;104:169–173.
19. Sasai Y, Horiike S, Misawa S, et al. Genotype of glutathione S-transferase and other genetic configurations in myelodysplasia. *Leuk Res* 1999;23:975–981.
20. Arruda VR, Lima CS, Grignoli CR, et al. Increased risk for acute myeloid leukaemia in individuals with glutathione S-transferase mu 1 (GSTM1) and theta 1 (GSTT1) gene defects. *Eur J Haematol* 2001;66:383–388.