

9-2018

Asymmetric Relativistic Fermi Gas Model for Quasielastic Lepton-Nucleus Scattering

M. B. Barbaro

A. De Pace

T. W. Donnelly

J. A. Caballero

G. D. Megias

See next page for additional authors

Follow this and additional works at: https://digitalcommons.odu.edu/physics_fac_pubs

 Part of the [Nuclear Commons](#)

Repository Citation

Barbaro, M. B.; De Pace, A.; Donnelly, T. W.; Caballero, J. A.; Megias, G. D.; and Van Orden, J. W., "Asymmetric Relativistic Fermi Gas Model for Quasielastic Lepton-Nucleus Scattering" (2018). *Physics Faculty Publications*. 246.
https://digitalcommons.odu.edu/physics_fac_pubs/246

Original Publication Citation

Barbaro, M. B., De Pace, A., Donnelly, T. W., Caballero, J. A., Megias, G. D., & Van Orden, J. W. (2018). Asymmetric relativistic Fermi gas model for quasielastic lepton-nucleus scattering. *Physical Review C*, 98(3), 035501. doi:10.1103/PhysRevC.98.035501

Authors

M. B. Barbaro, A. De Pace, T. W. Donnelly, J. A. Caballero, G. D. Megias, and J. W. Van Orden

Asymmetric relativistic Fermi gas model for quasielastic lepton-nucleus scattering

M. B. Barbaro,^{1,2} A. De Pace,² T. W. Donnelly,³ J. A. Caballero,⁴ G. D. Megias,^{4,5} and J. W. Van Orden^{6,7}

¹*Dipartimento di Fisica, Università di Torino, Italy*

²*INFN, Sezione di Torino, Via P. Giuria 1, 10125 Torino, Italy*

³*Center for Theoretical Physics, Laboratory for Nuclear Science and Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

⁴*Departamento de Física Atómica, Molecular y Nuclear, Universidad de Sevilla, 41080 Sevilla, Spain*

⁵*IRFU, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France*

⁶*Department of Physics, Old Dominion University, Norfolk, Virginia 23529, USA*

⁷*Jefferson Laboratory, 12000 Jefferson Avenue, Newport News, Virginia 23606, USA*



(Received 22 June 2018; published 4 September 2018)

We develop an asymmetric relativistic Fermi gas model for the study of the electroweak nuclear response in the quasielastic region. The model takes into account the differences between neutron and proton densities in asymmetric ($N > Z$) nuclei, as well as differences in the neutron and proton separation energies. We present numerical results for both neutral and charged-current processes, focusing on nuclei of interest for ongoing and future neutrino oscillation experiments. We point out some important differences with respect to the commonly employed symmetric Fermi gas model.

DOI: [10.1103/PhysRevC.98.035501](https://doi.org/10.1103/PhysRevC.98.035501)

I. INTRODUCTION

The relativistic Fermi gas (RFG) is currently the main model used in the analysis of long baseline experiments, aiming at high-precision measurements of the neutrino oscillation parameters (see [1] for a recent review of the subject). Since the detectors of these experiments are made of complex nuclei, a good control of the nuclear effects is essential in order to achieve this goal. In particular, the next generation experiment DUNE [2] will be using liquid argon time-projection chamber technology; hence a reliable description of neutrino interactions with ^{40}Ar is needed.

Although the RFG is clearly inadequate to describe the details of the nuclear dynamics at the required level of precision, and several more refined calculations have been developed in recent years to provide a better modeling of neutrino-nucleus scattering [1,3–13], the model has some merits in providing the gross features of inclusive lepton-nucleus cross sections, which are the subject of the present work. First, it is fully relativistic, and relativity plays an essential role in the kinematical domain of present and future neutrino experiments, covering an energy range of few to several GeV. Second, in spite of its simplicity, the RFG is capable of predicting the behavior of the inclusive cross section as a function of the momentum transfer q and of the Fermi momentum k_F , the so-called scaling of first and second kind [14–16]. This feature is the basis of the SuSAv2 model [4,17,18], which has been proved to give an excellent description of both electron and neutrino cross sections on symmetric ($N = Z$) nuclei, in particular carbon and oxygen. The extension of the RFG model to asymmetric ($N \neq Z$) nuclei will provide the basis for the development of the SuSAv2 model for these nuclei.

With these motivations, in this work we develop the formalism of a fully relativistic Fermi gas model for asymmetric

nuclear matter and apply this to the study of quasielastic (QE) electron and neutrino scattering on a selected set of nuclei. A similar approach was taken in Refs. [19–23], however, using a nonrelativistic model, to study various response functions. Specifically, we focus on a typical $N = Z$ nucleus ^{12}C , on a slightly asymmetric nucleus ^{40}Ar , both of which are of practical interest for studies of neutrino oscillations [1], and on a very asymmetric case, ^{208}Pb . Neutral current electron and (anti)neutrino scattering results are given for these three nuclei, although other cases may be treated in a similar manner. For charge-changing neutrino and antineutrino reactions, of course, the neighboring nuclei $^{12}\text{B}/^{12}\text{N}$, $^{40}\text{Cl}/^{40}\text{K}$, and $^{208}\text{Tl}/^{208}\text{Bi}$, respectively, are also modeled using the same asymmetric Fermi gas approach. Here we focus on QE processes, although the formalism can easily be extended to include the one-particle–one-hole (1p1h) inelastic spectrum, namely, meson production, production of baryon resonances, and Deep Inelastic Scattering, following the developments in Ref. [24]. The 2p2h excitations will be treated in future work.

The paper is organized as follows. In Sec. II we introduce the formalism for the asymmetric relativistic Fermi gas (ARFG) model, both for neutral current (Sec. II A) and charged-current (Sec. II B) reactions. In Sec. III we present a selection of numerical results, again for neutral (Sec. III A) and charged-current (Sec. III B) reactions. Finally, in Sec. IV we draw our conclusions.

II. FORMALISM: ARFG MODEL

We consider a nucleus (N, Z) having N neutrons and Z protons, where N and Z need not be equal, and assume that the nuclear volume V for protons and neutrons is the same. Here we are assuming that the cases typically of interest in such

studies are in the valley of stability where the equal-volume assumption is a reasonable approximation (in the case of lead, for instance, this is known to be the case); cases far from the valley of stability where, for instance, neutron skins may come into play, go beyond the present modeling and are not considered here. Thus we can define two different Fermi spheres for protons and neutrons, with corresponding Fermi momenta

$$k_F^p(Z) = \left(\frac{3\pi^2 Z}{V} \right)^{1/3}, \quad k_F^n(N) = \left(\frac{3\pi^2 N}{V} \right)^{1/3}. \quad (1)$$

Defining the ratio between the two Fermi momenta

$$\rho_0 \equiv \frac{k_F^n(N)}{k_F^p(Z)} = \left(\frac{N}{Z} \right)^{1/3} \quad (2)$$

and the average Fermi momentum

$$k_F^0(N, Z) \equiv \frac{1}{N+Z} [Zk_F^p(Z) + Nk_F^n(N)], \quad (3)$$

we can write

$$k_F^n(N) = \rho_0 k_F^p(Z) = \frac{\rho_0(Z+N)}{Z+\rho_0 N} k_F^0(N, Z). \quad (4)$$

In the study of charged-current (CC) quasielastic neutrino (or antineutrino) scattering, where a neutron is converted into a proton (or vice versa), one also has to consider the neighboring nuclei, $(N+1, Z-1)$ and $(N-1, Z+1)$. The corresponding Fermi momenta for protons and neutrons can be written as

$$k_F^p(Z \pm 1) = k_F^p(Z) \left(1 \pm \frac{1}{Z} \right)^{1/3}, \quad (5)$$

$$k_F^n(N \pm 1) = k_F^n(N) \left(1 \pm \frac{1}{N} \right)^{1/3}. \quad (6)$$

In the RFG model a nucleon having 3-momentum $\mathbf{k}$ has on-shell energy

$$E^{p(n)}(k) = \sqrt{k^2 + m_{p(n)}^2}. \quad (7)$$

A well-known shortcoming of the RFG is that the model has states starting with 3-momentum equal to zero (energy equal to the proton or neutron mass) and going up to the Fermi levels. This corresponds to an unrealistic negative separation energy [25]. In fact, the Fermi levels in a bound nucleus are negative and have positive separation energies $S_n(N)$, $S_p(Z)$, $S_n(N+1)$, $S_p(Z-1)$, $S_n(N-1)$, and $S_p(Z+1)$, respectively, for the six cases of interest. In order to correct for this flaw, we shift the energies of the protons and neutrons in the triplet of nuclei using the following prescriptions:

$$\begin{aligned} H^n(N; k) &= E^n(k) - D^n(N), \\ H^p(Z; k) &= E^p(k) - D^p(Z), \\ H^n(N+1; k) &= E^n(k) - D^n(N+1), \\ H^p(Z-1; k) &= E^p(k) - D^p(Z-1), \\ H^n(N-1; k) &= E^n(k) - D^n(N-1), \\ H^p(Z+1; k) &= E^p(k) - D^p(Z+1), \end{aligned} \quad (8)$$

TABLE I. Neutron and proton separation energies (S_n , S_p) and Fermi momenta (k_F) used in this work.

$X(A, Z, N)$	S_n (MeV)	S_p (MeV)	k_F^0 (MeV/c)	k_F^n (MeV/c)	k_F^p (MeV/c)
C(12,6,6)	18.72	15.96	228	228	228
B(12,5,7)	3.37	14.10		240.02	214.56
N(12,7,5)	15.04	0.60		214.56	240.02
Ar(40,18,22)	9.87	12.53	241	248.23	232.17
Cl(40,17,23)	5.83	11.68		251.93	227.78
K(40,19,21)	7.80	7.58		244.41	236.39
Pb(208,82,126)	7.37	8.00	248	261.77	226.85
Tl(208,81,127)	3.79	7.55		262.46	225.92
Bi(208,83,125)	6.89	3.71		261.07	227.76

where the offsets are given by

$$\begin{aligned} D^n(N) &= E_F^n(N) + S_n(N), \\ D^p(Z) &= E_F^p(Z) + S_p(Z), \\ D^n(N+1) &= E_F^n(N+1) + S_n(N+1), \\ D^p(Z-1) &= E_F^p(Z-1) + S_p(Z-1), \\ D^n(N-1) &= E_F^n(N-1) + S_n(N-1), \\ D^p(Z+1) &= E_F^p(Z+1) + S_p(Z+1), \end{aligned} \quad (9)$$

and where the usual RFG Fermi energies are given by

$$\begin{aligned} E_F^n(N) &\equiv E^n(k_F^n(N)), \\ E_F^p(Z) &\equiv E^p(k_F^p(Z)), \\ E_F^n(N+1) &\equiv E^n(k_F^n(N+1)), \\ E_F^p(Z-1) &\equiv E^p(k_F^p(Z-1)), \\ E_F^n(N-1) &\equiv E^n(k_F^n(N-1)), \\ E_F^p(Z+1) &\equiv E^p(k_F^p(Z+1)). \end{aligned} \quad (10)$$

Clearly when at the true Fermi surfaces the energies in Eq. (8) become minus the separation energies. For instance, when $k = k_F^n(N)$ one has

$$\begin{aligned} H^n(N; k_F^n(N)) &= E^n(k_F^n(N)) - [E_F^n(N) + S_n(N)] \\ &= -S_n(N). \end{aligned} \quad (11)$$

In this work the values of the parameter $k_F^0(N, Z)$, in terms of which all the different Fermi momenta can be calculated, are taken from the superscaling analysis [16] of electron scattering data, while $S_{p,n}$ are the measured proton and neutron separation energies, taken from the ENSDF database [26]. The numerical values for the cases considered in this work are listed in Table I. Note that although not explicitly indicated in our notation, for sake of simplicity, all separation energies depend on both Z and N .

Within this model, denoted as asymmetric relativistic Fermi gas (ARFG), we can now calculate the quasielastic double differential cross section with respect to the outgoing lepton momentum k' and scattering angle Ω corresponding to inclusive electron scattering, (e, e'), neutral-current (NC)

neutrino and antineutrino scattering, (ν, ν') and $(\bar{\nu}, \bar{\nu}')$, and to charged-current (CC) neutrino and antineutrino scattering, (ν, μ^-) and $(\bar{\nu}, \mu^+)$. In the Rosenbluth decomposition these can be expressed as

$$\left(\frac{d^2\sigma}{d\Omega dk'}\right)^{(e,e')} = \sigma_M [v_L R_{p \rightarrow p}^{L,em} + v_L R_{n \rightarrow n}^{L,em} + v_T R_{p \rightarrow p}^{T,em} + v_T R_{n \rightarrow n}^{T,em}] \quad (12)$$

$$\left(\frac{d^2\sigma}{d\Omega dk'}\right)^{(\nu,\nu')} = \sigma_0^{(NC)} [v_L R_{n \rightarrow n}^{L,w} + v_T R_{n \rightarrow n}^{T,w} + v_{T'} R_{n \rightarrow n}^{T',w}] \quad (13)$$

$$\left(\frac{d^2\sigma}{d\Omega dk'}\right)^{(\bar{\nu},\bar{\nu}')} = \sigma_0^{(NC)} [v_L R_{p \rightarrow p}^{L,w} + v_T R_{p \rightarrow p}^{T,w} - v_{T'} R_{p \rightarrow p}^{T',w}] \quad (14)$$

$$\left(\frac{d^2\sigma}{d\Omega dk'}\right)^{(\nu,\mu^-)} = \sigma_0^{(CC)} [V_{CC} R_{n \rightarrow p}^{CC,w} + V_{CL} R_{p \rightarrow p}^{CL,w} + V_{LL} R_{n \rightarrow p}^{LL,w} + V_T R_{n \rightarrow p}^{T,w} + V_{T'} R_{n \rightarrow p}^{T',w}] \quad (15)$$

$$\left(\frac{d^2\sigma}{d\Omega dk'}\right)^{(\bar{\nu},\mu^+)} = \sigma_0^{(CC)} [V_{CC} R_{p \rightarrow n}^{CC,w} + V_{CL} R_{p \rightarrow n}^{CL,w} + V_{LL} R_{n \rightarrow n}^{LL,w} + V_T R_{p \rightarrow n}^{T,w} - V_{T'} R_{p \rightarrow n}^{T',w}], \quad (16)$$

where v_K and V_K are leptonic kinematic factors (see [27,28] for their explicit expressions), σ_M is the Mott cross section, and $\sigma_0^{(NC)}$, $\sigma_0^{(CC)}$ the corresponding elementary weak cross sections for NC and CC reactions, respectively. The response functions $R^K \equiv R^K(q, \omega)$, where the labels “em” and w stand for “electromagnetic” and “weak”, respectively, embody the nuclear structure and dynamics and are functions of the momentum q and energy ω transferred to the nucleus. They are related to the specific components of the corresponding hadronic tensor $W^{\mu\nu}$:

$$R_L \equiv R_{CC} = W^{00}, \quad (17)$$

$$R_{CL} = -\frac{1}{2}(W^{03} + W^{30}), \quad (18)$$

$$R_{LL} = W^{33}, \quad (19)$$

$$R_T = W^{11} + W^{22}, \quad (20)$$

$$R_{T'} = -\frac{i}{2}(W^{12} - W^{21}). \quad (21)$$

The general expression for the nuclear tensor in the ARFG model is

$$\begin{aligned} W_{i \rightarrow f}^{\mu\nu}(q, \omega) &= \frac{3m^2 \mathcal{N}}{4\pi (k_F^i)^3} \int d\mathbf{h} \frac{\theta(k_F^i - |\mathbf{h}|) \theta(|\mathbf{h} + \mathbf{q}| - k_F^f)}{H^i(\mathbf{h}) H^f(\mathbf{h} + \mathbf{q})} \\ &\times f_{i \rightarrow f}^{\mu\nu}(\mathbf{h}, \mathbf{h} + \mathbf{q}) \delta[H^f(\mathbf{h} + \mathbf{q}) - H^i(\mathbf{h}) - \omega], \quad (22) \end{aligned}$$

where the superscripts i and f refer to the initial and final nucleons, respectively, m and $\mathcal{N}$ are the appropriate mass and number of nucleons in the target nucleus, $H^{i,f}$ are the nucleon energies defined in Eqs. (8) and $f_{i \rightarrow f}^{\mu\nu}$ is the corresponding single-nucleon tensor.

In the following sections we shall derive the explicit expression of $W^{\mu\nu}$ for the reactions listed in Eqs. (12)–(16), distinguishing between the two cases of neutral and charged-current reactions.

A. Electron scattering and NC (anti)neutrino scattering

In the case of neutral-current processes, Eqs. (12)–(14), mediated by the exchange of a photon or a Z^0 boson, the energy-conserving delta-function appearing in Eq. (22) involves the difference between the *on-shell* particle (p) and hole (h) energies, where the hole is assumed to have 3-momentum $\mathbf{h}$, while the particle has 3-momentum $\mathbf{h} + \mathbf{q}$,

$$\begin{aligned} E^{n \rightarrow n}(N) &= H^n(N; \mathbf{h} + \mathbf{q}) - H^n(N; \mathbf{h}) \\ &= E^n(\mathbf{h} + \mathbf{q}) - E^n(\mathbf{h}), \quad (23) \end{aligned}$$

$$\begin{aligned} E^{p \rightarrow p}(Z) &= H^p(Z; \mathbf{h} + \mathbf{q}) - H^p(Z; \mathbf{h}) \\ &= E^p(\mathbf{h} + \mathbf{q}) - E^p(\mathbf{h}), \quad (24) \end{aligned}$$

since the $D^n(N)$ and $D^p(N)$ offsets cancel in the particle-hole energy differences; namely, the nucleon separation energies introduced in the ARFG model have no impact on the results for neutral-current processes. The only difference with respect to the usual RFG arises from the different values of k_F for protons and neutrons. The corresponding nuclear tensors are then

$$\begin{aligned} W_{p \rightarrow p}^{\mu\nu}(q, \omega) &= \frac{3m_p^2 Z}{4\pi [k_F^p(Z)]^3} \int d\mathbf{h} \frac{\theta(k_F^p(Z) - |\mathbf{h}|) \theta(|\mathbf{h} + \mathbf{q}| - k_F^p(Z))}{E^p(\mathbf{h}) E^p(\mathbf{h} + \mathbf{q})} \\ &\times f_{p \rightarrow p}^{\mu\nu}(\mathbf{h}, \mathbf{h} + \mathbf{q}) \delta[E^p(\mathbf{h} + \mathbf{q}) - E^p(\mathbf{h}) - \omega] \quad (25) \end{aligned}$$

for protons and

$$\begin{aligned} W_{n \rightarrow n}^{\mu\nu}(q, \omega) &= \frac{3m_n^2 N}{4\pi [k_F^n(N)]^3} \int d\mathbf{h} \frac{\theta(k_F^n(N) - |\mathbf{h}|) \theta(|\mathbf{h} + \mathbf{q}| - k_F^n(N))}{E^n(\mathbf{h}) E^n(\mathbf{h} + \mathbf{q})} \\ &\times f_{n \rightarrow n}^{\mu\nu}(\mathbf{h}, \mathbf{h} + \mathbf{q}) \delta[E^n(\mathbf{h} + \mathbf{q}) - E^n(\mathbf{h}) - \omega] \quad (26) \end{aligned}$$

for neutrons, where

$$\begin{aligned} f_{j \rightarrow j}^{\mu\nu} &= -w_{1,j}(\tau_j) \left(g^{\mu\nu} - \frac{Q^\mu Q^\nu}{Q^2} \right) + w_{2,j}(\tau_j) V_j^\mu V_j^\nu \\ &\quad - \frac{i}{m_n} w_{3,n}(\tau_n) \epsilon^{\mu\nu\rho\sigma} Q_\rho V_{\sigma,j} \quad (27) \end{aligned}$$

(with $j = p, n$) are the single-nucleon tensors, with

$$\begin{aligned} \tau_j &\equiv \frac{|Q^2|}{4m_j^2} \quad \text{and} \\ V_j^\mu &= \frac{1}{m_j} \left(P^\mu - \frac{P \cdot Q}{Q^2} Q^\mu \right) = \frac{1}{m_j} \left(P^\mu + \frac{1}{2} Q^\mu \right), \quad (28) \end{aligned}$$

having used the on-shell condition $\frac{P \cdot Q}{Q^2} = -\frac{1}{2}$.

For (e, e') the electromagnetic structure functions are

$$w_{1,j}(\tau) = \tau G_{M,j}^2(\tau), \quad (29)$$

$$w_{2,j}(\tau) = \frac{G_{E,j}^2(\tau) + \tau G_{M,j}^2(\tau)}{1 + \tau}, \quad (30)$$

and $w_3 = 0$, while for (ν, ν') and $(\bar{\nu}, \bar{\nu}')$ they are

$$w_{1,j}(\tau) = \tau \tilde{G}_{M,j}^2(\tau) + (1 + \tau) \tilde{G}_{A,j}^2(\tau), \quad (31)$$

$$w_{2,j}(\tau) = \frac{\tilde{G}_{E,j}^2(\tau) + \tau \tilde{G}_{M,j}^2(\tau)}{1 + \tau} + \tilde{G}_{A,j}^2(\tau), \quad (32)$$

$$w_{3,j}(\tau) = \tilde{G}_{M,j}(\tau) \tilde{G}_{A,j}(\tau), \quad (33)$$

with $\tau = \tau_{n,p}$ as is appropriate. By performing the angular integration in Eqs. (25) and (26) one gets

$$W_{p \rightarrow p}^{\mu\nu}(q, \omega) = \frac{3m_p^2 Z}{2[k_F^p(Z)]^3 q} \int_{E_0^p(Z)}^{E_F^p(Z)} dE [f_{p \rightarrow p}^{\mu\nu}]_{x=x_0^p(E)}, \quad (34)$$

$$W_{n \rightarrow n}^{\mu\nu}(q, \omega) = \frac{3m_n^2 N}{2[k_F^n(N)]^3 q} \int_{E_0^n(N)}^{E_F^n(N)} dE [f_{n \rightarrow n}^{\mu\nu}]_{x=x_0^n(E)}, \quad (35)$$

where

$$E_0^p(Z) = \max \left\{ E_F^p(Z) - \omega, \frac{q}{2} \sqrt{1 + \frac{1}{\tau_p}} - \frac{\omega}{2} \right\}, \quad (36)$$

$$E_0^n(N) = \max \left\{ E_F^n(N) - \omega, \frac{q}{2} \sqrt{1 + \frac{1}{\tau_n}} - \frac{\omega}{2} \right\}, \quad (37)$$

and

$$x_0^{n,p}(E) = \frac{\omega E - \frac{|Q^2|}{2}}{q \sqrt{E^2 - m_{n,p}^2}}. \quad (38)$$

Finally, the analytic integration over E yields

$$W_{p \rightarrow p}^{\mu\nu}(q, \omega) = \frac{3m_p^2 Z}{2[k_F(Z)]^3 q} [E_F^p(Z) - E_0^p(Z)] U_p^{\mu\nu}, \quad (39)$$

$$W_{n \rightarrow n}^{\mu\nu}(q, \omega) = \frac{3m_n^2 N}{2[k_F(N)]^3 q} [E_F^n(N) - E_0^n(N)] U_n^{\mu\nu}. \quad (40)$$

In particular, the relevant components for the calculation of the L , T , and T' responses turn out to be

$$U_j^{00} = \frac{\kappa^2}{\tau_j} [-w_{1,j}(\tau_j) + (1 + \tau_j) w_{2,j}(\tau_j) + w_{2,j}(\tau_j) \Delta_j], \quad (41)$$

$$U_j^{11} + U_j^{22} = 2w_{1,j}(\tau_j) + w_{2,j}(\tau_j) \Delta_j, \quad (42)$$

$$U_j^{12} = 2i \sqrt{\tau_j(1 + \tau_j)} w_{3,j}(\tau_j) (1 + \Delta_j'), \quad (43)$$

with

$$\Delta_j = \frac{\tau_p}{\kappa^2} \frac{1}{m_j^2} \left[\frac{1}{3} (E_F^{j2}(Z) + E_0^j(Z) E_F^j(Z) + E_0^{j2}(Z)) + \frac{\omega}{2} (E_F^j(Z) + E_0^j(Z)) + \frac{\omega^2}{4} \right] - 1 - \tau_j, \quad (44)$$

$$\Delta_j' = \frac{1}{\kappa} \sqrt{\frac{\tau_j}{1 + \tau_j}} \frac{1}{m_j} \left[\frac{\omega}{2} + \frac{1}{2} (E_F^j(Z) + E_0^j(Z)) \right] - 1. \quad (45)$$

The corresponding numerical results will be shown in Sec. III.

B. CC (anti)neutrino scattering

In the case of (anti)neutrino-induced CC reactions, where neutrons (protons) are converted into protons (neutrons) through the absorption of a W^+ (W^-) boson, the energy differences appearing in the delta function of Eq. (22) are

$$\begin{aligned} E^{n \rightarrow p}(N, Z; ph) &= H^p(Z + 1; p) - H^n(N; h) \\ &= [E^p(p) - E^n(h)] + \Delta D^{n \rightarrow p}(N, Z), \end{aligned} \quad (46)$$

with

$$\begin{aligned} \Delta D^{n \rightarrow p}(N, Z) &\equiv D^n(N) - D^p(Z + 1) \\ &= [E_F^n(N) - E_F^p(Z + 1)] \\ &\quad + [S_n(N) - S_p(Z + 1)], \end{aligned} \quad (47)$$

for CC neutrino reactions, and

$$\begin{aligned} E^{p \rightarrow n}(N, Z; k'k) &= H^n(N + 1; k') - H^p(Z; k) \\ &= [E^n(k') - E^p(k)] + \Delta D^{p \rightarrow n}(N, Z), \end{aligned} \quad (48)$$

with

$$\begin{aligned} \Delta D^{p \rightarrow n}(N, Z) &\equiv D^p(Z) - D^n(N + 1) \\ &= [E_F^p(Z) - E_F^n(N + 1)] \\ &\quad + [\varepsilon_s^p(Z) - \varepsilon_s^n(N + 1)], \end{aligned} \quad (49)$$

for CC antineutrino reactions. The numerical values for these energy offsets are given in Table II.

TABLE II. Energy offsets used in this work for CC neutrino ($D^{n \rightarrow p}$) and antineutrino ($D^{p \rightarrow n}$) scattering.

$X(A, Z, N)$	$D^{n \rightarrow p}$ (MeV)	$D^{p \rightarrow n}$ (MeV)
C(12,6,6)	15.21	9.68
Ar(40,18,22)	5.25	1.77
Pb(208,82,126)	12.24	-4.77

The corresponding nuclear tensors are then

$$W_{n \rightarrow p}^{\mu\nu}(q, \omega) = \frac{3m_n^2 N}{4\pi [k_F^n(N)]^3} \int d\mathbf{h} \frac{\theta(k_F^n(N) - |\mathbf{h}|) \theta(|\mathbf{h} + \mathbf{q}| - k_F^p(Z+1))}{E^n(\mathbf{h}) E^p(\mathbf{h} + \mathbf{q})} \times f_{n \rightarrow p}^{\mu\nu}(\mathbf{h}, \mathbf{h} + \mathbf{q}) \delta[E^p(\mathbf{h} + \mathbf{q}) - E^n(\mathbf{h}) + \Delta D^{n \rightarrow p}(N, Z) - \omega], \quad (50)$$

$$W_{p \rightarrow n}^{\mu\nu}(q, \omega) = \frac{3m_p^2 Z}{4\pi [k_F^p(Z)]^3} \int d\mathbf{h} \frac{\theta(k_F^p(Z) - |\mathbf{h}|) \theta(|\mathbf{h} + \mathbf{q}| - k_F^n(N-1))}{E^p(\mathbf{h}) E^n(\mathbf{h} + \mathbf{q})} \times f_{p \rightarrow n}^{\mu\nu}(\mathbf{h}, \mathbf{h} + \mathbf{q}) \delta[E^n(\mathbf{h} + \mathbf{q}) - E^p(\mathbf{h}) + \Delta D^{p \rightarrow n}(N, Z) - \omega], \quad (51)$$

for neutrino and antineutrino scattering, respectively, where the elementary isovector tensor $f_{n \rightarrow p}^{\mu\nu} = f_{p \rightarrow n}^{\mu\nu} \equiv f^{\mu\nu(1)}$ is

$$f^{\mu\nu(1)} = -w_1^{(1)}(\tau) \left(g^{\mu\nu} - \frac{Q^\mu Q^\nu}{Q^2} \right) + w_2^{(1)}(\tau) V^\mu V^\nu + u_1^{(1)}(\tau) \frac{Q^\mu Q^\nu}{Q^2} - \frac{i}{m} w_3^{(1)}(\tau) \epsilon^{\mu\nu\rho\sigma} Q_\rho V_\sigma, \quad (52)$$

and the structure functions w_i are the appropriate isovector ones:

$$w_1^{(1)}(\tau) = \tau [G_M^{(1)}(\tau)]^2 + (1 + \tau) [G_A^{(1)}(\tau)]^2, \quad (53)$$

$$w_2^{(1)}(\tau) = \frac{[G_E^{(1)}(\tau)]^2 + \tau [G_{M,i}^{(1)}(\tau)]^2}{1 + \tau} + [G_A^{(1)}(\tau)]^2, \quad (54)$$

$$u_1^{(1)}(\tau) = -[G_A^{(1)}(\tau)]^2, \quad (55)$$

$$w_3^{(1)}(\tau) = G_M^{(1)}(\tau) G_A^{(1)}(\tau), \quad (56)$$

where

$$G_A^{(1)}(\tau) = G_A^{(1)}(\tau) - \tau G_p^{(1)}(\tau). \quad (57)$$

In the case of CC ν reactions we set $m_n \cong m_p \cong m \equiv (m_n + m_p)/2$ and define a single dimensionless 4-momentum transfer $\tau \equiv |Q^2|/4m^2$.

Similar to what we did for the NC case, we perform the angular integral, obtaining

$$W_{n \rightarrow p}^{\mu\nu}(q, \omega) = \frac{3m_n^2 N}{2[k_F^n(N)]^3 q} \int_{E_0^{n \rightarrow p}(N)}^{E_F^n(N)} dE [f_{n \rightarrow p}^{\mu\nu}]_{x=x_0^{n \rightarrow p}(E)}, \quad (58)$$

$$W_{p \rightarrow n}^{\mu\nu}(q, \omega) = \frac{3m_p^2 Z}{2[k_F^p(Z)]^3 q} \int_{E_0^{p \rightarrow n}(Z)}^{E_F^p(Z)} dE [f_{p \rightarrow n}^{\mu\nu}]_{x=x_0^{p \rightarrow n}(E)}, \quad (59)$$

where now

$$x_0^{n \rightarrow p}(E) = \frac{\tilde{\omega}_{n \rightarrow p} E - \frac{|\tilde{Q}_{n \rightarrow p}^2|}{2}}{q \sqrt{E^2 - m_n^2}} \quad \text{and} \quad x_0^{p \rightarrow n}(E) = \frac{\tilde{\omega}_{p \rightarrow n} E - \frac{|\tilde{Q}_{p \rightarrow n}^2|}{2}}{q \sqrt{E^2 - m_p^2}}, \quad (60)$$

having defined

$$\tilde{\omega}_{n \rightarrow p} \equiv \omega - \Delta D^{n \rightarrow p}(N, Z) \quad \text{and} \quad \tilde{Q}_{n \rightarrow p}^2 \equiv \tilde{\omega}_{n \rightarrow p}^2 - q^2, \quad (61)$$

$$\tilde{\omega}_{p \rightarrow n} \equiv \omega - \Delta D^{p \rightarrow n}(N, Z) \quad \text{and} \quad \tilde{Q}_{p \rightarrow n}^2 \equiv \tilde{\omega}_{p \rightarrow n}^2 - q^2. \quad (62)$$

The lower limits of integration in Eqs. (58) and (59) are

$$E_0^{n \rightarrow p}(N) = \max \{ E_F^p(Z+1) - \tilde{\omega}_{n \rightarrow p}, \Gamma^{n \rightarrow p} \}, \quad (63)$$

$$E_0^{p \rightarrow n}(Z) = \max \{ E_F^n(N+1) - \tilde{\omega}_{p \rightarrow n}, \Gamma^{p \rightarrow n} \}, \quad (64)$$

with

$$\Gamma^{n \rightarrow p} = \frac{q}{2} \sqrt{1 + \frac{1}{\tilde{\tau}_{n \rightarrow p}}} - \frac{\tilde{\omega}_{n \rightarrow p}}{2} \quad \text{and} \quad \Gamma^{p \rightarrow n} = \frac{q}{2} \sqrt{1 + \frac{1}{\tilde{\tau}_{p \rightarrow n}}} - \frac{\tilde{\omega}_{p \rightarrow n}}{2}, \quad (65)$$

together with $\tilde{\tau}_{n \rightarrow p} \equiv \frac{|\tilde{Q}_{n \rightarrow p}^2|}{4m^2}$ and $\tilde{\tau}_{p \rightarrow n} \equiv \frac{|\tilde{Q}_{p \rightarrow n}^2|}{4m^2}$.

Although in principle also in this case, as in the NC one, it is possible to obtain fully analytic results, for practical purposes it is easier to perform the energy integral numerically. The corresponding results will be shown in the next section.




FIG. 1. Electromagnetic response functions of ^{40}Ar in the SRFG and ARFG for momentum transfer $q = 300$ (left column) and 800 (right column) MeV/c. The separate contributions of protons and neutrons are also displayed.

III. RESULTS

In this section we present and compare the nuclear response functions evaluated in the symmetric (SRFG) and asymmetric (ARFG) relativistic Fermi gas models.

As anticipated, in the case of neutral-current reactions, (e, e') , (ν, ν') , and $(\bar{\nu}, \bar{\nu}')$, the ARFG results differ from the SRFG ones only due to the different neutron and proton Fermi momenta. On the contrary, for charged-current reactions,

(ν, μ^-) and $(\bar{\nu}, \mu^+)$, the energy offsets related to the different separation energies in the initial and final nuclei also play a role.

A. Neutral-current reactions

In Figs. 1 and 2 we show the electromagnetic (e, e') longitudinal and transverse response functions for ^{40}Ar and




FIG. 2. Electromagnetic response functions of ^{208}Pb in the SRFG and ARFG. The separate contributions of protons and neutrons are also displayed.




FIG. 3. Weak NC response functions for ν - ^{40}Ar in the SRFG and ARFG. The separate contributions of protons and neutrons are also displayed.

^{208}Pb as functions of the energy transfer ω and two values of the momentum transfers q . We also show the separate contributions of protons and neutrons. We observe that, as expected from the different values of the Fermi momentum, the proton ARFG responses are higher than the SRFG ones and limited to a narrower region of ω , whereas the opposite occurs for neutrons. These two effects tend to cancel in the total transverse nuclear response R^T , which is affected only mildly by the N/Z asymmetry. On the other hand, in the longitudinal channel R^L , where the proton response dominates, asymmetry effects can be non-negligible. Specifically, at $q = 800 \text{ MeV}/c$ the ratios of the ARFG/SRFG responses at the quasielastic peak for argon are of the order of 1.03 (1.01) for L (T) and are of the order of 1.07 (1.03) for L (T) for lead. Roughly, the ratios are similar as functions of the momentum transfer.

In Figs. 3 and 4 we show the weak neutral-current (WNC) longitudinal (L) and transverse (both T and T') response functions for ^{40}Ar and ^{208}Pb , obtained in the usual way [27,28] by replacing the EM couplings by WNC couplings and adding the T' VA -interference response; recall that the last enters with the opposite sign for neutrinos and antineutrinos in the total cross section. Also, note that in this study we have ignored the effects from strangeness content in the nucleons.

The effects are similar to what was found above for electron scattering, but not exactly the same for the cases which can be directly compared (viz., L and T), implying that for asymmetric nuclei there are effects to be taken into account in using input from electron scattering to obtain parts of the WNC cross section, as is often done in scaling analyses. The purpose of the present study is to get some idea about how significant such effects can be. Specifically, again at $q = 800 \text{ MeV}/c$ the ratios of the ARFG/SRFG responses at the quasielastic peak for argon are of the order of 0.98 (1.01) for L (T) and are of the order of 0.96 (1.02) for L (T) for lead, while the ratios for the T' response are of the order of 1.00 and 1.01 for argon and lead, respectively. Furthermore, in (ν, ν') the ARFG responses are lower and more extended than the SRFG ones, while the opposite occurs for $(\bar{\nu}, \bar{\nu}')$.

B. Charged-current reactions

Let us now consider CC neutrino and antineutrino reactions. In this case the inclusive cross section is the combination of five, instead of three, response functions, as a consequence of the nonconservation of the axial current and of the nonvanishing mass of the outgoing charged lepton.




FIG. 4. Weak NC response functions for ν - ^{208}Pb in the SRFG and ARFG. The separate contributions of protons and neutrons are also displayed.

The numerical results for (ν_μ, μ^-) and $(\bar{\nu}_\mu, \mu^+)$ scattering corresponding to carbon, argon, and lead targets are shown in Figs. 5–10. Although not shown here, oxygen and iron, used as well in neutrino oscillation experiments, have also been considered. The results are very similar to those obtained for carbon and argon, respectively.

The main observation here is that the differences between the SRFG and ARFG results are much larger than was seen above for the NC processes. As noted in Sec. II A, for NC processes in the ARFG model the energy offsets cancel and the differences between the SRFG and ARFG arise entirely from the different Fermi momenta entering for protons and neutrons. In contrast, for charge-changing weak interactions this is not the case because the relative offsets for protons and neutrons (which are usually different) do enter. Hence the results corresponding to symmetric target nuclei, such as ^{12}C and ^{16}O , will also differ from the usual SRFG results, since the final nucleus will have $Z \neq N$. In order to disentangle these two effects, we also show in each plot the results, labeled as “ARFG, no shift,” where the separation energies S_p and S_n are set to zero.

We observe the following:

- (i) For ^{12}C the effect of the energy offsets is large, as could be anticipated looking at Table II, while the difference in k_F plays a minor role; these effects are generally larger for neutrinos than for antineutrinos.
- (ii) For ^{40}Ar , which is relevant for neutrino oscillation studies, the two effects are comparable and both contribute to a slight shift of the responses to higher energy transfers, in particular for low q .
- (iii) For ^{208}Pb the main differences between SRFG and ARFG are due to Fermi momentum effects, which shift the neutrino (antineutrino) responses to higher (lower) energy transfers.

As was done in the NC case, to get an idea of the importance of the asymmetry effects we list in Table III the ratios between the ARFG and SRFG responses for neutrino and antineutrino scattering at $q = 800 \text{ MeV}/c$. We observe that the effects are minor for argon, while they are important for carbon and lead. Moreover, due to the different origin of the effects illustrated above, in the case of carbon the asymmetric model yields higher responses at the quasielastic peak for both neutrino




FIG. 5. Neutrino charged-current weak response functions per neutron of ^{12}C in the SRFG and ARFG. The ARFG results with no energy shift (see text) are also shown. Each column corresponds to a fixed value of the momentum transfer q .




FIG. 6. Same as Fig. 5, but for the antineutrino functions.




FIG. 7. Neutrino CC weak response functions per neutron of ^{40}Ar in the SRFG and ARFG. The ARFG results with no energy shift (see text) are also shown. Each column corresponds to a fixed value of the momentum transfer q .




FIG. 8. Same as Fig. 7, but for the antineutrino functions.




FIG. 9. Neutrino CC weak response functions per neutron of ^{208}Pb in the SRFG and ARFG. The ARFG results with no energy shift (see text) are also shown. Each column corresponds to a fixed value of the momentum transfer q .




FIG. 10. Same as Fig. 9, but for antineutrino functions.

TABLE III. Ratios between the ARFG and SRFG weak response functions at the QEP for $q = 800 \text{ MeV}/c$ for neutrino (left table) and antineutrino (right table) CC scattering.

(ν_μ, μ^-)	CC	CL	LL	T	T'	$(\bar{\nu}_\mu, \mu^+)$	CC	CL	LL	T	T'
^{12}C	1.24	1.31	1.37	1.02	1.03	^{12}C	1.15	1.19	1.22	1.02	1.01
^{40}Ar	1.06	1.08	1.11	0.98	0.98	^{40}Ar	1.05	1.05	1.05	1.04	1.04
^{208}Pb	1.15	1.21	1.27	0.96	0.97	^{208}Pb	0.98	0.95	0.92	1.09	1.08

and antineutrino scattering and in all five channels, while for lead the asymmetry effects have the opposite sign for neutrinos and antineutrinos and they depend on the channel: they increase (decrease) the neutrino (antineutrino) charge and/or longitudinal responses, whereas the opposite occurs for the transverse T and T' responses.

IV. CONCLUSIONS

In the present study we have developed an extension of the familiar relativistic Fermi gas model for inclusive semileptonic electroweak processes. Specifically, we have developed the model for neutral-current scattering of electrons or (anti)neutrinos and for charge-changing (anti)neutrino reactions with nuclei. The new element in this work has been to allow for effects arising from differences between protons and neutrons within the context of the relativistic Fermi gas in order to evaluate how significant such effects may be in progressing from light $N = Z$ (i.e., symmetric) nuclei to asymmetric nuclei having $N > Z$. We denote the usual relativistic Fermi gas model, typically abbreviated RFG, to be the symmetric relativistic Fermi gas (SRFG), while the new extension to asymmetric nuclei we denote as the asymmetric relativistic Fermi gas (ARFG).

Two types of extensions have been studied: first, we consider only nuclei in the valley of stability where typically the volume occupied by protons and neutrons in the ground states of such nuclei is the same, and hence where the densities scale by the numbers of neutrons and protons. In the context of the Fermi gas this implies that the Fermi momenta for n and p will be different, scaling by $(N/Z)^{1/3}$. Second, we adjust the Fermi energies of the proton and neutron gases of the parent nucleus, and its neighbors in the case of CC (anti)neutrino reactions, to agree with the measured values. We note that this is a basic assumption in the present ARFG model and not the only way one might proceed. For instance, one might develop a different model where the energy offsets involved are allowed to be chosen by forcing agreement with experiment. The motivation in the present work is to explore the typical size of these second effects to see if they are typically negligible or if they should be taken seriously in future more sophisticated modeling.

One conclusion is that the density effect (leading to different Fermi momenta for neutrons and protons) plays no role at all for NC scattering from symmetric nuclei and a relatively minor role for CC (anti)neutrino reactions in such systems where there is some effect since neighboring nuclei which have slightly different Fermi momenta are involved. In contrast, for very asymmetric nuclei such as ^{208}Pb the effects from having differing neutron and proton Fermi momenta are somewhat

larger, although still relatively minor, for NC scattering, but much more significant for CC (anti)neutrino reactions.

A second observation is that in NC scattering (electrons or neutrinos) the energy offsets do not play a role; simply put, only energy differences between particles and holes enter and, since the 1p1h states involve only excitations of protons or neutrons individually, the offsets cancel. In contrast, for CC (anti)neutrino reactions neutrons change into protons or vice versa and thus the offsets do play a role. For the last type of reaction one sees that the energy offset effect is dominant in light symmetric nuclei such as ^{12}C , roughly comparable to the density effect for ^{40}Ar , and subdominant to the density effect in very asymmetric nuclei such as ^{208}Pb .

The ARFG model developed in this study can be extended straightforwardly to include inelastic processes following previous work done along these lines for the SRFG. Finally, while much more involved than the 1p1h focus of the present work, it is possible to extend the previous 2p2h SRFG studies of two-body Meson-Exchange Currents contributions [29,30] to incorporate asymmetric nuclei; such a study is in progress.

We believe that this study will give valuable indications on how to extend more sophisticated nuclear models to asymmetric nuclei. Moreover, it will provide relatively simple recipes for the implementation of asymmetry effects in Monte Carlo generators used to analyze neutrino oscillation experiments.

ACKNOWLEDGMENTS

This work was partially supported by the INFN under project MANYBODY, by the University of Turin under contract BARM-RILO-17, by the Spanish Ministerio de Economía y Competitividad and ERDF (European Regional Development Fund) under contracts FIS2014-53448-C2-1 and FIS2017-88410-P, by the Junta de Andalucía Grant No. FQM160, and part (T.W.D.) by the US Department of Energy under cooperative agreement DE-FC02-94ER40818. M.B.B. acknowledges support from the “Emilie du Châtelet” program of the P2IO LabEx (ANR-10-LABX-0038). G.D.M. acknowledges support from a Junta de Andalucía fellowship (FQM7632, Proyectos de Excelencia 2011). J.W.V.O. acknowledges support by the US Department of Energy under Contract No. DE-AC05-06OR23177, and by the US Department of Energy cooperative research agreement DE-AC05-84ER40150.

This article is authored by Jefferson Science Associates, LLC under US DOE Contract No. DE-AC05-06OR23177. The US Government retains a nonexclusive, paid-up, irrevocable, worldwide license to publish or reproduce this manuscript for US Government purposes.

- [1] L. Alvarez-Ruso *et al.*, *Prog. Part. Nucl. Phys.* **100**, 1 (2018).
- [2] R. Acciarri *et al.* (DUNE Collaboration), [arXiv:1512.06148](#).
- [3] G. D. Megias, T. W. Donnelly, O. Moreno, C. F. Williamson, J. A. Caballero, R. González-Jiménez, A. De Pace, M. B. Barbaro, W. M. Alberico, M. Nardi *et al.*, *Phys. Rev. D* **91**, 073004 (2015).
- [4] G. D. Megias, J. E. Amaro, M. B. Barbaro, J. A. Caballero, T. W. Donnelly, and I. R. Simo, *Phys. Rev. D* **94**, 093004 (2016).
- [5] N. Rocco, A. Lovato, and O. Benhar, *Phys. Rev. Lett.* **116**, 192501 (2016).
- [6] A. M. Ankowski and J. T. Sobczyk, *Phys. Rev. C* **77**, 044311 (2008).
- [7] A. M. Ankowski, O. Benhar, C. Mariani, and E. Vagnoni, *Phys. Rev. D* **93**, 113004 (2016).
- [8] A. Meucci, F. Capuzzi, C. Giusti, and F. D. Pacati, *Phys. Rev. C* **67**, 054601 (2003).
- [9] A. Meucci and C. Giusti, *Phys. Rev. D* **91**, 093004 (2015).
- [10] J. Nieves, I. Ruiz Simo, and M. J. Vicente Vaca, *Phys. Lett. B* **707**, 72 (2012).
- [11] M. Martini, M. Ericson, G. Chanfray, and J. Marteau, *Phys. Rev. C* **81**, 045502 (2010).
- [12] A. Lovato, S. Gandolfi, J. Carlson, S. C. Pieper, and R. Schiavilla, *Phys. Rev. C* **91**, 062501 (2015).
- [13] V. Pandey, N. Jachowicz, M. Martini, R. González-Jiménez, J. Ryckebusch, T. Van Cuyck, and N. Van Dessel, *Phys. Rev. C* **94**, 054609 (2016).
- [14] T. W. Donnelly and I. Sick, *Phys. Rev. Lett.* **82**, 3212 (1999).
- [15] T. W. Donnelly and I. Sick, *Phys. Rev. C* **60**, 065502 (1999).
- [16] C. Maieron, T. W. Donnelly, and I. Sick, *Phys. Rev. C* **65**, 025502 (2002).
- [17] R. González-Jiménez, G. D. Megias, M. B. Barbaro, J. A. Caballero, and T. W. Donnelly, *Phys. Rev. C* **90**, 035501 (2014).
- [18] G. D. Megias, J. E. Amaro, M. B. Barbaro, J. A. Caballero, and T. W. Donnelly, *Phys. Rev. D* **94**, 013012 (2016).
- [19] W. M. Alberico, A. Drago, and C. Villavechia, *Nucl. Phys. A* **505**, 309 (1989).
- [20] W. M. Alberico, G. Chanfray, M. Ericson, and A. Molinari, *Nucl. Phys. A* **475**, 233 (1987).
- [21] D. Davesne, A. Pastore, and J. Navarro, *Phys. Rev. C* **89**, 044302 (2014).
- [22] E. Lipparini and F. Pederiva, *Phys. Rev. C* **88**, 024318 (2013).
- [23] E. Lipparini and F. Pederiva, *Phys. Rev. C* **94**, 024323 (2016).
- [24] M. B. Barbaro, J. A. Caballero, T. W. Donnelly, and C. Maieron, *Phys. Rev. C* **69**, 035502 (2004).
- [25] R. Cenni, T. W. Donnelly, and A. Molinari, *Phys. Rev. C* **56**, 276 (1997).
- [26] J. Tepel, *Comput. Phys. Commun.* **33**, 129 (1984).
- [27] J. E. Amaro, M. B. Barbaro, J. A. Caballero, T. W. Donnelly, A. Molinari, and I. Sick, *Phys. Rev. C* **71**, 015501 (2005).
- [28] J. E. Amaro, M. B. Barbaro, J. A. Caballero, and T. W. Donnelly, *Phys. Rev. C* **73**, 035503 (2006).
- [29] I. Ruiz Simo, J. E. Amaro, M. B. Barbaro, A. De Pace, J. A. Caballero, and T. W. Donnelly, *J. Phys. G* **44**, 065105 (2017).
- [30] J. E. Amaro, M. B. Barbaro, J. A. Caballero, A. De Pace, T. W. Donnelly, G. D. Megias, and I. Ruiz Simo, *Phys. Rev. C* **95**, 065502 (2017).