

2019

Conformal Invariance of Transverse-Momentum Dependent Parton Distributions Rapidity Evolution

Ian Balitsky
Old Dominion University, ibalitsk@odu.edu

Giovanni A. Chirilli

Follow this and additional works at: https://digitalcommons.odu.edu/physics_fac_pubs



Part of the [Elementary Particles and Fields and String Theory Commons](#), and the [Nuclear Commons](#)

Original Publication Citation

Balitsky, I., & Chirilli, G. A. (2019). Conformal invariance of transverse-momentum dependent parton distributions rapidity evolution. *Physical Review D: Particles, Fields, Gravitation & Cosmology*, 100(5), 051504. doi:10.1103/PhysRevD.100.051504

This Article is brought to you for free and open access by the Physics at ODU Digital Commons. It has been accepted for inclusion in Physics Faculty Publications by an authorized administrator of ODU Digital Commons. For more information, please contact digitalcommons@odu.edu.

Conformal invariance of transverse-momentum dependent parton distributions rapidity evolution

Ian Balitsky

*Physics Department, Old Dominion University, Norfolk, Virginia 23529, USA
and Theory Group, JLAB, 12000 Jefferson Avenue, Newport News, Virginia 23606, USA*

Giovanni A. Chirilli

Institut für Theoretische Physik, Universität Regensburg, D-93040 Regensburg, Germany



(Received 17 June 2019; published 20 September 2019)

We discuss conformal properties of TMD operators and present the result of the conformal rapidity evolution of TMD operators in the Sudakov region.

DOI: [10.1103/PhysRevD.100.051504](https://doi.org/10.1103/PhysRevD.100.051504)

I. INTRODUCTION

In recent years, the transverse-momentum dependent parton distributions (TMDs) [1–4] have been widely used in the analysis of processes like semi-inclusive deep inelastic scattering or particle production in hadron-hadron collisions (for a review, see Ref. [5]).

The TMDs are defined as matrix elements of quark or gluon operators with attached lightlike gauge links (Wilson lines) going to either $+\infty$ or $-\infty$ depending on the process under consideration. It is well known that these TMD operators exhibit rapidity divergencies due to infinite lightlike gauge links and the corresponding rapidity/UV divergences should be regularized. There are two schemes on the market: the most popular is based on Collins-Soper-Sterman [2] or soft-collinear effective theory [6] formalism, and the second one is adopted from the small- x physics [7,8]. The obtained evolution equations differ even at the leading-order level and need to be reconciled, especially in view of the future electron-ion collider accelerator which will probe the TMDs at values of Bjorken x between small- x and $x \sim 1$ regions.

In our opinion, a good starting point is to obtain conformal leading-order evolution equations. It is well known that at the leading-order perturbative QCD (pQCD) is conformally invariant, so there is hope of getting any evolution equation without explicit running coupling from conformal considerations. In our case, since TMD operators are defined with attached lightlike Wilson lines, formally they will transform covariantly under the subgroup of the full conformal group which preserves this lightlike direction. However, as we mentioned, the TMD

operators contain rapidity divergencies which need to be regularized. At present, there is no rapidity cutoff which preserves conformal invariance, so the best one can do is to find the cutoff which is conformal at the leading order in perturbation theory. In higher orders, one should not expect conformal invariance since it is broken by the running of QCD coupling. However, if one considers corresponding correlation functions in $\mathcal{N} = 4$ super Yang-Mills (SYM), one should expect conformal invariance. After that, the results obtained in $\mathcal{N} = 4$ SYM theory can be used as a starting point of QCD calculation. Typically, the result in $\mathcal{N} = 4$ theory gives the most complicated part of the pQCD result, i.e., the one with maximal transcendentality. Thus, the idea is to find the TMD operator conformal in $\mathcal{N} = 4$ SYM and use it in QCD. This scheme was successfully applied to the rapidity evolution of color dipoles. At the leading order, the Balitsky-Kovchegov evolution of color dipoles [9–12] is invariant under $SL(2, \mathbb{C})$ (Möbius) group. At the next-to-leading order (NLO), the “conformal dipole” with the α_s correction [13] makes NLO Balitsky-Kovchegov evolution Möbius invariant for $\mathcal{N} = 4$ SYM, and the corresponding QCD kernel [14] differs by terms proportional to the β function.

II. CONFORMAL INVARIANCE OF TMD OPERATORS

For definiteness, we will talk first about gluon operators with lightlike Wilson lines stretching to $-\infty$ in the $+$ direction. The gluon TMD (unintegrated gluon distribution) is defined as [15]

$$\begin{aligned} \mathcal{D}(x_B, k_\perp, \eta) &= \int d^2 z_\perp e^{i(k, z)_\perp} \mathcal{D}(x_B, z_\perp, \eta), \\ g^2 \mathcal{D}(x_B, z_\perp, \eta) &= \frac{-x_B^{-1}}{2\pi p^-} \int dz^+ e^{-ix_B p^- z^+} \\ &\quad \times \langle P | \mathcal{F}_\xi^a(z) [z - \infty n, -\infty n]^{ab} \mathcal{F}_\xi^b(0) | P \rangle \Big|_{z^-=0}, \end{aligned} \quad (1)$$

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

where $|P\rangle$ is an unpolarized target with momentum $p \simeq p^-$ (typically proton) and $n = (\frac{1}{\sqrt{2}}, 0, 0, \frac{1}{\sqrt{2}})$ is a lightlike vector in the “+” direction. Hereafter, we use the notation

$$\mathcal{F}^{\xi,a}(z_\perp, z^+) \equiv gF^{-\xi,m}(z)[z, z - \infty n]^{ma}|_{z^-=0}, \quad (2)$$

where $[x, y]$ denotes straight-line gauge link connecting points x and y :

$$[x, y] \equiv \text{Pe}^{ig \int du(x-y)^\mu A_\mu(ux+(1-u)y)}. \quad (3)$$

To simplify one-loop evolution, we multiplied $F_{\mu\nu}$ by a coupling constant. Since the gA_μ is renormalization invariant, we do not need to consider self-energy diagrams (in the background-Feynman gauge). Note that $z^- = 0$ is fixed by the original factorization formula for particle production [5] (see also the discussion in Refs. [16,17]).

The algebra of full conformal group $SO(2,4)$ consists of four operators P^μ , six $M^{\mu\nu}$, four special conformal generators K^μ , and dilatation operator D . It is easy to check that in the leading order the following 11 operators act on gluon TMDs covariantly,

$$P^i, P^-, M^{12}, M^{-i}, D, K^i, K^-, M^{-+}, \quad (4)$$

while the action of operators P^+ , M^{+i} , and K^+ do not preserve the form of the operator (2). The action of the generators (4) on the operator (2) is the same as the action on the field F^{-i} without gauge link attachments. The corresponding group consists of transformations which leave the hyperplane $z^- = 0$ and vector n invariant. Those include shifts in transverse and + directions, rotations in the transverse plane, Lorentz rotations/boosts created by M^{-i} , dilatations, and special conformal transformations

$$z'_\mu = \frac{z_\mu - a_\mu z^2}{1 - 2a \cdot z + a^2 z^2} \quad (5)$$

with $a = (a^+, 0, a_\perp)$. In terms of “embedding formalism” [18–21] defined in six-dimensional space, this subgroup is isomorphic to the “Poincaré + dilatations” group of the four-dimensional subspace orthogonal to our physical lightlike + and “−” directions.

As we noted, infinite Wilson lines in the definition (2) of TMD operators make them divergent. As we discussed above, it is very advantageous to have a cutoff of these divergencies compatible with approximate conformal invariance of tree-level QCD. The evolution equation with such a cutoff should be invariant with respect to transformations described above.

In the next section, we demonstrate that the “small- x ” rapidity cutoff enables us to get a conformally invariant evolution of TMD in the so-called Sudakov region.

III. TMD FACTORIZATION IN THE SUDAKOV REGION

The rapidity evolution of the TMD operator (1) is very different in the region of large and small longitudinal separations z^+ . The evolution at small z^+ is linear and double-logarithmic, while at large z^+ , the evolution becomes nonlinear due to the production of color dipoles typical for small- x evolution. It is convenient to consider as a starting point the simple case of TMD evolution in the so-called Sudakov region corresponding to small longitudinal distances.

First, let us specify what we call a Sudakov region. A typical factorization formula for the differential cross section of particle production in hadron-hadron collision is [5,22]

$$\frac{d\sigma}{d\eta d^2q_\perp} = \sum_f \int d^2b_\perp e^{i(q,b)_\perp} \mathcal{D}_{f/A}(x_A, b_\perp, \eta) \times \mathcal{D}_{f/B}(x_B, b_\perp, \eta) \sigma(ff \rightarrow H) + \dots, \quad (6)$$

where $\eta = \frac{1}{2} \ln \frac{q^+}{q^-}$ is the rapidity, $\mathcal{D}_{f/h}(x, z_\perp, \eta)$ is the TMD density of a parton f in hadron h , and $\sigma(ff \rightarrow H)$ is the cross section of production of particle H of invariant mass $m_H^2 = q^2 \equiv Q^2$ in the scattering of two partons. (One can keep in mind Higgs production in the approximation of the pointlike gluon-gluon-Higgs vertex). The Sudakov region is defined by $Q \gg q_\perp \gg 1$ GeV since at such kinematics there is a double-log evolution for transverse momenta between Q and q_\perp . In the coordinate space, TMD factorization (6) looks like

$$\begin{aligned} & \langle p_A, p_B | g^2 F_{\mu\nu}^a F^{a\mu\nu}(z_1) g^2 F_{\lambda\rho}^b F^{b\lambda\rho}(z_2) | p_A, p_B \rangle \\ &= \frac{1}{N_c^2 - 1} \langle p_A | \tilde{\mathcal{O}}_{ij}(z_1^-, z_{1\perp}; z_2^-, z_{2\perp}) | p_A \rangle^{\sigma_A} \\ & \times \langle p_B | \mathcal{O}^{ij}(z_1^+, z_{1\perp}; z_2^+, z_{2\perp}) | p_B \rangle^{\sigma_B} + \dots, \end{aligned} \quad (7)$$

where

$$\begin{aligned} \mathcal{O}_{ij}(z_1^+, z_{1\perp}; z_2^+, z_{2\perp}) &= \mathcal{F}_i^a(z_1)[z_1 - \infty n, z_2 - \infty n]^{ab} \\ & \times \mathcal{F}_j^b(z_2)|_{z_1^-=z_2^-=0}, \end{aligned} \quad (8)$$

$$\begin{aligned} \tilde{\mathcal{O}}_{ij}(z_1^-, z_{1\perp}; z_2^-, z_{2\perp}) &= \mathcal{F}_i^a(z_1)[z_1 - \infty n', z_2 - \infty n']^{ab} \\ & \times \mathcal{F}_j^b(z_2)|_{z_1^+=z_2^+=0}, \end{aligned}$$

$$\mathcal{F}^{i,a}(z_\perp, z^-) \equiv gF^{+i,m}(z)[z, z - \infty n']^{ma}|_{z^+=0}. \quad (9)$$

Here, $p_A = \sqrt{s}n + \frac{p_A^2}{\sqrt{2}s}n'$, $p_B = \sqrt{s}n' + \frac{p_B^2}{\sqrt{2}s}n$, and $n' = (\frac{1}{\sqrt{2}}, 0, 0, -\frac{1}{\sqrt{2}})$. Our metric is $x^2 = 2x^+x^- - x_\perp^2$.

As we mentioned, TMD operators exhibit rapidity divergencies due to infinite lightlike gauge links. The “small- x style” rapidity cutoff for longitudinal divergencies is imposed as the upper limit of k^+ components of gluons emitted from the Wilson lines. As we will see below, to get the conformal invariance of the leading-order evolution, we

need to impose the cutoff of k^+ components of gluons correlated with transverse size of TMD in the following way:

$$\mathcal{F}^{i,a}(z_\perp, z^+)^\sigma \equiv gF^{-i,m}(z)[Pe^{ig\int_{-\infty}^{z^+} dz^+ A^{-\sigma}(up_1+x_\perp)}]^{ma},$$

$$A_\mu^\sigma(x) = \int \frac{d^4k}{16\pi^4} \theta\left(\frac{\sigma\sqrt{2}}{z_{12\perp}} - |k^+|\right) e^{-ik \cdot x} A_\mu(k). \quad (10)$$

Similarly, the operator $\tilde{\mathcal{O}}$ in Eq. (9) is defined with the rapidity cutoff for β integration imposed as $\theta(\frac{\tilde{\sigma}\sqrt{2}}{z_{12\perp}} - |k^-|)$.

The Sudakov region $Q^2 \gg q_\perp^2$ in the coordinate space corresponds to

$$z_{12\parallel}^2 \equiv 2z_{1\perp}^- z_{2\perp}^+ \ll z_{12\perp}^2, \quad (11)$$

where $z_{12} \equiv z_1 - z_2$. In the leading log approximation, the upper cutoff for k^+ integration in the target matrix element in Eq. (7) is $\sigma_B = \frac{1}{\sqrt{2}} \frac{z_{12\perp}^2}{z_{12}^+}$, and similarly the β -integration cutoff in the projectile matrix element is $\sigma_A = \frac{1}{\sqrt{2}} \frac{z_{12\perp}^2}{z_{12}^-}$ [23].

In the next section, we demonstrate that the rapidity cutoff (10) enables us to get a conformally invariant evolution of TMD in the Sudakov region (11).

IV. ONE-LOOP EVOLUTION OF TMDs

A. Evolution of gluon TMD operators in the Sudakov region

In this section, we derive the evolution of gluon TMD operator (8) with respect to cutoff σ in the leading log approximation. As usual, to get an evolution equation, we integrate over momenta $\frac{\sigma_2\sqrt{2}}{z_{12\perp}} > k^+ > \frac{\sigma_1\sqrt{2}}{z_{12\perp}}$. To this end, we calculate diagrams shown in Fig. 1 in the background field of gluons with $k^+ < \frac{\sigma_1\sqrt{2}}{z_{12\perp}}$. The calculation is easily done by method developed in Refs. [24,25], and the result is

$$\mathcal{O}^{\sigma_2}(z_1^+, z_2^+) = \frac{\alpha_s N_c}{2\pi} \int_{\frac{\sigma_1\sqrt{2}}{z_{12\perp}}}^{\frac{\sigma_2\sqrt{2}}{z_{12\perp}}} \frac{dk^+}{k^+} K \mathcal{O}^{\sigma_1}(z_1^+, z_2^+), \quad (12)$$

where the kernel K is given by

$$K\mathcal{O}(z_1^+, z_2^+) = \mathcal{O}(z_1^+, z_2^+) \int_{-\infty}^{z_1^+} \frac{dz^+}{z_2^+ - z^+} e^{-i\frac{z_{12\perp}\sigma}{\sqrt{2}(z_2-z)^+}} \\ + \mathcal{O}(z_1^+, z_2^+) \int_{-\infty}^{z_2^+} \frac{dz^+}{z_1^+ - z^+} e^{-i\frac{z_{12\perp}\sigma}{\sqrt{2}(z_1-z)^+}} \\ - \int_{-\infty}^{z_1^+} dz^+ \frac{\mathcal{O}(z_1^+, z_2^+) - \mathcal{O}(z^+, z_2^+)}{z_1^+ - z^+} \\ - \int_{-\infty}^{z_2^+} dz^+ \frac{\mathcal{O}(z_1^+, z_2^+) - \mathcal{O}(z_1^+, z^+)}{z_2^+ - z^+}, \quad (13)$$




FIG. 1. Typical diagrams for production (a) and virtual (b) contributions to the evolution kernel. The dashed lines denote gauge links.

where we suppress arguments $z_{1\perp}$ and $z_{2\perp}$ since they do not change during the evolution in the Sudakov regime. The first two terms in the kernel K come from the “production” diagram in Fig. 1(a), while the last two terms come from the “virtual” diagram in Fig. 1(b). The result (13) can be also obtained from Ref. [25] by Fourier transformation of Eq. (5.9) with the help of Eqs. (3.12) and (3.30) therein. The approximations for diagrams in Fig. 1 leading to Eq. (13) are valid as long as

$$k^+ \gg \frac{z_{12\perp}^2}{z_{12\perp}^2}, \quad (14)$$

which gives the region of applicability of Sudakov-type evolution.

Evolution equation (12) can be easily integrated using Fourier transformation. Since

$$K e^{-ik^- z_1^+ + ik'^- z_2^+} = \left[-2 \ln \sigma z_{12\perp} - \ln(ik^-) - \ln(-ik'^-) \right. \\ \left. + \ln 2 - 4\gamma_E + O\left(\frac{z_{12\perp}^2}{z_{12\perp}^2 \sigma}\right) \right] e^{-ik^- z_1^+ + ik'^- z_2^+}, \quad (15)$$

one easily obtains

$$\mathcal{O}^{\sigma_2}(z_1^+, z_2^+) = e^{-2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1} [\ln \sigma_1 \sigma_2 + 4\gamma_E - \ln 2]} \\ \times \int dz_1'^+ dz_2'^+ \mathcal{O}^{\sigma_1}(z_1'^+, z_2'^+) z_{12\perp}^{-2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1}} \\ \times \frac{1}{4\pi^2} \left[\frac{i\Gamma(1 - 2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1})}{(z_1^+ - z_1'^+ + i\epsilon)^{1-2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1}} + \text{c.c.}} \right] \\ \times \left[\frac{i\Gamma(1 - 2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1})}{(z_2^+ - z_2'^+ + i\epsilon)^{1-2\bar{\alpha}_s \ln \frac{\sigma_2}{\sigma_1}} + \text{c.c.}} \right], \quad (16)$$

where we introduced notation $\bar{\alpha}_s \equiv \frac{\alpha_s N_c}{4\pi}$. It should be mentioned that the factor $4\gamma_E$ is “scheme dependent”;

if one introduces to α integrals smooth cutoff $e^{-\alpha/a}$ instead of rigid cutoff $\theta(a > \alpha)$, the value $4\gamma_E$ changes to $2\gamma_E$.

It is easy to see that the rhs of Eq. (16) transforms covariantly under all transformations (4) except the Lorentz boost generated by M^{+-} . The reason is that the Lorentz boost in the z direction changes cutoffs for the evolution. To understand that, note that Eq. (15) is valid until $\sigma > \frac{z_{12}^+}{z_{12\perp}^2}$, so the linear evolution (16) is applicable in the region between

$$\sigma_2 = \sigma_B = \frac{z_{12\perp}}{z_{12}^+ \sqrt{2}} \quad \text{and} \quad \sigma_1 = \frac{z_{12}^+ \sqrt{2}}{z_{12\perp}}. \quad (17)$$

From Eq. (16), it is easy to see that Lorentz boost $z^+ \rightarrow \lambda z^+$, $z^- \rightarrow \frac{1}{\lambda} z^-$ changes the value of target matrix element $\langle p_A | \mathcal{O} | p_B \rangle$ by $\exp\{4\lambda \bar{\alpha}_s \ln \frac{z_{12\perp}^2}{z_{12\perp}^2}\}$, but simultaneously it will change the result of similar evolution for projectile matrix element $\langle p_A | \tilde{\mathcal{O}} | p_A \rangle$ by $\exp\{-4\lambda \bar{\alpha}_s \ln \frac{z_{12\perp}^2}{z_{12\perp}^2}\}$, so the overall result for the amplitude (7) remains intact.

B. Evolution of quark TMD operators

A simple calculation of evolution of quark operator

$$\mathcal{O}_q(z_1^+, z_{1\perp}; z_2^+, z_{2\perp}) \equiv g^{\frac{2C_F}{b}} \bar{\psi}(z_\perp + un)[un + z_\perp, -\infty n] \not{n}[z_\perp - \infty n, -\infty n][\infty n, 0]\psi(0) \quad (18)$$

yields the same evolution (16) as for the gluon operators with trivial replacement $N_c \rightarrow C_F$ [26]. The factor $g^{\frac{2C_F}{b}}$ ($b \equiv \frac{11}{3}N_c - \frac{2}{3}n_f$) is added to avoid taking into account quark self-energy.

C. Evolution beyond Sudakov region

As we mentioned above, the TMD factorization formula (6) for particle production at $q_\perp \ll Q$ translates to the coordinate space as Eq. (7) with the requirement $z_{12\perp}^2 \ll z_{12\perp}^2$. During the evolution (16), the transverse separation between gluon operators \mathcal{F}_i and \mathcal{F}_j remains intact, while the longitudinal separation increases. As discussed in Refs. [24,25], the Sudakov approximation can be trusted until the upper cutoff in α integrals is greater than $\frac{q_\perp^2}{x_B s}$, which is equivalent to Eq. (14) in the coordinate space. If $x_B \sim 1$ and $q_\perp \sim m_N$, the relative energy between Wilson-line operators \mathcal{F} and the target nucleon at the final point of evolution is approximately m_N^2 , so one should use phenomenological models of TMDs with this low rapidity cutoff as a starting point of the evolution (16). If, however, $x_B \ll 1$, this relative energy is $\frac{q_\perp^2}{x_B s} \gg m_N^2$, so one can continue the rapidity evolution in the region $\frac{q_\perp^2}{x_B s} > \sigma > \frac{m_N^2}{s}$ beyond the Sudakov region into the small- x region. The evolution in a “proper” small- x region is known [27]—the TMD operator,

known also as Weizsäcker-Williams distribution, will produce a hierarchy of color dipoles as a result of the nonlinear evolution. However, the transition between the Sudakov region and small- x region is described by a rather complicated interpolation formula [24]. In the coordinate space, this means the study of operator \mathcal{O} at $z_\parallel^2 \sim z_\perp^2$, and we hope that conformal considerations can help us to obtain the TMD evolution in that region.

V. DISCUSSION

As we mentioned in the Introduction, TMD evolution is analyzed by very different methods at small x and moderate $x \sim 1$. In view of the future electron-ion collider accelerator, which will probe the region between small x and $x \sim 1$, we need a universal description of TMD evolution valid at both limits. Since the two formalisms differ even at the leading order where QCD is conformally invariant, our idea is to make this universal description first in $\mathcal{N} = 4$ SYM. In a first step, we found a conformally invariant evolution in the Sudakov region using our small- x cutoff with the “conformal refinement” (10).

To compare with conventional TMD analysis, let us write down the evolution of “generalized TMD” [28,29]

$$D^\sigma(x, \xi) = \int dz^+ e^{-ix\sqrt{\xi}z^+} \left\langle p'_B \left| \mathcal{O}^\sigma \left(-\frac{z^+}{2}, \frac{z^+}{2} \right) \right| p_B \right\rangle,$$

where $\xi = -\frac{p'_B - p_B}{\sqrt{2}s}$. From Eq. (16), one easily obtains

$$\frac{D^{\sigma_2}(x, \xi)}{D^{\sigma_1}(x, \xi)} = e^{-2\bar{\alpha}_s \ln \frac{Q^2}{\sigma_1} [\ln \sigma_2 \sigma_1 (x^2 - \xi^2) s z_{12\perp}^2 + 4\gamma_E - \ln 2]}. \quad (19)$$

For usual TMD at $\xi = 0$ with the limits of Sudakov evolution set by Eq. (17), one obtains

$$\frac{D^{\sigma_2}(x, q_\perp)}{D^{\sigma_1}(x, q_\perp)} = e^{-2\bar{\alpha}_s \ln \frac{Q^2}{q_\perp^2} [\ln \frac{Q^2}{q_\perp^2} + 4\gamma_E - \ln 2]}, \quad (20)$$

which coincides with usual one-loop evolution of TMDs [30] up to replacement $4\gamma_E - 2 \ln 2 \rightarrow 4\gamma_E - 4 \ln 2$. As we discussed, such a constant depends on the way of cutting k^- integration, which should be coordinated with the cutoffs in the “coefficient function” $\sigma(ff \rightarrow H)$ in Eq. (6). Thus, the discrepancy is just like using two different schemes for usual renormalization. It should be mentioned, however, that at $\xi \neq 0$ the result (19) differs from the conventional one-loop result, which does not depend on ξ ; see, e.g., Ref. [31].

VI. CONCLUSIONS

The first result of our paper is finding the subgroup of $SO(2, 4)$, which formally leaves TMD operators invariant. Although there was some discussion of conformal invariance of the TMD approach in the literature [32,33], to the best of our knowledge, we present the first complete description of that subgroup.

The second result is related to the fact that conformal invariance is violated by the rapidity cutoff (even in $\mathcal{N} = 4$ SYM). As we mentioned above, since tree-level QCD is conformally invariant, it is convenient to have a leading-order evolution which respects that symmetry so the NLO corrections can be sorted out as conformal plus proportional to the β function. We have studied the TMD evolution in the Sudakov region of intermediate x and demonstrated that the rapidity cutoff used in small- x literature preserves all generators of our subgroup except the Lorentz boost, which is related to the change of that cutoff. It should be mentioned that usually the analysis of TMD evolution in the $x \sim 1$ region is performed with a combination of UV and rapidity cutoffs, which gives two evolution equations, in μ^2 and ζ (related to rapidity). However, although the results of these two evolutions are known at two- [34–36] and three-loop [37] levels, their relation to conformal properties of TMD operators is not obvious. It would be interesting to check if our cutoff corresponds to some conformal evolution path in the two-dimensional (μ^2, ζ) plane [38].

Our main outlook is to try to connect to the small- x region, first in $\mathcal{N} = 4$ SYM and then in QCD. As we mentioned above, although the TMD evolution in a small- x region is conformal with respect to the $SL(2, C)$ group, and our evolution (16) is also conformal [albeit with respect to a different group of which $SL(2, C)$ is a subgroup], the transition between the Sudakov region and small- x region is described by a rather complicated interpolation formula [24] which is not conformally invariant. Our hope is that in a conformal theory one can simplify that transition using the conformal invariance requirement. The study is in progress.

ACKNOWLEDGMENTS

We thank V.M. Braun, A. Vladimirov, and A. Tarasov for discussions. The work of I. Balitsky is supported by Jefferson Science Associates, LLC under U.S. Contract No. DE-AC05-06OR23177 and by U.S. Grant No. DE-FG02-97ER41028.

-
- [1] J. C. Collins and D. E. Soper, *Nucl. Phys.* **B194**, 445 (1982).
 - [2] J. C. Collins, D. E. Soper, and G. F. Sterman, *Nucl. Phys.* **B250**, 199 (1985).
 - [3] X.-d. Ji, J.-p. Ma, and F. Yuan, *Phys. Rev. D* **71**, 034005 (2005).
 - [4] M. G. Echevarria, A. Idilbi, and I. Scimemi, *J. High Energy Phys.* **07** (2012) 002.
 - [5] J. Collins, *Foundations of Perturbative QCD* (Cambridge University Press, Cambridge, England, 2013).
 - [6] I. Z. Rothstein and I. W. Stewart, *J. High Energy Phys.* **08** (2016) 025.
 - [7] L. N. Lipatov, *Phys. Rep.* **286**, 131 (1997).
 - [8] Y. V. Kovchegov and E. Levin, *Quantum Chromodynamics at High Energy*, Cambridge Monographs on Particle Physics, Nuclear Physics and Cosmology (Cambridge University Press, Cambridge, England, 2012).
 - [9] I. Balitsky, *Nucl. Phys.* **B463**, 99 (1996).
 - [10] I. Balitsky, *Phys. Rev. D* **60**, 014020 (1999).
 - [11] Y. V. Kovchegov, *Phys. Rev. D* **60**, 034008 (1999).
 - [12] Y. V. Kovchegov, *Phys. Rev. D* **61**, 074018 (2000).
 - [13] I. Balitsky and G. A. Chirilli, *Nucl. Phys.* **B822**, 45 (2009).
 - [14] I. Balitsky and G. A. Chirilli, *Phys. Rev. D* **77**, 014019 (2008).
 - [15] P. J. Mulders and J. Rodrigues, *Phys. Rev. D* **63**, 094021 (2001).
 - [16] I. Balitsky and A. Tarasov, *J. High Energy Phys.* **07** (2017) 095.
 - [17] I. Balitsky and A. Tarasov, *J. High Energy Phys.* **05** (2018) 150.
 - [18] P. A. M. Dirac, *Ann. Math.* **37**, 429 (1936).
 - [19] G. Mack and A. Salam, *Ann. Phys. (N.Y.)* **53**, 174 (1969).
 - [20] S. Ferrara, A. F. Grillo, and R. Gatto, *Ann. Phys. (N.Y.)* **76**, 161 (1973).
 - [21] S. Rychkov, *EPFL Lectures on Conformal Field Theory in $D \geq 3$ Dimensions*, Springer Briefs in Physics (Springer, Cham, 2016).
 - [22] J. Collins and T. Rogers, *Phys. Rev. D* **91**, 074020 (2015).
 - [23] Hereafter, we use the simplified notation $z_{12\perp} \equiv |z_{12\perp}|$.
 - [24] I. Balitsky and A. Tarasov, *J. High Energy Phys.* **10** (2015) 017.
 - [25] I. Balitsky and A. Tarasov, *J. High Energy Phys.* **06** (2016) 164.
 - [26] We assume that \mathcal{O}^q is defined with the same type of cutoff (10) imposed on quarks and gluons emitted as a result of evolution of \mathcal{O}^q .
 - [27] F. Dominguez, A. H. Mueller, S. Munier, and B.-W. Xiao, *Phys. Lett. B* **705**, 106 (2011).
 - [28] S. Meissner, A. Metz, and M. Schlegel, *J. High Energy Phys.* **08** (2009) 056.
 - [29] C. Lorce and B. Pasquini, *J. High Energy Phys.* **09** (2013) 138.
 - [30] S. M. Aybat and T. C. Rogers, *Phys. Rev. D* **83**, 114042 (2011).
 - [31] M. G. Echevarria, A. Idilbi, K. Kanazawa, C. Lorce, A. Metz, B. Pasquini, and M. Schlegel, *Phys. Lett. B* **759**, 336 (2016).
 - [32] A. Vladimirov, *J. High Energy Phys.* **04** (2018) 045.
 - [33] A. A. Vladimirov, *Phys. Rev. Lett.* **118**, 062001 (2017).
 - [34] M. G. Echevarria, I. Scimemi, and A. Vladimirov, *Phys. Rev. D* **93**, 054004 (2016).

-
- [35] Y. Li, D. Neill, and H. X. Zhu, [arXiv:1604.00392](#) [Phys. Rev. D (to be published)].
- [36] T. Luebbert, J. Oredsson, and M. Stahlhofen, [J. High Energy Phys.](#) 03 (2016) 168.
- [37] M. G. Echevarria, I. Scimemi, and A. Vladimirov, [J. High Energy Phys.](#) 09 (2016) 004.
- [38] I. Scimemi and A. Vladimirov, [J. High Energy Phys.](#) 08 (2018) 003.