

2022

## Editorial: Advances in Understanding Lateral Blue Carbon Export From Coastal Ecosystems

Kai Xiao

Nengwang Chen

Zhaohui Aleck Wang

Joseph James Tamborski  
*Old Dominion University, jtambors@odu.edu*

Damien Troy Maher

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.odu.edu/oeas\\_fac\\_pubs](https://digitalcommons.odu.edu/oeas_fac_pubs)



Part of the [Oceanography Commons](#), [Terrestrial and Aquatic Ecology Commons](#), and the [Water Resource Management Commons](#)

---

### Original Publication Citation

Xiao, K., Chen, N., Wang, Z. A., Tamborski, J. J., Maher, D. T., & Yu, X. (2022). Editorial: Advances in understanding lateral blue carbon export from coastal ecosystems. *Frontiers in Marine Science*, 9, 1-3, Article 1060958. <https://doi.org/10.3389/fmars.2022.1060958>

This Editorial is brought to you for free and open access by the Ocean & Earth Sciences at ODU Digital Commons. It has been accepted for inclusion in OES Faculty Publications by an authorized administrator of ODU Digital Commons. For more information, please contact [digitalcommons@odu.edu](mailto:digitalcommons@odu.edu).

---

**Authors**

Kai Xiao, Nengwang Chen, Zhaohui Aleck Wang, Joseph James Tamborski, Damien Troy Maher, and Xuan Yu



## OPEN ACCESS

## EDITED AND REVIEWED BY

Angel Borja,  
Technology Center Expert in Marine  
and Food Innovation (AZTI), Spain

## \*CORRESPONDENCE

Xuan Yu  
yuxuan7@mail.sysu.edu.cn

## SPECIALTY SECTION

This article was submitted to  
Marine Ecosystem Ecology,  
a section of the journal  
Frontiers in Marine Science

RECEIVED 04 October 2022

ACCEPTED 21 October 2022

PUBLISHED 27 October 2022

## CITATION

Xiao K, Chen N, Wang ZA,  
Tamborski JJ, Maher DT and Yu X  
(2022) Editorial: Advances in  
understanding lateral blue carbon  
export from coastal ecosystems.  
*Front. Mar. Sci.* 9:1060958.  
doi: 10.3389/fmars.2022.1060958

## COPYRIGHT

© 2022 Xiao, Chen, Wang, Tamborski,  
Maher and Yu. This is an open-access  
article distributed under the terms of  
the [Creative Commons Attribution  
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution  
or reproduction in other forums is  
permitted, provided the original  
author(s) and the copyright owner(s)  
are credited and that the original  
publication in this journal is cited, in  
accordance with accepted academic  
practice. No use, distribution or  
reproduction is permitted which does  
not comply with these terms.

# Editorial: Advances in understanding lateral blue carbon export from coastal ecosystems

Kai Xiao<sup>1</sup>, Nengwang Chen<sup>2</sup>, Zhaohui Aleck Wang<sup>3</sup>,  
Joseph James Tamborski<sup>4</sup>, Damien  
Troy Maher<sup>5</sup> and Xuan Yu<sup>6\*</sup>

<sup>1</sup>State Environmental Protection Key Laboratory of Integrated Surface Water-Groundwater Pollution Control, School of Environmental Science and Engineering, Southern University of Science and Technology, Shenzhen, China, <sup>2</sup>Key Laboratory of the Coastal and Wetland Ecosystems, College of the Environment and Ecology, Xiamen University, Xiamen, China, <sup>3</sup>Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, Woods Hole, MA, United States, <sup>4</sup>Department of Ocean & Earth Sciences, Old Dominion University, Norfolk, VA, United States, <sup>5</sup>School of Environment, Science and Engineering, Southern Cross University, Lismore, NSW, Australia, <sup>6</sup>Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai, China

## KEYWORDS

saltmarsh, mangrove, kelp forests, groundwater, outwelling, carbon cycle, intertidal porewater, tidal flat

## Editorial on the Research Topic

### Advances in understanding lateral blue carbon export from coastal ecosystems

'Blue Carbon' refers to the carbon captured by the coastal systems or ocean and was coined about a decade ago (Nellemann et al., 2009), emphasizing the carbon sequestration capacity of coastal vegetated ecosystems (e.g., macroalgae/kelp, seagrass beds, saltmarshes, and mangroves). These blue carbon systems only cover <0.1% of the ocean area, but may account for >50% of the carbon storage in marine environments, representing a large carbon sink comparable to the global river input (Alongi, 2014). The fluxes of terrestrial-derived carbon including dissolved organic carbon (DOC), dissolved inorganic carbon (DIC), and particulate organic carbon (POC) transported through surface river runoff to the ocean were well known and quantified (Ludwig et al., 1996; Regnier et al., 2022). However, increasing evidence suggests that tidal exchange dominates the transport of significant dissolved carbon from coastal ecosystems to adjacent estuarine and shelf waters (e.g., Maher et al., 2013; Tait et al., 2016; Wang et al., 2016; Chen et al., 2021). This mechanism is commonly named as carbon 'outwelling' or lateral carbon export (e.g., Teal, 1962; Odum, 1968; Wang and Cai, 2004; Sippo et al., 2017; Cabral et al., 2021; Santos et al., 2021; Tamborski et al., 2021).

Outwelling represents a potentially overlooked source of 'blue carbon', which may rival or even exceed the sedimentary burial rates of blue carbon in coastal

ecosystems (Correa et al., 2022), and yet, this pathway remains understudied to date due to a lack of robust, coherent measurements quantifying lateral blue carbon exports in highly dynamic coastal ecosystems. The aim of this Research Topic was to summarize the recent advances in understanding lateral blue carbon export from coastal ecosystems. Five articles were finally collected in this Research Topic as summarized below.

Zhu et al. calculated the porewater exchange rate and fluxes of carbon outwelling and greenhouse gas emissions, based on a  $^{222}\text{Rn}$  mass balance model in a subtropical multi-species saltmarsh in Hangzhou Bay, China. They found that DIC was the most dominant (~90%) carbon species exported through outwelling and porewater exchange, which were 3.2 and 1.2 times that of carbon burial. The emissions of  $\text{CO}_2$  ( $54.6 \pm 0.5 \text{ mmol m}^{-2} \text{ d}^{-1}$ ) and  $\text{CH}_4$  ( $0.19 \pm 0.01 \text{ mmol m}^{-2} \text{ d}^{-1}$ ) from creek water can offset 16% of sedimentary carbon burial. Further, the isotopic signal of  $\delta^{13}\text{C}$  and ratios of C/N reveal that the organic carbon were mainly originated from the C3 plant (i.e., *Scirpus mariqueter*) rather than the C4 plant (i.e., *Spartina alterniflora*). This study emphasized that the porewater-derived carbon outwelling acts as a critical role in the long-term carbon sink, providing a scientific basis for protecting blue carbon ecosystems.

Yuan et al. assessed the annual lateral exchange of organic carbon between the adjacent water and salt marsh at the Yangtze estuary, China, to determine whether the salt marsh acted as a net source or sink for estuarine carbon. They found that the concentrations of DOC and POC peaked in autumn ( $3.54 \text{ mg L}^{-1}$  and  $4.19 \text{ mg L}^{-1}$ , respectively) and declined to the lowest in winter and spring ( $1.87 \text{ mg L}^{-1}$  and  $1.51 \text{ mg L}^{-1}$ , respectively), and their fluxes were significantly correlated in different seasons. In different seasons the tidal creek showed the export of organic carbon with the flux range from  $-12.65$  to  $4.04 \text{ g C m}^{-2}$ . Further, the flux of organic carbon varied with tidal pattern and were significantly higher during spring tides than that during neap tides. Last, this study indicates that the salt marshes acted as a net source through lateral carbon export for the estuary.

Wang et al. investigated the submarine groundwater discharge (SGD) and its effect on the carbon cycle in a highly urbanized and river-dominated coastal area, i.e., the Guangdong-HongKong-Macao Greater Bay Area in China. They found that SGD-derived fluxes of DIC and DOC were  $(0.77\text{--}3.29) \times 10^{10} \text{ g d}^{-1}$  and  $(0.60\text{--}9.94) \times 10^9 \text{ g d}^{-1}$ , respectively, which were nearly ~2 times larger than riverine inputs. Further, they found that SGD acted as a potential source of atmospheric  $\text{CO}_2$  with a flux of  $1.46 \times 10^9 \text{ g C d}^{-1}$ . These additional inputs of carbon and nutrients were expected to enhance biological pump efficiency, stimulate new primary production, and regulate the balance of the carbonate system in marine waters. This study emphasized that SGD is important as rivers, both of which plays a significant role in carbon budgets at the regional and global scales.

Kim et al. explored the spatial patterns of DIC and total alkalinity ( $\text{Alk}_T$ ) productions and DIC in a shallow beach aquifer in Cape Shores, Delaware, USA. They found that the substantial changes of DIC and  $\text{Alk}_T$  can occur along the subsurface

flowpaths due to the anoxic reactions which led to the additional fluxes of DIC ( $191 \text{ mmol d}^{-1}$ ) and  $\text{Alk}_T$  ( $134 \text{ mmol d}^{-1}$ ) to the ocean per meter length of shoreline, respectively. In the saltwater-freshwater mixing zone, the ratios of DIC:  $\text{Alk}_T$  and  $\text{dDIC:dAlk}_T$  relative to the theoretical dilution line revealed that both aerobic and anaerobic reactions can actively contribute to the productions of DIC and  $\text{Alk}_T$  beneath the beach surface. This study suggests that the beach aquifers (as carbon-poor sites) can support the transformation of inorganic carbon and should not be overlooked as an important source of DIC and  $\text{Alk}_T$  like carbon-rich mangrove wetlands.

Pan et al. proposed an effective approach to track the dynamics of detached macrophytes on a semi-sheltered beach in Odense Fjord, Denmark. They conducted a monitoring survey using multiple technologies including real-time camera trap, deep learning with a network architecture and partial least squares regression analysis. The camera trap can be used as a labor-saving approach to track the spatiotemporal dynamics of detached macrophytes deposited on the beach. Further, the application of deep learning provides an important aid in image identification and ecological survey and environmental management.

Overall, these articles presented in this Research Topic represent important progress, datasets, as well as novel methodologies in understanding of lateral carbon export from various coastal ecosystems (e.g., beach, salt marsh, estuary, and coastal bay). We are convinced that this Research Topic will continue to inspire improvement of quantifying lateral carbon fluxes and developing effective strategies for management and protection of the blue carbon ecosystems.

## Author contributions

KX invited the other guest editors NC, ZW, JT, DM and XY to design this Research Topic. All guest editors have edited and reviewed the editorial article, and approved the submitted version.

## Acknowledgments

We would like to thank the authors for their contributions and patience, and the reviewers for their help in the review process. KX thanks the fundings from National Natural Science Foundation of China (No. 41907162), GuangDong Basic and Applied Basic Research Foundation (No. 2022A1515010572), and Shenzhen Science and Technology Program (No. JSGG20210802153535002).

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## References

- Alongi, D. M. (2014). Carbon cycling and storage in mangrove forests. *Ann. Rev. Mar. Sci.* 6 (1), 195–219. doi: 10.1146/annurev-marine-010213-135020
- Cabral, A., Dittmar, T., Call, M., Scholten, J., de Rezende, C. E., Asp, N., et al. (2021). Carbon and alkalinity outwelling across the groundwater-creek-shelf continuum off Amazonian mangroves. *Limnol. Oceanogr. Lett.* 6 (6), 369–378. doi: 10.1002/lol2.10210
- Chen, X., Santos, I. R., Call, M., Reithmaier, G. M. S., Maher, D., Holloway, C., et al. (2021). The mangrove CO<sub>2</sub> pump: Tidally driven pore-water exchange. *Limnol. Oceanogr.* 66 (4), 1563–1577. doi: 10.1002/lno.11704
- Correa, R. E., Xiao, K., Conrad, S., Wadnerkar, P., Wilson, A. M., Sanders, C., et al. (2022). Groundwater carbon exports reduce the sediment carbon burial potential of a salt marsh. *Estuaries Coasts* 45, 1545–1561. doi: 10.1007/s12237-021-01021-1
- Ludwig, W., Amiotte-Suchet, P., and Probst, J. L. (1996). River discharges of carbon to the world's oceans: Determining local inputs of alkalinity and of dissolved and particulate organic carbon. *Comptes Rendus l'Academie Sci. - Ser. Ila Sci. la Terre Des. Planetes* 323, 1007–1014.
- Maher, D. T., Santos, I. R., Golsby-Smith, L., Gleeson, J., and Eyre, B. D. (2013). Groundwater-derived dissolved inorganic and organic carbon exports from a mangrove tidal creek: The missing mangrove carbon sink? *Limnol. Oceanogr.* 58 (2), 475–488. doi: 10.4319/lno.2013.58.2.0475
- Nellemann, C., Corcoran, E., Duarte, C. M., Valdés, L., De Young, C., Fonseca, L., et al. (2009). "Blue carbon: A rapid response assessment," *Blue Carbon* (Arendal, Norway: United Nations Environment Programme, GRID-Arendal).
- Odum, E. P. (1968). "A research challenge: Evaluating the productivity of coastal and estuarine water," in *Proceedings of the Second Sea Grant Conference* (Univ. of Rhode Island) 63–64.
- Regnier, P., Resplandy, L., Najjar, R. G., and Ciais, P. (2022). The land-to-Ocean loops of the global carbon cycle. *Nature* 603, 401–410. doi: 10.1038/s41586-021-04339-9
- Santos, I. R., Burdige, D. J., Jennerjahn, T. C., Bouillon, S., Cabral, A., Serrano, O., et al. (2021). The renaissance of odum's outwelling hypothesis in 'Blue carbon' science. *Estuar. Coast. Shelf Sci.* 255, 107361. doi: 10.1016/j.ecss.2021.107361
- Sippo, J. Z., Maher, D. T., Tait, D. R., Ruiz-Halpern, S., Sanders, C. J., and Santos, I. R. (2017). Mangrove outwelling is a significant source of oceanic exchangeable organic carbon. *Limnol. Oceanogr. Lett.* 2 (1), 1–8. doi: 10.1002/lol2.10031
- Tait, D. R., Maher, D. T., Macklin, P. A., and Santos, I. R. (2016). Mangrove pore water exchange across a latitudinal gradient. *Geophys. Res. Lett.* 43 (7), 3334–3341. doi: 10.1002/2016GL068289
- Tamborski, J. J., Eagle, M., Kurylyk, B. L., Kroeger, K. D., Wang, Z. A., Henderson, P., et al. (2021). Pore water exchange-driven inorganic carbon export from intertidal salt marshes. *Limnol. Oceanogr.* 66, 1774–1792. doi: 10.1002/lno.11721
- Teal, J. M. (1962). Energy-flow in salt-marsh ecosystem of Georgia. *Ecology* 43, 614–624. doi: 10.2307/1933451
- Wang, Z. A., and Cai, W. J. (2004). Carbon dioxide degassing and inorganic carbon export from a marsh-dominated estuary (the duplin river): A marsh CO<sub>2</sub> pump. *Limnol. Oceanogr.* 49, 341–354. doi: 10.4319/lno.2004.49.2.0341
- Wang, Z. A., Kroeger, K. D., Ganju, N. K., Gonneea, M. E., and Chu, S. N. (2016). Intertidal salt marshes as an important source of inorganic carbon to the coastal ocean. *Limnol. Oceanogr.* 61, 1916–1931. doi: 10.1002/lno.10347