

2021

## First Record of Blacknape Large-Eye Bream *Gymnocranius satoi* (Perciformes: Lethrinidae) in the Philippines

Nicko Amor Flores

*National Fisheries Research and Development Institute, Philippines*

Jade Tiffany Rey

*National Fisheries Research and Development Institute, Philippines*

Jeffrey T. Williams

*Smithsonian Institution*

Kent Carpenter

*Old Dominion University, kcarpent@odu.edu*

Mudjekeewis Santos

*National Fisheries Research and Development Institute, Philippines*

Follow this and additional works at: [https://digitalcommons.odu.edu/biology\\_fac\\_pubs](https://digitalcommons.odu.edu/biology_fac_pubs)



Part of the [Aquaculture and Fisheries Commons](#), [Marine Biology Commons](#), and the [Natural Resources and Conservation Commons](#)

---

### Original Publication Citation

Flores, N. A., Rey, J. T., Williams, J. T., Carpenter, K., & Santos, M. (2021). First record of blacknape large eye bream *Gymnocranius satoi* (Perciformes: Lethrinidae) in the Philippines. *Philippine Journal of Systematic Biology*, 15(1), 8 pp. [https://asbp.org.ph/wp-content/uploads/2022/05/PJSB\\_2021-007.pdf](https://asbp.org.ph/wp-content/uploads/2022/05/PJSB_2021-007.pdf)

This Article is brought to you for free and open access by the Biological Sciences at ODU Digital Commons. It has been accepted for inclusion in Biological Sciences Faculty Publications by an authorized administrator of ODU Digital Commons. For more information, please contact [digitalcommons@odu.edu](mailto:digitalcommons@odu.edu).

# First record of blacknape large-eye bream *Gymnocranius satoi* (Perciformes: Lethrinidae) in the Philippines

Nicko Amor Flores<sup>1</sup>, Jade Tiffany Rey<sup>1</sup>, Jeffrey T. Williams<sup>2</sup>, Kent Carpenter<sup>3</sup>, and Mudjekeewis Santos<sup>1\*</sup>

## Abstract

The Philippines has been regarded as the center of the center of marine shorefish biodiversity, having the highest number of fish species per square area in the world. The blacknape large-eye bream, *Gymnocranius satoi*, has been reported to occur from Southern Japan, Taiwan to Northwestern Australia and to the Coral Sea, but has not previously been recorded from the Philippines. From 2011 - 2019, the National Fisheries Research and Development Institute (NFRDI) collaborated with the National Museum of Natural History of the Smithsonian Institution (NMNH/SI), USA, and the Old Dominion University (ODU), Virginia, USA, to inventory all commercial fish species sold in fish markets around the Philippines. During three (3) fish market surveys (Dumaguete City Market, Negros Oriental; Claveria Public Market, Cagayan, Northern Luzon; and Tabaco City Market, Albay, Southeastern Luzon in 2013, 2016 and 2017, respectively), we collected and eventually identified using morphological and DNA barcoding (COI) analysis, seven (7) specimens of *G. satoi*, representing the first records of this species from the country. Since the potential to discover new species and first records of fish species in the Philippines is high, further taxonomic study of the genus *Gymnocranius* is needed.

**Keywords:** Marine biodiversity, conservation, fisheries, DNA barcoding, taxonomy

## Introduction

The Philippines is endowed with highly diverse marine species resulting in its being considered as the epicenter of the world's marine shorefish biodiversity (Carpenter & Springer 2005; Sanciangco *et al.*, 2013). However, the country has also been cited as a hotspot for marine biodiversity conservation (Roberts *et al.*, 2002).

The genus *Gymnocranius* is one of the four genera of the large-eye seabreams or Monotaxinae subfamily of the Lethrinidae family (Sato, 1986; Carpenter & Allen, 1989). *Gymnocranius* currently includes 11 valid described species, and one undescribed species (Carpenter & Allen, 1989; Chen *et al.*, 2016, 2017) but the number could be as high as 15 with additional four (4) possibly undescribed species (Chen & Borsa, 2020).

*Gymnocranius satoi* Borsa, Béarez, Paijo & Chen (Borsa *et al.*, 2013) is known to occur from New Caledonia, West Papua (Raja Ampat), Southern Japan and Taiwan (Borsa *et al.*, 2013; Chen *et al.*, 2016, Chen *et al.*, 2017; Chen & Borsa, 2020). It is possible that it also occurs in the Great Barrier Reef, Solomon Sea, Bismarck Sea, New Guinea and Lesser Sunda Islands (Fricke *et al.*, 2021). A search of the Barcode of Life Data (BOLD) systems (<https://v3.boldsystems.org/>) yielded 2 published and released record of *G. satoi* from Taiwan (GenBank Accession Numbers: MT607083 and MT607084 with BIN Cluster ID BOLD:AAE3729).

The National Fisheries Research and Development Institute (NFRDI) collaborated with the National Museum of Natural History of the Smithsonian Institution (NMNH/SI) USA and the Old Dominion University (ODU), Virginia, USA from 2011 to 2019, to inventory and DNA barcode all fish species sold in fish markets around the country. The purpose of the study was to improve food safety and security by developing the capacity to identify fillets or other fish products that cannot be identified using normal taxonomic methods (Carpenter *et al.*, 2017).

DNA barcoding is a method widely used to identify species using the mitochondrial cytochrome c oxidase subunit 1 gene (COI) (Hebert *et al.*, 2003). It has been used in the

<sup>1</sup>Genetic Fingerprinting Laboratory, National Fisheries Research and Development Institute, 101 Mother Ignacia St., Quezon City, Philippines

<sup>2</sup>Department of Vertebrate Zoology, National Museum of Natural History of the Smithsonian Institution, Washington DC, USA

<sup>3</sup>Old Dominion University, Virginia, USA

\*Corresponding author: mudjiesantos@gmail.com

Date Submitted: 12 April 2021

Date Accepted: 28 August 2021

Philippines to describe novel fish species such as whitechin surgeonfish, *Acanthurus albimento* (Carpenter *et al.*, 2017) and Pogi perchlet, *Chelidoperca santosi* (Williams & Carpenter, 2015). It has been utilized to identify new records of indigenous fishes like Pacific bluefin tuna, *Thunnus orientalis* (Sarmiento *et al.*, 2016) and reef manta ray, *Manta alfredi* (Acebes *et al.*, 2016) as well as those that are non-native to the country such as the Midas cichlid, *Amphilophus citrinellus* (Poniente *et al.*, 2020) and Mayan cichlid *Cichlasoma urophthalmus* (Ordonez *et al.*, 2015). The present paper reports the first record of blacknape large-eye bream, *G. satoi*, from the Philippines using morphological analysis and DNA barcoding.

## Materials and Methods

### Sample collection and preservation


Commercial fish markets and fish landings in Dumaguete City market, Negros Oriental; Claveria Public Market, Cagayan, Northern Luzon; and Tabaco City Market, Albay, Southeastern Luzon were visited in 2013, 2016 and 2017, respectively, as part of the collaborative project of NMNH/SI, ODU and NFRDI to inventory species in fish markets around the country (Fig. 1).

Seven (7) voucher specimens of an unknown *Gymnocranius* species were observed and purchased from local market vendors and fishers (Table 1). Each specimen (voucher) was photographed to capture the fresh color pattern and their muscle tissues sampled in duplicate and preserved in ethanol and M2 buffer of the Autogen prep protocol for genetic analysis. The voucher specimens were then tagged with a unique identifying number, preserved in formalin and stored as reference vouchers at the NMNH/SI. All 7 specimens were labelled with place of origin/collection site and collection date.

### Sample identification

For morphological analysis, each of the 7 *Gymnocranius* sp. collected was identified based on identification keys in FAO Species Identification Guide for Fishery Purposes (Carpenter & Niem, 2001), in FishBase (Froese & Pauly 2010), the Market Fishes of Indonesia (White *et al.* 2013) and specifically the most recent compilation for identification purposes in Table 3 of Chen *et al.*, (2017).

For genetic analysis, DNA extraction and amplification from the 7 muscle tissues were performed following Smithsonian protocols of Baldwin *et al.*, (2011). Briefly, the CO1 sequences were amplified using primers FISH-BCL (5'-TCAACYAATCAYAAAGATATYGGCAC) and FISH-BCH (5'-TAAACTTCAGGGTGACCAAAAATCA) at the following thermal cycling procedure: 1 cycle of 5 min at 95°C; 35 cycles of 30 s at 95°C, 30 s at 52°C, and 45 s at 72°C; 1



**Figure 1.** Map showing the collection sites of *Gymnocranius satoi* in Philippines. Inset map approximates the known distribution of the species in the Indo-West Pacific (IWP).

cycle of 5 min at 72°C; and a hold at 10°C. DNA sequences were obtained using an ABI 3730XL automated DNA sequencer and sequence files were exported into Sequencher 4.7 (GeneCodes, Ann Arbor, MI).

Reference CO1 sequences of *Gymnocranius* spp. (*G. satoi*, *G. superciliosus*, *G. oblongus*, *G. griseus* and *G. obesus*) and of *Lethrinus* spp. (*L. lentjan* and *L. atkinsoni*) as outgroups were obtained from GenBank and BOLD (Fig 3.) Alignment of all CO1 sequences were done using ClustalW. MEGA6 software (Tamura *et al.* 2013) was employed to determine the extent of genetic distance between species by calculating average pairwise comparisons of CO1 sequence differences across all individuals using Kimura 2-Parameter (K2P) method (Kimura, 1980). Maximum Likelihood (ML) method using K2P was done to generate the phylogenetic tree following Carpenter *et al.* (2017).

## Results and Discussion

### Morphological identification

**Observations:** The seven (7) unknown fish specimens (Table 1 and Fig. 2) were initially identified based on their distinct morphological characteristics as belonging to the genus *Gymnocranius* of the family Lethrinidae (Emperors). The specimens had naked cheeks versus scaly compared with the other Lethrinid genera (Carpenter & Niem, 2001). Other characteristics observed included the following: bumpy forehead and the area immediately above eye forms a distinctive brownish to blackish eyebrow; a vertical dark bar crossing the eye; the lower edge of eye is above the body axis; scales above lateral line have dark-grey patch; caudal fin is forked and the lobes are convex on inner side; scales in 3 rows below lateral line are darker. The characteristic reddish to bright vermilion red fins, reddish to red upper lip and white lower lip (Borsa *et al.*, 2013) are unfortunately not visible in the photographs. Market specimens tend to lose color quickly.

**Remarks:** The morpho-anatomy of the unknown *Gymnocranius* specimens as mentioned above (Fig. 2) closely resemble the features of the holotype specimen of *G. satoi* as reported by Borsa *et al.* (2013)- with high body feature; bumpy forehead; shallowly forked caudal fin; flanks that are silvery; with distinctive brownish to blackish eyebrow and visible vertical dark bar crossing the eye. Taken together, these characteristics points to our specimens as *G. satoi*.

### Genetic identification

The COI sequences of the seven (7) putative *G. satoi* samples formed a distinct monophyletic clade with the three (3) *G. satoi* reference sequences using Maximum Likelihood method (Fig. 3). Bootstrap value at the node between the nearest clade (*G. superciliosus*) is significant at over 50 similar to reports of Carpenter *et al.*, 2017. In addition, pairwise genetic distance between the seven (7) putative *G. satoi* and three (3) reference *G. satoi* ranges from 0.002-0.003, which is significantly different from the nearest species *G. supecciliosus* (0.061-0.097) and *G. griseus* (0.067-0.069) (Table 2). Hence, both genetic analysis, like the morpho-anatomical features, strongly suggests that the 7 unknown *Gymnocranius* are in fact *G. satoi*.

### Distribution of *G. satoi*

*G. satoi* was previously reported as “*G. lethrinoides*” by Masudaka *et al.* (1984). It was Torao Sato, a Japanese

ichthyologist who first recognized that the said specimen was distinct from the other *Gymnocranius* species (Sato, 1986). The specimen was finally described as a new species named after Tarao Sato (Borsa *et al.*, 2013; Borsa *et al.*, 2010a, 2010b). *G. satoi* is known to be distributed in the southern lagoon of New Caledonia, Raja Ampat in western West Papua, Southern Japan, Northwestern Australia, Coral Sea and Taiwan (Borsa *et al.*, 2013; Chen *et al.*, 2016; Chen *et al.*, 2017; Chen & Borsa 2020). It could also be present in the Great Barrier Reef, Solomon Sea, Bismarck Sea, New Guinea and Lesser Sunda Islands (Fricke *et al.*, 2021). This study provides the first definitive evidence of *G. satoi* in the Philippines, completing its range distribution in the Indo-West Pacific (IWP). This information is valuable for the species’ global and regional IUCN Red List assessment, conservation and management.


### Conclusion

The discovery of the occurrence of *G. satoi* in the Philippines fills the large geographic gap distribution between Southern Japan and West Papua to Coral Sea, leading to the conclusion that *G. satoi* exhibits continuous distribution and is recurring between Southern Japan and Western Pacific Ocean. Preliminary analysis of COI sequences for fishes we have collected at Philippine fish markets suggest that there may be hundreds of yet to be described new species and first records of fishes occurring in the Philippines. Moreover, rarefaction and extrapolation species curves after 258 sampling events over 6 years indicates that around 1,500 species could eventually be sampled from fish markets in the Philippines (Carpenter *et al.*, 2017). Hence, further taxonomic study of fishes in the genus *Gymnocranius* is needed.

### Acknowledgments

The authors would like to thank Apollo Marco Lizano, Mikel Mendiola, Philip Jessele, Jay Diddiquin, Jin Kevin Dionisio, Clien Ivan Ongkiatko, Jasmine Meren and Criselda Williams, who worked with us tirelessly as field assistants. We also thank the National Stock Assessment Program (NSAP) enumerators who eagerly assisted us in collecting specimens and who guided our way to rural and more remote areas we visited. We want to thank the market vendors who gave us free fish samples even after we offered to pay for them. Special thanks to former BFAR 5 Director Dennis del Socorro and to Dr. Noemi SB. Lanuzela, staff and enumerators, BFAR 6 Director Remia Aparri and Sheryll Mesa, staff and enumerators for providing space to conduct lab work in Tabaco City, Albay and Ilo-ilo City during Panay and and Negros Islands sampling,






**Figure 2.** Photographs of the seven (7) *Gymnocranius satoi* collected from 3 areas in the Philippines (Table 1). **A.** USNM 403315, **B.** USNM 443548, **C.** USNM:FISH:445411, **D.** USNM 445413, **E.** USNM 445415, **F.** USNM:FISH:445417 and **G.** USNM:FISH:445419. Photos by J.T. Williams.

**Table 1.** *Gymnocranius satoi* samples from Philippines with corresponding specimen codes, standard lengths, sampling location, BOLD Barcode Index Numbers (BIN) Identification and GenBank Accession Numbers.

Sample	Voucher specimen code	Standard Length (SL) in cm	Tissue specimen code	Collection sites in Philippines	Barcode of Life Database (BOLD) BIN ID	GenBank Accession Number
A	USNM 403315	17	PI-0315 (PHILA318-13)	Dumaguete City Market, Negros Oriental	BOLD:AAE3729	MZ921405
B	USNM 443548	18.1	PHISH-708 (PHILA-2899-18)	Claveria Public Market, Cagayan, Northern Luzon	BOLD:AAE3729	MZ921404
C	USNM:FISH:445411	20	LUZ-194 (PHILU194-18)	Tabaco City Market, Albay, Southeastern Luzon	BOLD:AAE3729	MZ921403
D	USNM 445413	22.5	LUZ 196 (PHILU196-18)	Tabaco City Market, Albay, Southeastern Luzon	BOLD:AAE3729	MZ921407
E	USNM 445415	23.8	LUZ-198 (PHILU198-18)	Tabaco City Market, Albay, Southeastern Luzon	BOLD:AAE3729	MZ921406
F	USNM:FISH:445417	19	LUZ-200 (PHILU200-18)	Tabaco City Market, Albay, Southeastern Luzon	BOLD:AAE3729	MZ921408
G	USNM:FISH:445419	12.4	LUZ-202 (PHILU202-18)	Tabaco City Market, Albay, Southeastern Luzon	BOLD:AAE3729	MZ921409

**Table 2.** Computed pairwise distances between the samples and reference sequences from GenBank and BOLD databases using the Kimura 2-parameter model. The number of base substitutions per site from between sequences are shown. Standard error estimate(s) are shown above the diagonal.

SAMPLES	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
1 PHILU196-18_ <i>Gymnocranius</i> sp.		0.002	0.000	0.002	0.002	0.002	0.002	0.002	0.002	0.000	0.011	0.011	0.011	0.011	0.012	0.010	0.010	0.011	0.022	0.022
2 PHILU198-18_ <i>Gymnocranius</i> sp.	0.003		0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.011	0.011	0.011	0.011	0.012	0.010	0.010	0.011	0.022	0.022
3 PHILA2899-18_ <i>Gymnocranius</i> sp.	0.000	0.003		0.002	0.002	0.002	0.002	0.002	0.002	0.000	0.011	0.011	0.011	0.011	0.012	0.010	0.010	0.011	0.022	0.022
4 PHILA318-13_ <i>Gymnocranius</i> sp.	0.002	0.002	0.002		0.002	0.002	0.000	0.002	0.002	0.002	0.010	0.010	0.011	0.011	0.012	0.010	0.010	0.011	0.022	0.022
5 PHILU194-18_ <i>Gymnocranius</i> sp.	0.003	0.003	0.003	0.002		0.002	0.002	0.002	0.002	0.002	0.011	0.011	0.011	0.011	0.012	0.010	0.010	0.010	0.022	0.022
6 PHILU200-18_ <i>Gymnocranius</i> sp.	0.003	0.003	0.003	0.002	0.003		0.002	0.002	0.002	0.002	0.011	0.011	0.011	0.011	0.012	0.010	0.010	0.011	0.022	0.022
7 PHILU202-18_ <i>Gymnocranius</i> sp.	0.002	0.002	0.002	0.000	0.002	0.002		0.002	0.002	0.002	0.010	0.010	0.011	0.011	0.012	0.010	0.010	0.011	0.022	0.022
8 GBMNC14689-20_ <i>G. satoi</i>	0.003	0.003	0.003	0.002	0.003	0.003	0.002		0.000	0.002	0.011	0.011	0.011	0.011	0.012	0.010	0.010	0.011	0.022	0.022
9 MT607083.1_ <i>G. satoi</i>	0.003	0.003	0.003	0.002	0.003	0.003	0.002	0.000		0.002	0.011	0.011	0.011	0.011	0.012	0.010	0.010	0.011	0.022	0.022
10 MT607084.1_ <i>G. satoi</i>	0.000	0.003	0.000	0.002	0.003	0.003	0.002	0.003	0.003		0.011	0.011	0.011	0.011	0.012	0.010	0.010	0.011	0.022	0.022
11 MT607076.1_ <i>G. griseus</i>	0.069	0.069	0.069	0.067	0.069	0.069	0.067	0.069	0.069	0.069		0.000	0.011	0.011	0.011	0.011	0.011	0.012	0.022	0.022
12 MT607075.1_ <i>G. griseus</i>	0.069	0.069	0.069	0.067	0.069	0.069	0.067	0.069	0.069	0.069	0.000		0.011	0.011	0.011	0.011	0.011	0.012	0.022	0.022
13 MT607082.1_ <i>G. oblongus</i>	0.075	0.075	0.075	0.073	0.075	0.075	0.073	0.073	0.073	0.075	0.071	0.071		0.002	0.013	0.012	0.012	0.013	0.023	0.022
14 MT607080.1_ <i>G. oblongus</i>	0.077	0.077	0.077	0.075	0.077	0.077	0.075	0.075	0.075	0.077	0.069	0.069	0.002		0.013	0.012	0.012	0.013	0.023	0.022
15 MT607079.1_ <i>G. obesus</i>	0.080	0.080	0.080	0.078	0.080	0.080	0.078	0.076	0.076	0.080	0.076	0.076	0.089	0.087		0.013	0.013	0.013	0.021	0.021
16 MT607085.1_ <i>G. superciliosus</i>	0.063	0.063	0.063	0.061	0.060	0.063	0.061	0.063	0.063	0.063	0.067	0.067	0.080	0.082	0.091		0.002	0.003	0.022	0.022
17 MT607088.1_ <i>G. superciliosus</i>	0.065	0.065	0.065	0.063	0.061	0.065	0.063	0.065	0.065	0.065	0.069	0.069	0.082	0.084	0.093	0.002		0.003	0.022	0.022
18 MT607087.1_ <i>G. superciliosus</i>	0.069	0.069	0.069	0.067	0.065	0.069	0.067	0.069	0.069	0.069	0.073	0.073	0.086	0.088	0.097	0.005	0.007		0.022	0.022
19 MN870430.1_ <i>Lethrinus lentjan</i>	0.224	0.227	0.224	0.227	0.229	0.224	0.227	0.224	0.224	0.224	0.237	0.237	0.243	0.241	0.211	0.224	0.222	0.224		0.016
20 MN870331.1_ <i>Lethrinus atkinsoni</i>	0.216	0.219	0.216	0.219	0.221	0.221	0.219	0.216	0.216	0.216	0.216	0.216	0.223	0.225	0.203	0.223	0.225	0.218	0.122	



**Figure 3.** Maximum-likelihood (ML) Tree of CO1 sequences of *Gymnocranius* spp including the (7) specimens (red box) collected from this study (Table 1) and other species from GenBank (*G. satoi*, *G. superciliosus*, *G. oblongus*, *G. griseus* and *G. obesus*) with corresponding Accession Numbers. GenBank CO1 sequences of *Lethrinus lentjan* and *L. atkinsoni* (also under family Lethrinidae) were included as outgroup. Scale bar refers to substitution per site.

respectively. We appreciate Lee Weigt, Amy Driskell and Jeff Hunt of the Laboratories of Analytical Biology (Smithsonian Institution) for providing support and assistance with logistics and molecular analysis of samples throughout the project. We also thank Carole Baldwin Jerry Finan, Diane Pitassy, Erika Wilbur, Shirleen Smith, Kris Murphy, David Smith and Sandra Raredon of the National Museum of Natural History for their support and assistance. Major funding for this food safety and security project was provided by the United States Food and Drug Administration (FDA) through a program administered by Jonathan Deeds under an ongoing Interagency Agreement between the FDA and the NMNH. This study was part of a project entitled “*Collaboration on the Inventory and DNA Barcoding of Commercial Fishes of the Philippines for Food Safety and Biodiversity*” made possible by a multi-year (2011 - 2019) Memorandum of Agreement (MoA) between the Philippine Department of Agriculture, Bureau of Fisheries and Aquatic Resources-National Fisheries Research and

Development Institute (BFAR-NFRDI), and the National Museum of Natural History of the Smithsonian Institution-Department of Vertebrate Zoology, USA from .

### Literature Cited

- de Leon, A.M., J.J.D. Luangsa – ard, S.C. Karunarathna, K.D. Hyde, R.G. Reyes & T.E.E dela Cruz, 2013. Species listing, distribution, and molecular identification of macrofungi in six Aeta tribal communities in Central Luzon, Philippines. *Mycosphere*, 4 (3): 478 – 494.
- Hawksworth, D.L, 2001. The magnitude of fungal diversity: the 1.5 million species estimate revisited. *Mycological Research*, 105 (12): 1422-1432.
- Acebes, J.M.V, Y. Barr, J.M.R. Pereda & M.D. Santos, 2016. Characteristics of a previously undescribed fishery and habitat for *Manta alfredi* in the Philippines. *Marine Biodiversity Records*. 9: 97.



- Baldwin, C.C, C.I. Castillo, L.A. Weigt & B.C. Victor, 2011. Seven new species within western Atlantic *Starksia* atlantica, *S. lepicoelia*, and *S. sluiteri* (Teleostei, Labrisomidae), with comments on congruence of DNA barcodes and species. *ZooKeys*, 79, 21–72.
- Borsa P, Béarez P, Chen W-J (2010a) *Gymnocranius oblongus* (Teleostei: Lethrinidae), a new large-eye bream species from New Caledonia. *Comptes Rendus Biologies* 333, 241–247.
- Borsa P, Collet A, Carassou L, Ponton D, Chen W-J. (2010b) Multiple nuclear and mitochondrial genotyping identifies emperors and large-eye breams (Teleostei: Lethrinidae) from New Caledonia and reveals new large-eye bream species. *Biochemical Systematics and Ecology* 38, 370–389.
- Borsa, P, P. Béarez, S. Paijo & W.-J. Chen, 2013. *Gymnocranius superciliosus* and *Gymnocranius satoi*, two new large-eye breams (Sparoidea: Lethrinidae) from the Coral Sea and adjacent regions. *Comptes Rendus Biologies* 336: 233–240.
- Carpenter, K.E & G.R. Allen, 1989. FAO species catalogue, vol. 9, Emperor fishes and large-eye breams of the world (family Lethrinidae), an annotated and illustrated catalogue of lethrinid species known to date. FAO Species Synopsis No.125, Rome.
- Carpenter, K.E & V.G. Springer, 2005. The center of the center of marine shore fish biodiversity: the Philippine Islands. *Environmental Biology of Fishes*, 72(4), 467–480.
- Carpenter, K.E, J.T. Williams & M.D. Santos, 2017. *Acanthurus albimento*, a new species of surgeonfish (Acanthuriformes: Acanthuridae) from northeastern Luzon, Philippines, with comments on zoogeography. *Journal of the Ocean Science Foundation*, 25, 33–46
- Carpenter, K.E & V.H. Niem, 2001. The Living Marine Resources of the Western Central Pacific, Bony Fishes Part 3 (Menidae to Pomacentridae), vol. 5, (Ed.), FAO Species Identification Guide for Fishery Purposes, FAO, Rome, 2001, pp. 3004–3050.
- Chen, W.-J, H.C. Ho & P. Borsa, 2016. Taiwanese records of oblong large-eye seabream *Gymnocranius oblongus* (Teleostei: Lethrinidae) and other rare or undetermined large-eye seabreams. *Front. Mar. Sci.* 3, 107. <https://doi.org/10.3389/fmars.2016.00107>
- Chen, W.-J, R. Miki & P. Borsa, 2017. *Gymnocranius obesus*, a new large-eye seabream from the Coral Triangle. *C. R. Biol.* 340, 520–530. doi: 10.1016/j.crvi.2017.08.004
- Chen, W.-J & P. Borsa, 2020. Diversity, phylogeny, and historical biogeography of large-eye seabreams (Teleostei: Lethrinidae). *Molecular Phylogenetics and Evolution*, 151: 106902 doi: <https://doi.org/10.1016/j.ympev.2020.106902>
- Fricke, R., W.N. Eschmeyer & R. Van der Laan (eds), 2021. ESCHMEYER'S CATALOG OF FISHES: GENERA, SPECIES, REFERENCES. (<http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>). Electronic version accessed 21 August 2021.
- Froese, R. & D. Pauly. Editors. 2021. FishBase. World Wide Web electronic publication. [www.fishbase.org](http://www.fishbase.org), version (06/2021).
- Hebert, P.D.N, A. Cywinska, S.L. Ball & J.R. deWaard, 2003. Biological identifications through DNA barcodes. *Proceedings of the Royal Society B: Biological Sciences* 270(1512): 313–321.
- Kimura, M, 1980. A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* 16:111–120.
- Masudaka, H., K. Amaoka, C. Araga, T. Uyeno & T. Yoshino, 1984. The fishes of the Japanese Archipelago, Tokai Univ. Press, Tokyo, xxii+437 p., 370 pl.
- Ordóñez, J.F.F., A. M. J. M. Asis, B. J. Catacutan, J. dela Pena & M. D. Santos, 2015. First report on the occurrence of invasive black-chin tilapia *Sarotherodon melanotheron* (Ruppell, 1852) in Manila Bay and of Mayan cichlid *Cichlasoma urophthalmus* (Gunther, 1892) in the Philippines. *Bioinvasions Records*. Volume 4, Issue 2: 115–124. [dx.doi.org/10.3391/bir.2015.4.2.08](https://doi.org/10.3391/bir.2015.4.2.08).
- Poniente, J.A, J.T. Dela Peña, R.M. Pol, L.H. Zapanta & M.D. Santos, 2020. First report of the non-native Midas cichlid, *Amphilophus citrinellus* (Gunther, 1864), in Laguna de Bay, Philippines. *The Philippine Journal of Fisheries* 26: 55–60.
- Roberts, C.M., C.J. McClean, J.E.N. Veron, J.P. Hawkins, G.R. Allen, D.E. McAllister, C.G. Mittermeier, F.W. Schueler, M. Spalding, F. Wells, C. Vynne & T.B. Werner, 2002. Marine Biodiversity Hotspots and Conservation Priorities for Tropical Reefs. *Science* 295: 1280–1284.
- Sanciangco, J.C., K.E. Carpenter, P.J. Etnoyer & F. Moretzsohn, 2013. Habitat availability and heterogeneity and the Indo-Pacific warm pool as predictors of marine species richness in the tropical Indo-Pacific. *PLoS ONE* 8(2): e56245. doi:10.1371/journal.pone.0056245.
- Sarmiento, K.P, M.M. Ventolero, R. Ramiscal, W. Dela Cruz & M.D. Santos, 2016. First DNA record of Pacific bluefin tuna, *Thunnus orientalis* in Philippine waters. *Marine Biodiversity Records*. 9:25. doi:10.1186/s41200-016-0020-y
- Sato, T, 1986. A systematic review of the sparoid fishes of the subfamily Monotaxinae. In: Uyeno, T., Arai, R., Taniuchi, T., Matsuura, K. (Ed.), Indo-Pacific fish biology:



proceedings of the second international conference on Indo-Pacific fishes. Ichthyological Society of Japan, Tokyo, pp. 602–612.

Tamura, K, G. Stecher, D. Peterson, A. Filipski & S. Kumar, 2013. MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. *Molecular Biology and Evolution*, 2013. 30: 2725-2729.

Williams, J.T & K.E. Carpenter, 2015. A new fish species of the subfamily Serraninae (Perciformes, Serranidae) from the Philippines. *Zootaxa* 3911 (2): 287–293. [dx.doi.org/10.11646/zootaxa.3911.2.10](https://doi.org/10.11646/zootaxa.3911.2.10)

White W.T, P.R. Last, Dharmadi, R. Faizah, U. Chodrijah, B.I. Prisantoso, J.J. Pogonoski, M. Puckridge & S.J.M. Blaber, 2013. Market fishes of Indonesia (= Jenis-jenis ikan di Indonesia). ACIAR Monograph No. 155. Australian Centre for International Agricultural Research: Canberra. 438 pp.