

2019

Multistate Survey of American Dog Ticks (*Dermacentor variabilis*) for *Rickettsia* Species

Joy A. Hecht

Michelle E.J. Allerdice

Elizabeth A. Dykstra

Laura Mastel

Rebecca J. Eisen

See next page for additional authors

Follow this and additional works at: https://digitalcommons.odu.edu/biology_fac_pubs



Part of the [Biology Commons](#), [Parasitic Diseases Commons](#), and the [Parasitology Commons](#)

Original Publication Citation

Hecht, J. A., Allerdice, M. E. J., Dykstra, L. M., Eisen, R. J., Johnson, T. L., Gaff, H. D., Varela-Stokes, A. S., Goddard, J., Pagac, B. B., Paddock, C. D., & Karpathy, S. E. (2019). Multistate survey of American dog ticks (*Dermacentor variabilis*) for *Rickettsia* species. *Vector-Borne and Zoonotic Diseases*, 19(9), 652-657.
<https://doi.org/10.1089/vbz.2018.2415>

This Article is brought to you for free and open access by the Biological Sciences at ODU Digital Commons. It has been accepted for inclusion in Biological Sciences Faculty Publications by an authorized administrator of ODU Digital Commons. For more information, please contact digitalcommons@odu.edu.

Authors

Joy A. Hecht, Michelle E.J. Allerdice, Elizabeth A. Dykstra, Laura Mastel, Rebecca J. Eisen, Tammi L. Johnson, Holly D. Gaff, Andrea S. Varela-Stokes, Jerome Goddard, Benedict B. Pagac, Christopher D. Paddock, and Sandor E. Karpathy



HHS Public Access

Author manuscript

Vector Borne Zoonotic Dis. Author manuscript; available in PMC 2021 January 13.

Published in final edited form as:

Vector Borne Zoonotic Dis. 2019 September ; 19(9): 652–657. doi:10.1089/vbz.2018.2415.

Multistate Survey of American Dog Ticks (*Dermacentor variabilis*) for *Rickettsia* Species

Joy A. Hecht¹, Michelle E.J. Allerdice¹, Elizabeth A. Dykstra², Laura Mastel³, Rebecca J. Eisen⁴, Tammi L. Johnson^{4,*}, Holly D. Gaff⁵, Andrea S. Varela-Stokes⁶, Jerome Goddard⁷, Benedict B. Pagac⁸, Christopher D. Paddock¹, Sandor E. Karpathy¹

¹Rickettsial Zoonoses Branch, National Center for Emerging and Zoonotic Infectious Disease, Centers for Disease Control and Prevention, Atlanta, Georgia. ²Zoonotic Disease Program, Washington State Department of Health, Olympia, Washington. ³Division of Microbiology Lab Services, North Dakota Department of Health, Bismarck, North Dakota. ⁴Bacterial Diseases Branch, National Center for Emerging and Zoonotic Infectious Disease, Centers for Disease Control and Prevention, Ft. Collins, Colorado. ⁵Old Dominion University, Norfolk, Virginia. ⁶Department of Basic Sciences, College of Veterinary Medicine, Mississippi State University, Mississippi State, Mississippi. ⁷Department of Biochemistry, Molecular Biology, Entomology, and Plant Pathology, Mississippi State University, Mississippi State, Mississippi. ⁸Public Health Command-Atlantic, Ft. George G. Meade, Maryland.

Abstract

Dermacentor variabilis, a common human-biting tick found throughout the eastern half and along the west coast of the United States, is a vector of multiple bacterial pathogens. Historically, *D. variabilis* has been considered a primary vector of *Rickettsia rickettsii*, the causative agent of Rocky Mountain spotted fever. A total of 883 adult *D. variabilis*, collected between 2012 and 2017 from various locations in 12 states across the United States, were screened for rickettsial DNA. Tick extracts were evaluated using three real-time PCR assays; an *R. rickettsii*-specific assay, a *Rickettsia bellii*-specific assay, and a *Rickettsia* genus-specific assay. Sequencing of *ompA* gene amplicons generated using a seminested PCR assay was used to determine the rickettsial species present in positive samples not already identified by species-specific real-time assays. A total of 87 (9.9%) tick extracts contained *R. bellii* DNA and 203 (23%) contained DNA of other rickettsial species, including 47 (5.3%) with *Rickettsia montanensis*, 11 (1.2%) with *Rickettsia amblyommatis*, 2 (0.2%) with *Rickettsia rhipicephali*, and 3 (0.3%) with *Rickettsia parkeri*. Only 1 (0.1%) tick extract contained DNA of *R. rickettsii*. These data support multiple other

Address correspondence to: Joy A. Hecht., Rickettsial Zoonoses Branch, National Center for Emerging and Zoonotic, Infectious Disease, Centers for Disease Control and Prevention, Mail Stop H17-3, 1600 Clifton Road NE, Atlanta, GA 30329, jhecht@cdc.gov, .

*Current affiliation: Texas A&M AgriLife Research, Texas A&M University, Uvalde, Texas.

Disclaimer

The views expressed in this article are those of the authors and do not reflect the official policy or position of the Centers for Disease Control and Prevention or the U.S. Government. The authors, as employees of the U.S. Government, conducted the study as part of their official duties.

Author Disclosure Statement

No conflicting financial interests exist.

contemporary studies that indicate infrequent detection of *R. rickettsii* in *D. variabilis* in North America.

Keywords

Dermacentor variabilis; American dog tick; *Rickettsia rickettsii*; *Rickettsia bellii*; *Rickettsia parkeri*; *Rickettsia montanensis*

Introduction

The American dog tick, *Dermacentor variabilis*, is a vector of various viral and bacterial pathogens, most notably *Rickettsia rickettsii*, the causative agent of Rocky Mountain spotted fever (RMSF). This tick is found throughout most of the eastern half and along the west coast of the United States, as well as parts of northern Mexico and southern Canada (Price 1954, Dergousoff et al. 2013, James et al. 2015). The number of reported cases of spotted fever rickettsiosis in the United States steadily increased from 1996 to 2016 (Biggs et al. 2016). At the same time, multiple field surveys of *D. variabilis* revealed a rarity or absence of *R. rickettsii* DNA detected among thousands of American dog ticks (Pretzman et al. 1990, Dergousoff et al. 2009, Stromdahl et al. 2011, Goddard et al. 2014, Gleim et al. 2016, Wood et al. 2016). In this context, infections caused by other less pathogenic *Rickettsia* species could be responsible for many cases of spotted fever rickettsiosis reported as “RMSF” (Openshaw et al. 2010). With *R. rickettsii* detected in less than 1% of the American dog ticks screened (Pretzman et al. 1990, Dergousoff et al. 2009, Stromdahl et al. 2011, Goddard et al. 2014, Wood et al. 2016, Trout Fryxell et al. 2017), it remains unclear why this pathogen is so elusive in *D. variabilis* populations. As suggested by investigations of *D. variabilis* and other *Dermacentor* species (Burgdorfer et al. 1981, Macaluso et al. 2002, Sakai et al. 2014), it is possible that the presence of other *Rickettsia* species affects the infrequency with which *R. rickettsii* is identified in American dog ticks. To further evaluate the occurrence of *R. rickettsii* and other *Rickettsia* species in this common human-biting tick, we screened collections of questing, adult *D. variabilis* from 12 U.S. states using real-time PCR.

Materials and Methods

Tick processing and DNA extraction

Between 2012 and 2017, a total of 883 adult *D. variabilis* ticks were collected from 39 counties and 1 independent city within 12 U.S. states: California, Georgia, Kansas, Kentucky, Maryland, Mississippi, Minnesota, New York, North Dakota, Pennsylvania, Virginia, and Washington. Ticks were identified morphologically using standard taxonomic keys and sent to the CDC preserved in ethanol (70–95%). DNA was extracted from ticks using a DNeasy Blood and Tissue Kit (QIAGEN, Valencia, CA) and eluted into a final volume of 200 μ L per the manufacturer’s instructions. For some specimens, the morphological identification was validated by sequencing of the tick 12S ribosomal RNA gene (Beati and Keirans 2001) or 16S ribosomal RNA gene (Black and Piesman 1994).

Rickettsia screening

All extracts were screened for the presence of *R. rickettsii* and *Rickettsia bellii* DNA using two different real-time PCR assays, an *R. rickettsii*-specific TaqMan assay targeting the gene of hypothetical protein A1G_04230, and an *R. bellii*-specific TaqMan assay targeting *gltA*, the citrate synthase gene (Kato et al. 2013, Hecht et al. 2016). The *R. rickettsii*-specific assay was performed as previously described except that 0.4 μM of each primer and 12.5 μL QuantiTect Multiplex PCR Master Mix (QIAGEN) were used in each reaction. Four microliters of template DNA was used in each *R. rickettsii*-specific reaction and 5 μL of template DNA was used in the *R. bellii*-specific reactions. All real-time PCRs were performed in duplicate on a BioRad CFX 96 thermal cycler with a final reaction volume of 25 μL . We considered samples positive if one of the duplicates had a cycle threshold (Ct) <40. Two sets of negative controls and one set of positive controls were included on each plate, where water was used as the negative nontemplate control and DNA from cultured *R. rickettsii* or an *R. bellii* plasmid were used as positive controls (Hecht et al. 2016).

Additional screening was performed on all 883 tick extracts to identify other rickettsial species present using a *Rickettsia* genus-specific TaqMan real-time assay targeting the *gltA* gene (Denison et al. 2014). The reactions were performed as described using 0.2 μM of each primer and 4 μL of DNA. Samples positive by the *gltA* real-time assay were further screened using a seminested PCR targeting the *ompA* gene of spotted fever group *Rickettsia* (Regnery et al. 1991, Eremeeva et al. 1994, Roux et al. 1996). PCRs were performed using 1 μM of each primer (Rr190.70, Rr190.602, Rr190.701), 10 μL of Taq PCR Master Mix (QIAGEN), 2 μL of sample DNA in the primary reaction or 4 μL of the primary reaction product in the secondary reaction, and water to bring the final reaction volume to 20 μL . DNA amplicons were visualized on 1.5% agarose gels containing 0.1 $\mu\text{g}/\text{mL}$ ethidium bromide. Amplicons were extracted and purified using the Promega Wizard SV Gel and PCR Clean-up System (Promega, Madison, WI). Products were bidirectionally sequenced using a BigDye Terminator v3.1 kit on an ABI 3130xl genetic analyzer (Applied BioSystems, Carlsbad, CA) and assembled using Geneious version 7.0.4. (<http://geneious.com>, Kearse et al. 2012). A BLAST analysis comparing the assembled sequences with sequences available in GenBank (www.ncbi.nlm.nih.gov) identified the *Rickettsia* species present in the tick sample.

Results

Rickettsial DNA was detected using the *Rickettsia* genus-specific *gltA* real-time assay in 203 (23%) of the 883 total DNA extracts. *Rickettsia*-positive tick extracts were identified from each state included in this survey (Table 1). Of these, 75 (36.9%) produced *ompA* amplicons that could be used for sequencing. Attempts to amplify additional gene targets were unsuccessful, with the rickettsial DNA concentration being too low in the other samples for sequencing. Amplicons from 63 of these 75 samples were successfully sequenced, identifying 11 (1.2%) samples positive for DNA of *Rickettsia amblyommatis*, 47 (5.3%) for *Rickettsia montanensis*, 2 (0.2%) for *Rickettsia rhipicephali*, and 3 (0.3%) for *Rickettsia parkeri* (Table 1). There were 87 (9.9%) tick samples positive for *R. bellii*. Only one (0.1%) tick, a male specimen collected from Muhlenberg County in Kentucky, contained

DNA of *R. rickettsii*. This same tick was the only coinfecting tick sample identified, and was found to contain both *R. bellii* and *R. rickettsii* DNA.

Discussion

Specimens evaluated in this study were procured from general acarological surveys rather than collection efforts that specifically targeted *D. variabilis*. For this reason, sample sizes varied greatly and were often represented by fewer than 50 ticks from a particular county, precluding generalizations for individual regions or states. Nonetheless, we detected DNA of *R. rickettsii* in only 1 (0.1%) of the 883 *D. variabilis* ticks sampled from a total of 40 jurisdictions in 12 U.S. states. By comparison, we identified rickettsial agents of lesser or unknown pathogenicity, including *R. bellii*, *R. montanensis*, *R. parkeri*, and *R. rhipicephali* in 150 (17%) of the specimens.

The infrequent detection of *R. rickettsii* among questing adult American dog ticks is consistent with many other contemporary studies that either failed to identify this pathogen in *D. variabilis* or found very low rates of infection (Wikswa et al. 2008, Dergousoff et al. 2009, Moncayo et al. 2010, Williamson et al. 2010, Fritzen et al. 2011, Stromdahl et al. 2011, Goddard et al. 2014, Henning et al. 2014, Nadolny et al. 2014, Pagac et al. 2014, Gleim et al. 2016, Mitchell et al. 2016, Wood et al. 2016). Despite the low infection frequency of *R. rickettsii* in *D. variabilis* ticks collected in nature, this tick has been shown to be an efficient vector for *R. rickettsii* (Maver 1911, Burgdorfer 1975) and several isolates of *R. rickettsii* have been made from wild *D. variabilis*, proving that this tick does indeed become infected with *R. rickettsii* in the environment (Cox 1941, Karpathy et al. 2007). In addition, many of the human RMSF cases reported each year in the United States are from areas where *D. variabilis* is endemic (Biggs et al. 2016). Previously, it was reported that *R. rickettsii* has a lethal effect on immature *Dermacentor andersoni* ticks that significantly diminished the number of infected adult ticks (Niebylski et al. 1999), and it was proposed that this deleterious effect may explain the low prevalence of infected ticks in nature. However, Schumacher et al. (2016) recently demonstrated that in the laboratory *R. rickettsii* had no detrimental effects on the survival of infected *D. variabilis* ticks at any life stage.

Approximately 10% of the ticks evaluated in this study contained DNA of *R. bellii*, although the frequency of infection varied considerably state to state. For example, 88% of the 69 ticks collected from one site in Yolo County in northern California contained *R. bellii*, whereas none of the 196 ticks collected from Kansas, Maryland, Minnesota, New York, and North Dakota were infected with this species. *R. bellii* is often missed in prevalence studies because many of the molecular assays used to screen for *Rickettsia* target the rickettsial *ompA* gene, which is absent in *R. bellii* (Ogata et al. 2006). Inclusion of species-specific screening for *R. bellii* provides a more complete picture of the rickettsial species present within these tick populations and enabled us to identify the dually infected tick from Kentucky (Table 1). This could be important when considering the infrequency of *R. rickettsii* in some populations of *D. variabilis*. One mechanism by which *R. bellii* is maintained in tick populations is by transovarial transmission from an infected female to her offspring, and a primary *R. bellii* infection has been shown to inhibit the transovarial transmission of a secondarily acquired *Rickettsia* species (Sakai et al. 2014). Therefore, the

high frequency of *R. bellii* in California may play a significant inhibitory role in the maintenance of other pathogenic *Rickettsia* species and reduce significantly the occurrence of *R. rickettsii* in *D. variabilis* populations in that area. Although *R. bellii* may play an important inhibitory role within this tick population in Yolo County, California, this is a regionally specific finding that does not apply to the other tick populations where no *R. bellii* was identified.

R. montanensis has also been found to inhibit transovarial transmission of a secondarily acquired *Rickettsia* species in *D. variabilis* ticks (Macaluso et al. 2002). Similar to *R. bellii*, the frequency of *R. montanensis* varied greatly among tick samples, from ~32% of the 72 ticks collected from 2 counties in Minnesota to only 2.9% among the remaining 811 specimens collected from 11 other states. Infection rates of between 0% and 10.5% have been described in previously published studies (Wikswa et al. 2008, Dergousoff et al. 2009, Moncayo et al. 2010, Williamson et al. 2010, Fritzen et al. 2011, Stromdahl et al. 2011, Henning et al. 2014, Nadolny et al. 2014, Pagac et al. 2014, Gleim et al. 2016, Mitchell et al. 2016, Wood et al. 2016). *R. montanensis*, a potential human pathogen (McQuiston et al. 2012), may play an inhibitory role in the maintenance of other pathogenic *Rickettsia* species in these Minnesota tick populations.

R. amblyommatis was identified in ~1–8% of the ticks collected from Kentucky, New York, and Virginia, and absent from 533 specimens collected from the remaining nine states, notably the northern and western states of California, Minnesota, North Dakota, and Washington where the ranges of *Amblyomma americanum*, the primary host of *R. amblyommatis* in the United States does not extend. Although *D. variabilis* is not the primary host for *R. amblyommatis*, it has been previously identified in 0.4–2.5% of ticks collected from Georgia, Kentucky, Tennessee, Texas, and Virginia (Moncayo et al. 2010, Williamson et al. 2010, Fritzen et al. 2011, Stromdahl et al. 2011, Henning et al. 2014, Gleim et al. 2016), and one recent study suggests that *R. amblyommatis* may cause mild or subclinical infections in humans (Apperson et al. 2008).

R. rhipicephali was identified in only two specimen: one from North Dakota and one from Washington. To our knowledge this rickettsial species has not been previously reported in *D. variabilis* from these states. *R. parkeri* was identified in ~2.4% of the ticks collected from Muhlenberg County, Kentucky. *R. parkeri* is the causative agent of a spotted fever rickettsiosis found throughout the southern United States and transmitted predominantly by the Gulf Coast tick, *Amblyomma maculatum* (Paddock and Goddard 2015). Previous surveys have identified *R. parkeri* in *D. variabilis* ticks from Kentucky (0.6%), Virginia (0.7%), and Texas (2.3%) (Williamson et al. 2010, Fritzen et al. 2011, Henning et al. 2014), and recently *D. variabilis* has been shown to both acquire and transstadially transmit *R. parkeri* in a laboratory setting (Harris et al. 2017). At this time, it remains unclear what role this tick species plays in transmitting *R. parkeri* in nature; nonetheless, in the regions of the United States where *R. parkeri* has been identified, the frequencies with which this pathogen occurs in *D. variabilis* exceeds those observed for *R. rickettsii* (Williamson et al. 2010, Fritzen et al. 2011, Henning et al. 2014).

The well-documented scarcity of *R. rickettsii* in *D. variabilis* populations calls into question just how significant a role this tick plays in the maintenance of *R. rickettsii*; it thus becomes important to consider the role of other tick species in the disease ecology of RMSF. The most recent outbreak of RMSF in the United States occurred in eastern Arizona where the tick species *Rhipicephalus sanguineus* sensu lato was identified as the vector of *R. rickettsii* (Demma et al. 2005). This tick species found throughout the United States is typically not an aggressive human-biting tick; however, incidents of *Rh. sanguineus* biting humans have been reported in Alabama, Georgia, Kansas, Kentucky, Louisiana, Maryland, New Jersey, North Carolina, Oklahoma, Pennsylvania, Rhode Island, South Carolina, Texas, and Virginia (Estrada-Pena and Jongejan 1999, Stromdahl et al. 2011). Of note, North Carolina is the state with the highest number of reported spotted fever group rickettsioses cases from 2000 to 2012 (Openshaw et al. 2010, Drexler et al. 2016).

Other tick species for consideration as a bridging vector include *Amblyomma americanum* and *Haemaphysalis leporispalustris*. Parker reported isolating *R. rickettsii* from *H. leporispalustris*, collected from the Bitter Root Valley of Montana in 1951 (Parker et al. 1951), and it was later isolated from *H. leporispalustris* collected from Costa Rica (Fuentes et al. 1985). Although this tick has not been associated with any human cases in the United States and is generally not considered a frequent human biter (Merten and Durden 2000), *H. leporispalustris* has been identified as a competent vector of *R. rickettsii* under laboratory conditions (Freitas et al. 2009) and could play a role in enzootic maintenance of *R. rickettsii*. Lastly, *A. americanum* is not only a competent vector of *R. rickettsii* under laboratory conditions (Levin et al. 2017), it has also been implicated as the vector in a human case of RMSF in North Carolina (Breitschwerdt et al. 2011). Although reports of *R. rickettsii* in U.S. populations of *A. americanum* are scarce (Berrada et al. 2011), this aggressive human-biting tick species is found throughout the eastern half of the United States and regularly accounts for a majority of the ticks removed from humans in surveys of southeastern states (Estrada-Pena and Jongejan 1999, Gleim et al. 2016, Mitchell et al. 2016). The role of these other tick species in the maintenance of *R. rickettsii* may be greater than previously thought and our efforts to better understand the disease ecology of RMSF should include further investigation into alternative tick vectors.

Acknowledgments

The authors thank Melissa Bell, Gail Moraru (Mississippi State University), Roman Ganta (College of Veterinary Medicine, Kansas State University), David Kangiser, Susan Brush, Lauren Sherman, Michelle Jodziewicz, and Ashlee-Rose Ferguson (Washington State Department of Health) for their assistance in collecting or contributing specimens of *D. variabilis* evaluated in this study. This study was made possible by funding from the Centers for Disease Control and Prevention.

References

- Apperson CS, Engber B, Nicholson WL, Mead DG, et al. Tickborne diseases in North Carolina: Is “*Rickettsia amblyommii*” a possible cause of rickettsiosis reported as Rocky Mountain spotted fever? *Vector Borne Zoonotic Dis* 2008; 8:597–606. [PubMed: 18447622]
- Beati L, Keirans JE. Analysis of the systematic relationships among ticks of the genera *Rhipicephalus* and *Boophilus* (Acari: Ixodidae) based on mitochondrial 12S ribosomal DNA gene sequences and morphological characters. *J Parasitol* 2001; 87: 32–48. [PubMed: 11227901]

- Berrada ZL, Goethert HK, Cunningham J, Telford SR. *Rickettsia rickettsii* (Rickettsiales: Rickettsiaceae) in *Amblyomma americanum* (Acari: Ixodidae) From Kansas. *J Med Entomol* 2011; 48:461–467. [PubMed: 21485390]
- Biggs HM, Behravesh CB, Bradley KK, Dahlgren FS, et al. Diagnosis and management of tickborne rickettsial diseases: Rocky Mountain spotted fever and other spotted fever group rickettsioses, ehrlichioses and anaplasmosis—United States. *MMWR Morb Mortal Wkly Rep* 2016; 65:1–44. [PubMed: 26766396]
- Black WC, Piesman J. Phylogeny of hard- and soft-tick taxa (Acari: Ixodida) based on mitochondrial 16S rDNA sequences. *Proc Natl Acad Sci U S A* 1994; 91:10034–10038.
- Breitschwerdt EB, Hegarty BC, Maggi RG, Lantos PM, et al. *Rickettsia rickettsii* transmission by a lone star tick, North Carolina. *Emerg Infect Dis* 2011; 17:873–875. [PubMed: 21529399]
- Burgdorfer W A review of Rocky Mountain spotted fever (tickborne typhus), its agent, and its tick vectors in the United States. *J Med Entomol* 1975; 12:269–278. [PubMed: 810584]
- Burgdorfer W, Hayes SF, Mavros AJ. Nonpathogenic rickettsiae in *Dermacentor andersoni*: A limiting factor for the distribution of *Rickettsia rickettsii* In: Burgdorfer W, Anacker RL, eds. *Rickettsiae and Rickettsial Diseases*. New York, NY: Academic Press, 1981:585–594.
- Cox HR. Cultivation of Rickettsiae of the Rocky Mountain spotted fever, typhus and Q fever groups in the embryonic tissues of developing chicks. *Science* 1941; 94:399–403. [PubMed: 17798222]
- Demma LJ, Traeger MS, Nicholson WL, Paddock CD, et al. Rocky Mountain spotted fever from an unexpected tick vector in Arizona. *N Engl J Med* 2005; 353:587–594. [PubMed: 16093467]
- Denison AM, Amin BD, Nicholson WL, Paddock CD. Detection of *Rickettsia rickettsii*, *Rickettsia parkeri*, and *Rickettsia akari* in skin biopsy specimens using a multiplex real-time polymerase chain reaction assay. *Clin Infect Dis* 2014; 59: 635–642. [PubMed: 24829214]
- Dergousoff SJ, Gajadhar AJA, Chilton NB. Prevalence of *Rickettsia* species in Canadian populations of *Dermacentor andersoni* and *D. variabilis*. *Appl Environ Microbiol* 2009; 75:1786–1789. [PubMed: 19151178]
- Dergousoff SJ, Galloway TD, Lindsay LR, Curry PS, et al. Range expansion of *Dermacentor variabilis* and *Dermacentor andersoni* (Acari: Ixodidae) near their northern distributional limits. *J Med Entomol* 2013; 50:510–520. [PubMed: 23802445]
- Drexler NA, Dahlgren FS, Heitman KN, Massung RF, et al. National surveillance of spotted fever group rickettsioses in the United States, 2008–2012. *Am J Trop Med Hyg* 2016; 94:26–34. [PubMed: 26324732]
- Eremeeva M, Yu X, Raoult D. Differentiation among spotted fever group rickettsiae species by analysis of restriction fragment length polymorphism of PCR-amplified DNA. *J Clin Microbiol* 1994; 32:803–810. [PubMed: 7910831]
- Estrada-Peña A, Jongejan F. Ticks feeding on humans: A review of records on human-biting Ixodea with reference to pathogen transmission. *Exp Appl Acarol* 1999; 23:685–715. [PubMed: 10581710]
- Freitas LHT, Faccini JLH, Labruna MB. Experimental infection of the rabbit tick, *Haemaphysalis leporispalustris*, with the bacterium *Rickettsia rickettsii*, and comparative biology of infected and uninfected tick lineages. *Exp Appl Acarol* 2009; 47:321–345. [PubMed: 19067185]
- Fritzen CM, Huang J, Westby K, Freye JD, et al. Infection prevalences of common tick-borne pathogens in adult lone star ticks (*Amblyomma americanum*) and American dog ticks (*Dermacentor variabilis*) in Kentucky. *Am J Trop Med Hyg* 2011; 85:718–723. [PubMed: 21976578]
- Fuentes L, Calderón A, Hun L. Isolation and identification of *Rickettsia rickettsii* from the rabbit tick (*Haemaphysalis leporispalustris*) in the Atlantic zone of Costa Rica. *Am J Trop Med Hyg* 1985; 34:564–567. [PubMed: 3923853]
- Gleim ER, Garrison LE, Vello MS, Savage MY, et al. Factors associated with tick bites and pathogen prevalence in ticks parasitizing humans in Georgia, USA. *Parasit Vectors* 2016; 9:125. [PubMed: 26935205]
- Goddard J, Goltz L, Edwards KT, Smith W, et al. Survey of adult *Dermacentor variabilis* (Say) (Acari: Ixodidae) collected in northern Mississippi for spotted fever group rickettsiae. *Midsouth Entomologist* 2014; 7:8–14.

- Harris EK, Verhoeve VI, Banajee KH, Macaluso JA, et al. Comparative vertical transmission of *Rickettsia* by *Dermacentor variabilis* and *Amblyomma maculatum*. *Ticks Tick Borne Dis* 2017; 8:598–604. [PubMed: 28433729]
- Hecht JA, Allerdice ME, Krawczak FS, Labruna MB, et al. Development of a *Rickettsia bellii*-specific TaqMan assay targeting the citrate synthase gene. *J Med Entomol* 2016; 53:1492–1495. [PubMed: 27473178]
- Henning TC, Orr JM, Smith JD, Arias JR, et al. Spotted fever group rickettsiae in multiple hard tick species from Fairfax County, Virginia. *Vector Borne Zoonotic Dis* 2014; 14:482–485. [PubMed: 24978651]
- James AM, Burdett C, McCool MJ, Fox A, et al. The geographic distribution and ecological preferences of the American dog tick, *Dermacentor variabilis* (Say), in the U.S.A. *Med Vet Entomol* 2015; 29:178–188. [PubMed: 25684582]
- Karpathy SE, Dasch GA, Ereemeeva ME. Molecular typing of isolates of *Rickettsia rickettsii* by use of DNA sequencing of variable intergenic regions. *J Clin Microbiol* 2007; 45:2545–2553. [PubMed: 17553977]
- Kato CY, Chung IH, Robinson LK, Austin AL, et al. Assessment of real-time PCR assay for detection of *Rickettsia* spp. and *Rickettsia rickettsii* in banked clinical samples. *J Clin Microbiol* 2013; 51:314–317. [PubMed: 23135935]
- Kearse M, Moir R, Wilson A, Stones-Havas S, et al. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 2012; 28:1647–1649. [PubMed: 22543367]
- Levin ML, Zemtsova GE, Killmaster LF, Snellgrove A, et al. Vector competence of *Amblyomma americanum* (Acari: Ixodidae) for *Rickettsia rickettsii*. *Ticks Tick Borne Dis* 2017; 8:615–622. [PubMed: 28433728]
- Macaluso KR, Sonenshine DE, Ceraul SM, Azad AF. Rickettsial infection in *Dermacentor variabilis* (Acari: Ixodidae) inhibits transovarial transmission of a second *Rickettsia*. *J Med Entomol* 2002; 39:809–813. [PubMed: 12495176]
- Maver MB. Transmission of spotted fever by other than Montana and Idaho ticks. *J Infect Dis* 1911; 8:322–326.
- McQuiston JH, Zemtsova G, Perniciaro J, Hutson M, et al. Afebrile spotted fever group *Rickettsia* infection after a bite from a *Dermacentor variabilis* tick infected with *Rickettsia montanensis*. *Vector Borne Zoonotic Dis* 2012; 12:1059–1061. [PubMed: 23153005]
- Merten HA, Durden LA. A state-by-state survey of ticks recorded from humans in the United States. *J Vector Ecol* 2000; 25:102–113. [PubMed: 10925803]
- Mitchell EA, Williamson PC, Billingsley PM, Seals JP, et al. Frequency and distribution of rickettsiae, borreliae, and ehrlichiae detected in human-parasitizing ticks, Texas, USA. *Emerg Infect Dis* 2016; 22:312–315. [PubMed: 26811941]
- Moncayo AC, Cohen SB, Fritzen CM, Huang E, et al. Absence of *Rickettsia rickettsii* and occurrence of other spotted fever group rickettsiae in ticks from Tennessee. *Am J Trop Med Hyg* 2010; 83:653–657. [PubMed: 20810834]
- Nadolny RM, Wright CL, Sonenshine DE, Hynes WL, et al. Ticks and spotted fever group rickettsiae of southeastern Virginia. *Ticks Tick Borne Dis* 2014; 5:53–57. [PubMed: 24201057]
- Niebylski ML, Peacock MG, Schwan TG. Lethal effect of *Rickettsia rickettsii* on its tick vector (*Dermacentor andersoni*). *Appl Environ Microbiol* 1999; 65:773–778. [PubMed: 9925615]
- Ogata H, La Scola B, Audic S, Renesto P, et al. Genome sequence of *Rickettsia bellii* illuminates the role of amoebae in gene exchanges between intracellular pathogens. *PLoS Genet* 2006; 2:e76.
- Openshaw JJ, Swerdlow DL, Krebs JW, Holman RC, et al. Rocky Mountain spotted fever in the United States, 2000–2007: Interpreting contemporary increases in incidence. *Am J Trop Med Hyg* 2010; 83:174–182. [PubMed: 20595498]
- Paddock CD, Goddard J. The evolving medical and veterinary importance of the Gulf Coast tick (*Acari: Ixodidae*). *J Med Entomol* 2015; 52:230–252. [PubMed: 26336308]
- Pagac BB, Miller MK, Mazzei MC, Nielsen DH, et al. *Rickettsia parkeri* and *Rickettsia montanensis*, Kentucky and Tennessee, USA. *Emerg Infect Dis* 2014; 20:1750–1752. [PubMed: 25271771]

- Parker RR, Pickens EG, Lackman DB, Bell EJ, et al. Isolation and characterization of Rocky Mountain spotted fever rickettsiae from the rabbit tick *Haemaphysalis leporis-palustris* Packard. *Public Health Rep* 1951; 66:455–463. [PubMed: 14816519]
- Pretzman C, Daugherty N, Poetter K, Ralph D. The distribution and dynamics of *Rickettsia* in the tick population of Ohio. *Ann N Y Acad Sci* 1990; 590:227–236. [PubMed: 2378449]
- Price WH. The epidemiology of Rocky Mountain spotted fever II. Studies on the biological survival mechanism of *Rickettsia rickettsii*. *Am J Hyg* 1954; 60:292–319. [PubMed: 13207101]
- Regnery RL, Spruill CL, Plikaytis BD. Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *J Bacteriol* 1991; 173:1576–1589. [PubMed: 1671856]
- Roux V, Fournier PE, Raoult D. Differentiation of spotted fever group rickettsiae by sequencing and analysis of restriction fragment length polymorphism of PCR-amplified DNA of the gene encoding the protein rOmpA. *J Clin Microbiol* 1996; 34: 2058–2065. [PubMed: 8862558]
- Sakai RK, Costa FB, Ueno TEH, Ramirez DG, et al. Experimental infection with *Rickettsia rickettsii* in an *Amblyomma dubitatum* tick colony, naturally infected by *Rickettsia bellii*. *Ticks Tick Borne Dis* 2014; 5:917–923. [PubMed: 25108783]
- Schumacher L, Snellgrove A, Levin ML. Effect of *Rickettsia rickettsii* (Rickettsiales: Rickettsiaceae) infection on the biological parameters and survival of its tick vector-*Dermacentor variabilis* (Acari: Ixodidae). *J Med Entomol* 2016; 53:172–176. [PubMed: 26494822]
- Stromdahl EY, Jiang J, Vince M, Richards AL. Infrequency of *Rickettsia rickettsii* in *Dermacentor variabilis* removed from humans, with comments on the role of other human-biting ticks associated with spotted fever group rickettsiae in the United States. *Vector Borne Zoonotic Dis* 2011; 11:969–977. [PubMed: 21142953]
- Trout Fryxell RT, Hendricks BM, Pompo K, Mays SE, et al. Investigating the adult Ixodid tick populations and their associated *Anaplasma*, *Ehrlichia*, and *Rickettsia* bacteria at a Rocky Mountain spotted fever hotspot in western Tennessee. *Vector Borne Zoonotic Dis* 2017; 17:527–538. [PubMed: 28598270]
- Wikswa ME, Hu R, Dasch GA, Krueger L, et al. Detection and identification of spotted fever group rickettsiae in *Dermacentor* Species from Southern California. *J Med Entomol* 2008; 45:509–516. [PubMed: 18533446]
- Williamson PC, Billingsley PM, Teltow GJ, Seals JP, et al. *Borrelia*, *Ehrlichia*, and *Rickettsia* spp. in ticks removed from persons, Texas, USA. *Emerg Infect Dis* 2010; 16:441–446. [PubMed: 20202419]
- Wood H, Dillon L, Patel SN, Ralevski F. Prevalence of *Rickettsia* species in *Dermacentor variabilis* ticks from Ontario, Canada. *Ticks Tick Borne Dis* 2016; 7:1044–1046. [PubMed: 27318438]

Table 1.

RICKETTSIA SCREENING RESULTS DIVIDED BY STATE

State of collections (no. of counties/ unincorporated cities surveyed) ^a	Year(s) of collection	Total no. of ticks positive for rickettsial DNA/ total no. of ticks tested (%)	Total No. of ticks positive (%)					
			Rickettsia amblyommatis	Rickettsia montanensis	Rickettsia rhipicephali	Rickettsia parkeri	Rickettsia bellii	Rickettsia rickettsii
California (1)	2013, 2015	32/69 (46%)	0	0	0	0	61 (88.4%)	0
Georgia (5)	2015, 2017	7/71 (9.8%)	0	3 (4.2%)	0	0	2 (2.8%)	0
Kansas (1)	2012	48/86 (56%)	0	9 (10.5%)	0	0	0	0
Kentucky (2)	2014–2016	23/124 (19%)	1 (0.8%)	3 (2.4%)	0	3 (2.4%)	11 (8.9%)	1 (0.8%)
Maryland (1)	2014	1/3 (33%)	0	0	0	0	0	0
Minnesota (2)	2015	26/72 (36%)	0	23 (31.9%)	0	0	0	0
Mississippi (2)	2013–2015	2/51 (3.9%)	0	0	0	0	1 (2.0%)	0
New York (1)	2014–2015	2/12 (17%)	1 (8.3%)	1 (8.3%)	0	0	0	0
North Dakota (11)	2016	2/23 (8.7%)	0	1 (4.3%)	1 (4.3%)	0	0	0
Pennsylvania (1)	2014–2015	21/55 (38%)	0	2 (3.6%)	0	0	1 (1.8%)	0
Virginia (2)	2014–2015	25/214 (12%)	9 (4.2%)	3 (1.4%)	0	0	4 (1.9%)	0
Washington (11)	2012, 2014–2015	14/103 (14%)	0	2 (1.9%)	1 (1.0%)	0	7 (6.8%)	0
Total		203/883 (23%)	11 (1.2%)	47 (5.3%)	2 (0.2%)	3 (0.3%)	87 (9.9%)	1 (0.1%)

^aA list of the counties: California (Yolo), Georgia (Cobb, Coweta, Douglas, Gwinnett, Rockdale), Kansas (Osage), Kentucky (Edmonson, Muhlenberg), Maryland (Frederick), Mississippi (Oktibbeha), Minnesota (Beltrami, Hubbard), New York (Suffolk), North Dakota (Bottineau, Burleigh, Grant, Kidder, McHenry, Mercer, Mountrail, Nelson, Pierce, Walsh, Ward), Pennsylvania (Adams), Virginia (City of Chesapeake, Prince William), and Washington (Benton, Franklin, Grant, Kittitas, Klickitat, Lincoln, Okanogan, Spokane, Yakima, Walla Walla, Whitman).