

4-2014

Growth and Nitrogen Uptake Kinetics in Cultured *Procentrum donghaiense*

Zhangxi Hu

Chinese Academy of Sciences, Qingdao, China

Shunshan Duan

Jinan University, Guangzhou, Guangdong, China

Margaret R. Mulholland

Old Dominion University, mmulholl@odu.edu

Follow this and additional works at: https://digitalcommons.odu.edu/oeas_fac_pubs

 Part of the [Geochemistry Commons](#), [Marine Biology Commons](#), and the [Oceanography and Atmospheric Sciences and Meteorology Commons](#)

Repository Citation

Hu, Zhangxi; Duan, Shunshan; and Mulholland, Margaret R., "Growth and Nitrogen Uptake Kinetics in Cultured *Procentrum donghaiense*" (2014). *OEAS Faculty Publications*. 19.
https://digitalcommons.odu.edu/oeas_fac_pubs/19

Original Publication Citation

Hu, Z. X., Duan, S. S., Xu, N., & Mulholland, M. R. (2014). Growth and nitrogen uptake kinetics in cultured *Prorocentrum donghaiense*. *PLoS One*, 9(4). doi: 10.1371/journal.pone.0094030



Growth and Nitrogen Uptake Kinetics in Cultured *Prorocentrum donghaiense*

Zhangxi Hu^{1,2}, Shunshan Duan², Ning Xu^{2*}, Margaret R. Mulholland^{3*}

1 Key Laboratory of Marine Ecology and Environmental Sciences, Institute of Oceanology, Chinese Academy of Sciences, Qingdao, China, **2** Research Center of Hydrobiology, Jinan University, Guangzhou, Guangdong, China, **3** Department of Ocean, Earth and Atmospheric Sciences, Old Dominion University, Norfolk, Virginia, United States of America

Abstract

We compared growth kinetics of *Prorocentrum donghaiense* cultures on different nitrogen (N) compounds including nitrate (NO_3^-), ammonium (NH_4^+), urea, glutamic acid (glu), dialanine (diala) and cyanate. *P. donghaiense* exhibited standard Monod-type growth kinetics over a range of N concentrations ($0.5\text{--}500\ \mu\text{mol N L}^{-1}$ for NO_3^- and NH_4^+ , $0.5\text{--}50\ \mu\text{mol N L}^{-1}$ for urea, $0.5\text{--}100\ \mu\text{mol N L}^{-1}$ for glu and cyanate, and $0.5\text{--}200\ \mu\text{mol N L}^{-1}$ for diala) for all of the N compounds tested. Cultures grown on glu and urea had the highest maximum growth rates (μ_m , $1.51\pm 0.06\ \text{d}^{-1}$ and $1.50\pm 0.05\ \text{d}^{-1}$, respectively). However, cultures grown on cyanate, NO_3^- , and NH_4^+ had lower half saturation constants (K_m , $0.28\text{--}0.51\ \mu\text{mol N L}^{-1}$). N uptake kinetics were measured in NO_3^- -deplete and -replete batch cultures of *P. donghaiense*. In NO_3^- -deplete batch cultures, *P. donghaiense* exhibited Michaelis-Menten type uptake kinetics for NO_3^- , NH_4^+ , urea and algal amino acids; uptake was saturated at or below $50\ \mu\text{mol N L}^{-1}$. In NO_3^- -replete batch cultures, NH_4^+ , urea, and algal amino acid uptake kinetics were similar to those measured in NO_3^- -deplete batch cultures. Together, our results demonstrate that *P. donghaiense* can grow well on a variety of N sources, and exhibits similar uptake kinetics under both nutrient replete and deplete conditions. This may be an important factor facilitating their growth during bloom initiation and development in N-enriched estuaries where many algae compete for bioavailable N and the nutrient environment changes as a result of algal growth.

Citation: Hu Z, Duan S, Xu N, Mulholland MR (2014) Growth and Nitrogen Uptake Kinetics in Cultured *Prorocentrum donghaiense*. PLoS ONE 9(4): e94030. doi:10.1371/journal.pone.0094030

Editor: Senjie Lin, University of Connecticut, United States of America

Received: November 4, 2013; **Accepted:** March 12, 2014; **Published:** April 7, 2014

Copyright: © 2014 Hu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The authors acknowledge the financial support from the China Scholarship Council to Zhangxi Hu. In addition, this work was supported by NOAA and NSF grants to MRM, and Natural Science Foundation of China (NSFC) (Grant Numbers 40776078, 40876074, 41176104). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: txuning@163.com (NX); mmulholl@odu.edu (MRM)

Introduction

Harmful algal blooms (HABs) have increased in coastal waters worldwide [1–4], exerting serious economic impacts on marine fisheries and aquaculture, and threatening public health and aquatic ecosystems [1,5]. In Chinese coastal waters, HABs have increased in frequency, intensity, and duration in recent decades [6–8]. *Prorocentrum donghaiense* is a HAB species that frequently blooms during late spring and early summer in coastal waters around the East China Sea, including the Changjiang River Estuary, and coastal waters adjacent to Zhejiang in China [9–11], and coastal areas near Japan and South Korea [9]. Between 2000 and 2006, about 120 *P. donghaiense* bloom events (maximum cell densities of up to $3.6\times 10^8\ \text{cells}\cdot\text{L}^{-1}$) were reported from Chinese coastal waters (data from State Oceanic Administration, China) [2] and the spatial extent of these blooms ranged from several thousand to more than ten thousand km^2 . Blooms persisted several days to a month, during which time they caused serious economic impacts to marine fisheries, public health, and aquatic ecosystems [8,10,11]. Despite documentation regarding their impacts, the environmental factors promoting *P. donghaiense* bloom initiation and persistence are still unclear.

HABs are often linked to nutrient over-enrichment and consequent eutrophication of coastal waters including the East

China Sea [5,12–14]. The Changjiang River Estuary and East China Sea, where *P. donghaiense* blooms frequently occur, are hydrologically complex; they receive terrestrial nutrient inputs and freshwater through the watershed, but are also influenced by oceanic circulation that can affect particle transport in the estuary [15]. High nutrient loads to the watershed have been implicated as the proximate cause of *P. donghaiense* blooms in this system [16].

Dissolved inorganic N concentrations in the Changjiang River Estuary, mainly in the form of NO_3^- , increased several-fold in the past few decades, reaching as high as $131.6\ \mu\text{mol N L}^{-1}$ [15–18]. However, field studies have shown that NO_3^- is rapidly consumed during the initiation of *P. donghaiense* blooms leading to NO_3^- -depletion during blooms [16]. Our previous culture study determined that *P. donghaiense* can grow on multiple forms of N when they are supplied in saturating concentrations ($50\ \mu\text{mol N L}^{-1}$) [19]. However, it is not known whether *P. donghaiense* has a strong affinity for NO_3^- or sources of recycled N (e.g., NH_4^+ , urea, and amino acids) when they are present at lower concentrations, more realistic of environmental concentrations once blooms are established.

The dissolved N pool in aquatic systems is dynamic, consisting of varying concentrations of dissolved inorganic N (DIN) and dissolved organic N (DON) compounds, many of which remain to be identified [20–22]. In Chinese coastal waters DIN ($\text{NO}_3^- + \text{N-}$

$\text{O}_2^- + \text{NH}_4^+$ concentrations ranged 0–45.0 $\mu\text{mol N L}^{-1}$ in the surface waters of Southern Yellow Sea during 2006–2007 [23] and 0–60.0 $\mu\text{mol N L}^{-1}$ in waters adjacent to Hong Kong [24]. In the Changjiang River estuary and East China Sea, DIN (NO_3^- , NO_2^- and NH_4^+) concentrations ranged 0–131.6 $\mu\text{mol N L}^{-1}$ [15–18] over the same time period. In addition to DIN, urea can be an important component of the dissolved N pool and accounted for ~10–60% of the total dissolved N in Hong Kong waters in summer of 2008 [24]. In the East China Sea, DON concentrations ranged 5.78–25.26 $\mu\text{mol N L}^{-1}$, accounting for 44.0–88.5% of total dissolved nitrogen [16]. While concentrations of urea and DFAA were only a small component of the DON pool, 0.21–1.39 and 0.09–1.55 $\mu\text{mol N L}^{-1}$, respectively, these compounds are labile and rapidly regenerated and turned over in the environment [21,22].

To date, our understanding of the nitrogenous nutrition of *P. donghaiense* is limited. While previous research demonstrated that *P. donghaiense* can potentially utilize multiple inorganic and organic N compounds to grow [19], little is known about their affinity and capacity for growth on the diverse range of N compounds found in the environment. Therefore, we compared growth kinetics of *P. donghaiense* on medium supplied with NO_3^- , NH_4^+ , urea, glu, dialanine (diala) or cyanate as the sole source of N. In addition, we used stable isotopes as tracers to compare uptake kinetics for these compounds in cultures acclimated under NO_3^- -replete and -deplete culture conditions, conditions likely present in the environment during bloom initiation and maintenance phases, respectively.

Materials and Methods

2.1 Culture conditions

Prorocentrum donghaiense, was isolated from the East China Sea, and obtained from the Research Center of Hydrobiology, Jinan University, Guangzhou, China. Cultures were grown in artificial seawater enriched with, sterile, silicate-free f/2 trace metals and vitamins [25]. Concentrations of NO_3^- were adjusted to conform to different experimental treatments (see below). Cultures were maintained in an environmental room at constant temperature ($23 \pm 0.5^\circ\text{C}$) and irradiance ($60 \pm 5 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$) on a 12 h:12 h (light: dark) light cycle and transferred to fresh medium every week to ensure cells remained in exponential growth phase for the duration of the experiments.

For all experiments, dissolved nutrients were measured after filtering through a 0.2 μm Supor filter. Samples were stored frozen until analysis. NO_3^- and urea were measured on an Astoria Pacific autoanalyzer according to the manufacturer's specifications and using colorimetric methods [26]. NH_4^+ was determined using the manual phenol hypochlorite method [27]. Concentrations of dissolved free amino acids (DFAA) were measured by high performance liquid chromatography (HPLC) [28]. Cells were enumerated using a FACSCalibur flow cytometer (Becton Dickinson Instruments). Samples were also collected onto GF/F filters and frozen for analysis of chlorophyll *a* (Chl *a*). Chl *a* concentrations were measured fluorometrically within one week of sample collection after extraction of cells in 90% acetone [29].

2.2 Experimental design

2.2.1 Growth kinetics experiments. Growth kinetics of *P. donghaiense* were determined for six different N compounds. Cultures were grown in triplicate, capped, 50 ml Pyrex test tubes containing 35 ml of culture suspended in f/2 modified medium amended with 0.5, 2, 5, 20, 50, 100, 200, or 500 $\mu\text{mol N L}^{-1}$ supplied in the form of NO_3^- , NH_4^+ , urea, glu, diala, or cyanate.

Cultures were maintained in the environmental chamber under conditions described above for at least three generations prior to experiments in order to ensure cells were acclimated to treatment conditions. *In vivo* fluorescence was monitored to estimate chlorophyll biomass using a Turner Designs AU-10 fluorometer at the same time each day. Growth rates, μ (day^{-1}), were calculated using a least squares fit to a straight line after logarithmic transformation of *in vivo* fluorescence data as described by Guillard [30] using the equation:

$$\mu = \frac{\ln(N_1) - \ln(N_0)}{T_1 - T_0} \quad (1)$$

where N_1 and N_0 were the biomass (*in vivo* fluorescence) at time T_1 and T_0 , respectively, during the linear portion of exponential phase growth.

The relationship between specific growth rate and N concentration was fitted to a Monod growth kinetic model using the equation:

$$\mu = \frac{\mu_m \cdot S}{K_\mu + S} \quad (2)$$

where μ was the specific growth rate (day^{-1}) calculated during the linear portion of exponential phase growth (Equation 1), μ_m was the maximum specific growth rate (day^{-1}), S was the N concentration ($\mu\text{mol N L}^{-1}$), and K_μ (half saturation constant; $\mu\text{mol N L}^{-1}$) was the N concentration at $\mu_m/2$ [31].

2.2.2 N uptake kinetic experiments. For N uptake kinetic experiments, cultures of *P. donghaiense* were grown on f/2 medium modified to contain 50 $\mu\text{mol N L}^{-1}$ as NO_3^- in two, 5-liter bottles. Cell abundance and ambient NO_3^- concentrations were monitored daily. N-deplete uptake kinetic experiments were initiated after NO_3^- concentrations were below the detection limit (0.05 $\mu\text{mol N L}^{-1}$) for 5 consecutive days. Uptake kinetics experiments were initiated by dispensing 35 ml of NO_3^- -deplete culture into 50 ml Pyrex test tubes and adding $^{15}\text{NO}_3^-$, $^{15}\text{NH}_4^+$, ^{15}N - ^{13}C dually labeled urea, glu, or an algal amino acid mixture to duplicate test tubes at final substrate concentrations of 0.1, 0.2, 0.5, 1.0, 2.0, 5.0, 10.0, 20.0 and 50.0 $\mu\text{mol N L}^{-1}$. For NO_3^- -replete uptake kinetic experiments, 35 ml of culture was dispensed into a series of 50 ml Pyrex test tubes and each was amended with an additional 50.0 $\mu\text{mol N L}^{-1}$ as $^{14}\text{NO}_3^-$ to ensure the cultures were NO_3^- -replete. As for NO_3^- -replete treatments, $^{15}\text{NH}_4^+$, and ^{15}N - ^{13}C dually labeled urea, glu, or an algal amino acid mixture were added to duplicate tubes at the same concentrations indicated above, however, NO_3^- uptake kinetics could not be measured in these NO_3^- -replete cultures.

Uptake experiments were incubated at $\sim 23^\circ\text{C}$ and at an irradiance of $\sim 60 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$. After <1 hour, incubations were terminated by gentle filtration through pre-combusted (450°C for 2 hours) GF/F (nominal pore size $\sim 0.7 \mu\text{m}$) filters. Filters were immediately frozen in sterile polypropylene cryovials until analysis. Frozen samples were dried in an oven at 40°C for 2 days and pelletized into tin discs prior to analysis on a Europa Scientific 20–20 isotope ratio mass spectrometer equipped with an automated N and C analyzer preparation unit. Specific uptake rates were calculated using a mixing model [32]:

$$^{15}\text{N uptake}(V) = \frac{(\text{atom}\% \text{PN})_{\text{final}} - (\text{atom}\% \text{PN})_{\text{initial}}}{(\text{atom}\% \text{N source pool} - \text{atom}\% \text{PN})_{\text{initial}} \times \text{time}} \quad (3)$$

where the source pool was the dissolved N pool that was enriched during uptake experiments. Specific uptake rates (V ; h^{-1}) at each N concentration for each N substrate tested were fitted to a Michaelis-Menten model using the following equation:

$$V = \frac{V_m \cdot S}{K_s + S} \quad (4)$$

where V_m is the maximum specific uptake rate (h^{-1}); S the substrate concentration ($\mu\text{mol N L}^{-1}$); and K_s the half saturation constant, the concentration ($\mu\text{mol N L}^{-1}$) where V is equivalent to half of V_m , for the N substrate tested.

2.3 Statistical analysis

All statistical tests were carried out using Origin 7.0 and Microsoft Excel 2007 with the level of significance set at an α equal to 0.05. Differences between growth and uptake kinetic parameters were compared using a one-way ANOVA test. Monod and Michaelis-Menten curves were fitted using Origin 7.0.

Results

3.1 Growth kinetics of *P. donghaiense*

P. donghaiense exhibited standard Monod-type growth kinetics over N concentrations ranging from 0.5–500 $\mu\text{mol N L}^{-1}$ for NO_3^- and NH_4^+ , 0.5–50 $\mu\text{mol N L}^{-1}$ for urea, 0.5–100 $\mu\text{mol N L}^{-1}$ for glu and cyanate, and 0.5–200 $\mu\text{mol N L}^{-1}$ for diala (Fig. 1). Maximum specific growth rates (μ_m) ranged from 0.71 ± 0.04 to $1.51 \pm 0.06 \text{ d}^{-1}$ (Table 1). The highest maximum growth rates were observed in cultures growing with glu or urea as the sole N source; maximum growth rates were a factor of 2 lower in cultures grown with NO_3^- , NH_4^+ , cyanate, or diala supplied as the sole N source (Table 1). Half saturation constants (K_μ) ranged from 0.28 ± 0.08 to $5.25 \pm 0.40 \mu\text{mol N L}^{-1}$. The lowest K_μ concentrations were observed in *P. donghaiense* cultures growing on media containing NO_3^- , NH_4^+ , and cyanate as the sole source of N (0.28 – $0.51 \mu\text{mol N L}^{-1}$) (Table 1). In contrast, K_μ concentrations were an order of magnitude higher in cultures grown with urea, glu, or diala as the sole source of N suggesting higher affinities for cyanate, NO_3^- , NH_4^+ , and urea than for glu and diala (Table 1).

3.2 N uptake kinetics by *P. donghaiense*

Kinetics for NH_4^+ , urea, and algal amino acids conformed to Michaelis-Menten kinetics in both NO_3^- -deplete and -replete cultures of *P. donghaiense* (Table 2; Figs. 2 and 3). While NO_3^- uptake also conformed to Michaelis-Menten kinetics in NO_3^- -deplete cultures (Fig. 2a), NO_3^- uptake kinetics could not be measured in NO_3^- -replete cultures because of the excess NO_3^- in the growth media (Fig. 3a). Glu uptake did not conform to the Michaelis-Menten kinetic model in either NO_3^- -deplete (Fig. 2d) or -replete (Fig. 3d) cultures of *P. donghaiense* and higher glu uptake rates were observed at lower glu concentrations.

Uptake kinetics for NO_3^- , NH_4^+ , urea, and algal amino acids followed Michaelis-Menten type uptake kinetics in NO_3^- -deplete batch cultures of *P. donghaiense* (Table 2; Fig. 2). Although results were sufficient to fit a Michaelis-Menten model for NH_4^+ uptake, NH_4^+ uptake by NO_3^- -deplete *P. donghaiense* was not saturated even at the highest concentrations used in this experiment (Fig. 2b). Maximum specific uptake rates (V_m) of $(66.9 \pm 6.0) \times 10^{-3} \text{ h}^{-1}$ were calculated for NH_4^+ (Table 2), while V_m calculated for NO_3^- ($(34.4 \pm 0.9) \times 10^{-3} \text{ h}^{-1}$), urea ($(44.7 \pm 0.4) \times 10^{-3} \text{ h}^{-1}$), and the algal amino acid mixture ($(32.6 \pm 1.0) \times 10^{-3} \text{ h}^{-1}$) were lower (Table 2). The half saturation constants (K_s) for uptake of NO_3^- ,

NH_4^+ , urea, and algal amino acids by NO_3^- -deplete *P. donghaiense* were 1.3 ± 0.1 , 5.3 ± 1.1 , 0.13 ± 0.01 and $9.9 \pm 0.9 \mu\text{mol N L}^{-1}$, respectively (Table 2). The α values, which are often used as an index of nutrient affinity, were 27.7 ± 2.4 , 12.8 ± 1.6 , 339.4 ± 26.3 and 3.3 ± 0.2 for NO_3^- , NH_4^+ , urea, and algal amino acids, respectively (Table 2) suggesting NO_3^- -deplete *P. donghaiense* had the highest affinity for urea and the lowest for amino acids.

In NO_3^- -replete cultures of *P. donghaiense*, maximum specific uptake rates (V_m) of NH_4^+ were greatest and not significantly different than those observed in NO_3^- -deplete cultures ($p > 0.05$; Table 2) (Table 2). Similarly, V_m for urea and algal amino acids were lower than those observed for NH_4^+ but comparable to those observed in NO_3^- -deplete cultures (Table 2; Figs. 2 and 3). For urea, the V_m was significantly higher ($p < 0.05$) but for algal amino acids there was no significant difference ($p > 0.05$) between NO_3^- -replete and NO_3^- -deplete cultures of *P. donghaiense* (Table 2). Although there were significant differences between K_s values for NH_4^+ and algal amino acid uptake ($p < 0.05$) between NO_3^- -replete and -deplete cultures of *P. donghaiense*, the K_s values for NH_4^+ , urea, and the algal amino acid uptake in NO_3^- -replete cultures of *P. donghaiense* were 7.1 ± 0.4 , 0.1 ± 0.01 and $12.5 \pm 0.1 \mu\text{mol N L}^{-1} \text{ h}^{-1}$, respectively (Table 2), similar to and showing the same pattern as values measured in NO_3^- -deplete cultures. Similarly, the derived α values, a measure of nutrient affinity, were also similar for NO_3^- -replete and -deplete cultures of *P. donghaiense* even though there were significant differences ($p < 0.05$) in α between NO_3^- -replete and -deplete cultures for algal amino acids (Table 2).

Discussion

Nutrient enrichment has been implicated as a causal factor in the occurrence of HAB events worldwide [5,14,33]. Both the form of dissolved N and its concentration in coastal and marine environments are thought to be important to the formation and development of HABs and phytoplankton blooms in general [13,15,34,35]. However, different phytoplankton species and groups have different uptake capacities and affinities for specific inorganic and organic nitrogen compounds [19,20,36]. Many phytoplankton can also mobilize N from complex organic compounds such as peptides using extracellular enzymes thereby making N from these compounds available for uptake [32,37]. In coastal areas, the concentration and composition of the dissolved N pool and the relative concentrations of various N compounds vary temporally and spatially, likely impacting the phytoplankton community composition and abundance on both short and longer timescales [15,38,39].

P. donghaiense is a mixotrophic species capable of both autotrophic and heterotrophic metabolisms. Previous research demonstrated that *P. donghaiense* can utilize inorganic and organic N compounds including urea, amino acids, small peptides, and cyanate [16,19] as N sources. In addition, this organism can also ingest cyanobacteria, cryptophytes, and other dinoflagellates [40]. To better understand the N nutrition and capabilities of *P. donghaiense*, we conducted growth and N uptake kinetic experiments in cultured isolates to determine their affinity, preference, and capacity for growth on different N compounds and found that *P. donghaiense* had high maximum growth rates and exhibited high uptake capacities and affinities for a diverse suite of N compounds under both NO_3^- -deplete and -replete conditions.

4.1 Growth kinetics

The Monod equation [41] has been used to describe saturation kinetics for nutrient-limited phytoplankton growth in many field




Figure 1. Growth kinetics for *P. donghaiense*. Growth rate as a function of NO₃⁻ (a), NH₄⁺ (b), urea (c), glu (d), cyanate (e) and diala (f) concentrations in batch cultures of *P. donghaiense*. Solid lines were fitted iteratively to the data according to the Monod equation and kinetic parameters were calculated as described in the text.
doi:10.1371/journal.pone.0094030.g001

and laboratory studies [42–44]. Maximum specific growth rates (μ_m) and half saturation constants (K_μ) are two important kinetic parameters derived from this model that quantify organismal growth responses to environmental nutrient concentrations. The initial slope of the Monod function (μ_m/K_μ) is thought to be a competitive index and used to assess the affinity of phytoplankton for particular nutrient substrates [45]. This model provides several useful tools for evaluating fitness of co-occurring phytoplankton with respect to the external nutrient availability in the environment [46]. Theoretical studies of resource competition suggest that

the functional relationship between growth rate and the concentration of nutrient elements in the environment determines a species' ability to compete for that nutrient in the environment [46].

In our culture experiments, we found *P. donghaiense* grew well on DIN (NO₃⁻ and NH₄⁺) and DON (urea, glu, diala and cyanate) compounds when they were supplied as the sole source of N (Fig. 1). *P. donghaiense* had comparable maximum specific growth rates when growing on NO₃⁻, NH₄⁺, cyanate, or diala (0.71–0.83 d⁻¹); but maximum specific growth rates were a factor of 2

Table 1. Mean (\pm SD, n=3) growth kinetic parameters for *P. donghaiense* growing with NO_3^- , NH_4^+ , urea, glu, cyanate and diala as the sole source of N in the culture medium.

| Nitrogen species | μ_m (d^{-1}) | K_μ ($\mu\text{mol N L}^{-1}$) | Affinity (α) (μ_m/K_μ) | R^2 |
|------------------|-----------------------------|--------------------------------------|---|-------|
| NO_3^- | 0.74 ± 0.02 | 0.42 ± 0.02 | 1.74 ± 0.13 | 0.82 |
| NH_4^+ | 0.83 ± 0.02 | 0.51 ± 0.03 | 1.65 ± 0.10 | 0.92 |
| Urea | 1.50 ± 0.05 | 2.05 ± 0.04 | 0.73 ± 0.02 | 0.91 |
| Glu | 1.51 ± 0.06 | 5.25 ± 0.40 | 0.29 ± 0.01 | 0.98 |
| cyanate | 0.71 ± 0.04 | 0.28 ± 0.08 | 2.58 ± 0.49 | 0.85 |
| Diala | 0.82 ± 0.01 | 2.05 ± 0.02 | 0.40 ± 0.01 | 0.76 |

The maximum specific growth rate (μ_m), half saturation constant (K_μ) and affinity (α) for each nitrogen compound was calculated according to the Monod model.

doi:10.1371/journal.pone.0094030.t001

higher in cultures supplied with urea or glu as the sole source of N (1.50 and 1.51 d^{-1} , respectively; Table 1). In contrast, half saturation constants for growth (K_μ) were more variable (0.28 ± 0.08 to $5.25 \pm 0.40 \mu\text{mol N L}^{-1}$); K_μ values were lower (0.28 – $0.51 \mu\text{mol N L}^{-1}$) for cultures growing on NO_3^- , NH_4^+ , or cyanate, and were an order of magnitude higher (2.05 – $5.25 \mu\text{mol N L}^{-1}$) in cultures grown on urea, glu, or diala (Table 1), suggesting a higher affinity for the inorganic N compounds. So, while *P. donghaiense* had higher maximum growth rates when supplied urea or glu as a sole source of N, cells had a higher affinity for cyanate, NO_3^- , and NH_4^+ relative to urea, glu, and diala (Table 1) suggesting that the former compounds might be important sources N when nutrient concentrations are submicromolar.

Calculated growth kinetic parameters for *P. donghaiense* indicate that maximum specific growth rates and half saturation constants (K_μ) for NO_3^- , NH_4^+ , and urea, were comparable to each other and those measured for other bloom-forming dinoflagellate species, *Prorocentrum minimum*, *Cochlodinium polykrikoides* and for *Scrippsiella trochoidea* growing on urea (Table 3). Maximum growth rates for *P. donghaiense* growing with NO_3^- , NH_4^+ , urea or glu as the sole source of N were higher than those previously measured for *P. minimum*, *C. polykrikoides*, and *S. trochoidea* growing on the same N compounds (Table 3). *S. trochoidea*, *P. minimum* and *C. polykrikoides*

had comparable maximum specific growth rates when going on urea (0.44 – 0.47 d^{-1} ; Table 3), but these were all lower than the maximum specific growth rate observed for *P. donghaiense* in this study (1.50 d^{-1}) (Table 1). While *P. donghaiense* (this study) and *C. polykrikoides* [44] exhibited comparable maximum specific growth rates on NO_3^- , NH_4^+ , urea and glu, *P. minimum* exhibited higher specific growth rates in cultures growing on urea or NO_3^- (0.45 and $0.43 \pm 0.01 \text{ d}^{-1}$, respectively) relative to cultures growing on NH_4^+ or glu (0.27 ± 0.05 and $0.23 \pm 0.02 \text{ d}^{-1}$, respectively) [43].

But unlike *P. donghaiense*, *P. minimum* had the lowest half saturation constants for growth on urea, followed by NO_3^- and NH_4^+ ; both had the highest half saturation constants for glu (Table 3) [43]. In contrast, *C. polykrikoides* had higher half saturation constants for urea and NH_4^+ than for NO_3^- and glu [44]. The half saturation constant for *S. trochoidea* growing on urea was much higher than for the other three dinoflagellates suggesting a low affinity for growth on this compound [47].

In this study, *P. donghaiense* also exhibited maximum growth rates and half saturation constants comparable to those for growth on NH_4^+ and NO_3^- , when growing with cyanate and dialanine as the sole source of N (Fig. 1; Table 1). Because peptides and cyanate are degradation products of decaying cells [32,48], these compounds were likely available to *P. donghaiense* during bloom maintenance when NO_3^- concentrations are exhausted and recycling processes are important for maintaining cell biomass and turnover. The concentration and turnover of these organic compounds have not been assessed in natural systems where blooms of *P. donghaiense* occur. Comparisons of *P. donghaiense* growth kinetics with those of co-occurring species relative to the bioavailable DIN and DON compounds in the environment is warranted to elucidate how competitive interactions among phytoplankton populations contributes to bloom initiation and maintenance.

4.2 Uptake kinetics

The Michaelis-Menten equation is often used to describe the relationship between external nutrient concentrations and their uptake rates [49–51]. As for growth kinetics, kinetic parameters for N uptake have been used to assess the relative preference (and affinity) for different N substrates in the environment [50,52]. While kinetic parameters are usually measured in nutrient-deplete cultures, we conducted N uptake kinetic experiments in both NO_3^- -deplete and -replete batch cultures of *P. donghaiense* because NO_3^- concentrations are generally high in the Changjiang River

Table 2. Mean (\pm SD, n=2) uptake kinetic parameters (maximum specific uptake rate (V_m), half saturation constant (K_s) and affinity (α)) for *P. donghaiense* growing on NO_3^- deplete culture media.

| Nitrogen species | V_m ($\times 10^{-3} \text{ h}^{-1}$) | K_s ($\mu\text{mol N L}^{-1}$) | Affinity (α) (V_m/K_s) | R^2 |
|--|---|------------------------------------|-------------------------------------|-------|
| NO_3^- deplete cultures: | | | | |
| NO_3^- | 34.4 ± 0.9 | 1.3 ± 0.1 | 27.7 ± 2.4 | 0.97 |
| NH_4^+ | 66.9 ± 6.0 | 5.3 ± 1.1 | 12.8 ± 1.6 | 0.99 |
| Urea | 44.7 ± 0.4 | 0.13 ± 0.01 | 339.4 ± 26.3 | 0.61 |
| algal amino acids | 32.6 ± 1.0 | 9.9 ± 0.9 | 3.3 ± 0.2 | 0.93 |
| NO_3^- replete cultures ($50 \mu\text{mol L}^{-1} \text{NO}_3^-$): | | | | |
| NH_4^+ | 74.7 ± 3.4 | 7.1 ± 0.4 | 10.5 ± 0.1 | 0.99 |
| Urea | 39.7 ± 0.2 | 0.12 ± 0.01 | 332.9 ± 26.7 | 0.49 |
| algal amino acids | 31.4 ± 0.4 | 12.5 ± 0.1 | 2.5 ± 0.0 | 0.94 |

Results from glu uptake kinetic studies did not conform to Michaelis-Menten kinetics so kinetic parameters (Figs. 2d and 3d) so are not included here.

doi:10.1371/journal.pone.0094030.t002




Figure 2. N uptake kinetics for *P. donghaiense* growing in NO₃⁻-deplete batch cultures. Nitrogen uptake as a function of NO₃⁻ (a), NH₄⁺ (b), urea (c), glu (d), and algal amino acids (e) concentration in batch cultures of *P. donghaiense* growing on NO₃⁻-deplete media. Solid lines were fitted iteratively to the data according to the Michaelis-Menten equation and kinetic parameters were calculated as described in the text. doi:10.1371/journal.pone.0094030.g002

estuary when blooms of this organism initiate [15–18] but then become rapidly depleted as algal biomass increases. Here we compared N uptake kinetics for NO₃⁻ (N-deplete cultures only), NH₄⁺, urea, and algal amino acids in N-replete and -deplete cultures of *P. donghaiense* (Table 2; Figs. 2, 3).

Michaelis-Menten type uptake kinetics were observed for all N compounds tested except glu (Figs. 2d, 3d). For NH₄⁺, the uptake capacity (V_m) for exponentially growing *P. donghaiense* was ~1.5, 1.9, and 2.1 times greater than those for urea, NO₃⁻, and the algal amino acids, respectively, in both NO₃⁻-deplete and -replete cultures. The half saturation constants, which are also used to estimate nutrient affinity under nutrient-limiting conditions, suggest similar nutrient preferences under NO₃⁻-deplete and -replete conditions; urea >>>NO₃⁻>NH₄⁺> algal amino acids for NO₃⁻-deplete *P. donghaiense*, and urea >>>NH₄⁺> algal amino acids for NO₃⁻-replete *P. donghaiense* (Table 2; Figs. 2, 3).

Differences in kinetic parameters are often observed in cells growing under N-replete versus -deplete conditions [53]. Many

N-starved cells are capable of enhanced N uptake when nutrients are resupplied, and N-replete cells can have higher half-saturation constants than those grown under N-depleted conditions [36]. While our kinetics parameters conformed to this norm based on our statistical analyses, it was surprising that uptake kinetics for NH₄⁺, urea, and amino acids were similar in nutrient-replete and -deplete cultures of *P. donghaiense* (Table 2; Figs. 2&3). Ambient NO₃⁻ concentrations were ~43 μmol N L⁻¹ when we initiated our uptake experiments in the N-replete cultures and this should have been saturating for NO₃⁻ uptake (Table 2). However, kinetic parameters in NO₃⁻-replete and -deplete batch cultures were similar suggesting that even when NO₃⁻ concentrations are high, such as during bloom initiation, *P. donghaiense* has a high affinity for other N compounds that may available in the environment and this may give them a competitive edge over other phytoplankton.

NO₃⁻ and NH₄⁺ are the dominant DIN species thought to fuel the bulk of primary production in estuarine, coastal, and oceanic environments. In Chinese estuaries and coastal waters where *P.*




Figure 3. N uptake kinetics for *P. donghaiense* growing in NO₃⁻-replete batch cultures. Nitrogen uptake rates as a function of NO₃⁻ (a), NH₄⁺ (b), urea (c), glu (d), and algal amino acid (e) concentrations in batch cultures of *P. donghaiense* growing on NO₃⁻-replete media. Solid lines were fitted iteratively to the data according to the Michaelis-Menten equation and kinetic parameters were calculated as described in the text. doi:10.1371/journal.pone.0094030.g003

Table 3. Summary of mean (±SD) maximum specific growth rates (μ_{max} , d⁻¹) and half saturation constants (k_{μ} , μmol N L⁻¹) for nitrate (NO₃⁻), ammonium (NH₄⁺), urea, and glutamic acid (glu) estimated for different dinoflagellate species.

| Species | NO ₃ ⁻ | | NH ₄ ⁺ | | Urea | | glu | | Reference |
|-----------------------------------|------------------------------|-----------|------------------------------|-----------|-------------|-----------|-------------|-----------|------------|
| | μ_{max} | k_{μ} | μ_{max} | k_{μ} | μ_{max} | k_{μ} | μ_{max} | k_{μ} | |
| <i>Prorocentrum minimum</i> | 0.43±0.01 | 1.10±0.40 | 0.23±0.02 | 0.70±0.40 | 0.45 | <0.50 | 0.27±0.05 | 53.0±31.0 | [43] |
| <i>Cochlodinium polykrikoides</i> | 0.43±0.01 | 2.06±0.32 | 0.44±0.02 | 2.69±0.49 | 0.44±0.01 | 2.90±0.46 | 0.53±0.02 | 1.91±0.47 | [44] |
| <i>Scrippsiella trochoidea</i> | - | - | - | - | 0.47 | 21.83 | - | - | [47] |
| <i>P. donghaiense</i> | 0.74±0.02 | 0.42±0.02 | 0.83±0.02 | 0.51±0.03 | 1.50±0.05 | 2.05±0.04 | 1.51±0.06 | 5.25±0.40 | This study |

doi:10.1371/journal.pone.0094030.t003

Table 4. Summary of mean (\pm SD) maximum uptake rates (V_{\max} , $\times 10^{-3} \text{ h}^{-1}$) and half saturation constants (k_s , $\mu\text{mol N L}^{-1}$) for ammonium (NH_4^+), nitrate (NO_3^-), urea and amino acid (glutamic acid (glu), glutamate (gln), glycine (gly) or amino acid mixture) uptake by HAB dinoflagellate species.

| Species | Experiment type | NO_3^- | | NH_4^+ | | urea | | amino acids | | Reference |
|--|------------------------------------|---------------------------------|-----------------------------------|---------------------------------|---------------------------------|-----------------------------------|---------------------------------|---------------------------------------|-------------------------------|-------------------------|
| | | V_{\max} | k_s | V_{\max} | k_s | V_{\max} | k_s | V_{\max} | k_s | |
| <i>Heterosigma akashiwo</i> | Culture | 18.0 \pm 1.08 | 1.47 \pm 0.25 | 28.0 \pm 2.17 | 1.44 \pm 0.35 | 2.89 \pm 0.24 | 0.42 \pm 0.16 | - | - | [52] |
| <i>Pymnesium parvum</i> | Culture | - | - | - | - | - | - | 10.5–13.1 | 0.09–0.10 | [62] ^b |
| <i>Prorocentrum minimum</i> | Culture | - | - | 46–65 | 1.2–4.6 | 3E-01–8E-01 | 0.05–5.4 | 4–7 | 0.48–3.54 | [63] ^c |
| <i>Prorocentrum minimum</i> | Culture | 95.8 \pm 5.5–341.2 \pm 90.7 | 0.68 \pm 0.23–23.3.0 \pm 11.1 | 1070 \pm 120.5– \pm 88.7 | 2.48 \pm 0.8–6.23 \pm 0.47 | 149.2 \pm 12.3–480.5 \pm 13.4 | 0.86 \pm 0.16–1.82 \pm 0.59 | 271.6 \pm 35.6 517.1 \pm 125.8 | 14.1 \pm 6.8–22.6 \pm 9.9 | [55] ^{a, d} |
| <i>Prorocentrum minimum</i> (Choptank River) | Field | 18.4 \pm 4.3–53.8 \pm 5.7 | 1.4 \pm 0.2–7.1 \pm 5.5 | 327 \pm 172.5–868.6 \pm 159 | 2.4 \pm 0.3–9.8 \pm 2.6 | 38.2 \pm 16.9–492.6 \pm 71.5 | 6.6 \pm 4.0–17.9 \pm 5.1 | 84.2 \pm 12.9–1516 \pm 134 | 4.8 \pm 2.3–26.6 \pm 4.1 | [55] ^{a, d} |
| <i>Prorocentrum donghaiense</i> | Culture (NO_3^- deplete) | 34.4 \pm 0.9 | 1.3 \pm 0.1 | 66.9 \pm 6.0 | 5.3 \pm 1.1 | 44.7 \pm 0.4 | 0.13 \pm 0.01 | 32.6 \pm 1.0 | 9.9 \pm 0.9 | This study ^a |
| <i>Prorocentrum donghaiense</i> | Culture (NO_3^- replete) | - | - | 74.7 \pm 3.4 | 7.1 \pm 0.4 | 39.7 \pm 0.2 | 0.12 \pm 0.01 | 31.4 \pm 0.4 | 12.5 \pm 0.1 | This study ^a |
| <i>Prorocentrum donghaiense</i> dominant | Field | - | - | 700 | 3.9 | 17.0 | 5.3 | 25 | 1.8 | [16] ^c |
| <i>Karenia mikimotoi</i> | Field | 18.0 | 43.6 | 500 | 4.9 | 20.0 | 1.4 | 11 | 1.7 | [16] ^c |
| <i>Alexandrium minutum</i> | Culture | 0.29 \pm 0.01–0.70 | 0.22 \pm 0.02–0.28 \pm 0.06 | 0.65 \pm 0.01–1.49 | 0.25 \pm 0.04–0.38 \pm 0.04 | - | - | - | - | [58] ^e |
| <i>Alexandrium catenella</i> | Culture | 3–47 | 0.6–28.1 | 26.0 \pm 2.0 | 2.0 \pm 0.6 | 25 \pm 8 | 28.4 \pm 15.0 | - | - | [66] |
| <i>Alexandrium catenella</i> | Field | - | - | 14.9 \pm 0.8 | 2.52 \pm 0.36 | 3.5 \pm 0.2 | 0.65 \pm 0.12 | - | - | [39] |
| <i>Alexandrium catenella</i> | Field | 24 \pm 2 | 4.6 \pm 1.1 | 64 \pm 5 | 8.4 \pm 1.7 | 61 \pm 8 | 43.9 \pm 8.8 | - | - | [66] |
| <i>Dinophysis acuminata</i> | Field | 3.5 \pm 0.2 | 0.79 \pm 0.26 | 13.9 \pm 0.2 | 0.67 \pm 0.06 | 6.2 \pm 0.6 | 0.53 \pm 0.22 | - | - | [39] |
| <i>Lingulodinium polyedrum</i> | Field | 22.4 | 0.47 | 47.1 | 0.59 | 61.6 | 0.99 | - | - | [65] |
| <i>Cochlodinium</i> spp. | Field | 0.9 \pm 0.0 | 1.0 \pm 0.4 | - | - | 1.9 \pm 0.1–2.2 \pm 0.3 | 1.6 \pm 0.2–6.6 \pm 2.0 | - | - | [67] |
| Mixed dinoflagellate assemblages (Neuse Estuary) | Field | 4.0 \pm 1.6 | 0.5 \pm 0.1 | 52.9 \pm 1.7 | 4.9 \pm 0.5 | 5.8 \pm 0.5 | 0.4 \pm 0.1 | 2.3 \pm 0.5 | 2.3 \pm 1.7 | [55] ^{h, d} |

The ^a, ^b and ^c indicate that an algal amino acid mixture, glutamic acid (glu) or glycine (gly) used as N substrates, respectively. The ^d and ^e indicate the units are $\text{fmol N cell}^{-1} \text{ h}^{-1}$ and $\text{pmol N cell}^{-1} \text{ h}^{-1}$, respectively. doi:10.1371/journal.pone.0094030.t004

donghaiense blooms, NO_3^- and NH_4^+ concentrations are commonly 0–20.0 and 0–3.0 $\mu\text{mol N L}^{-1}$, respectively [16]. In this study, we found *P. donghaiense* had higher maximum uptake rates for NH_4^+ than NO_3^- , urea and amino acids in both NO_3^- -deplete and -replete batch cultures (Table 2; Fig. 2). However, because of the low half saturation constants for urea uptake (0.13 and 0.12, respectively), both N-deplete and -replete cultures of *P. donghaiense* had a much higher affinity for urea than for inorganic N or amino acids. This is consistent with results from a field study in the East China Sea during a *P. donghaiense* dominant bloom [11,16]. As we know, DON can constitute more than 80% of the TDN in surface waters in coastal systems and the open ocean [21] where it can also contribute to primary productivity [20]. In the East China Sea, DON accounted for 44.0%–88.5% of the TDN, and urea and DFAA accounted for 1.7%–36.3% of the DON, before and during a *P. donghaiense* bloom [16]. In our experiments, *P. donghaiense* had similar maximum specific uptake rates for urea and amino acids under NO_3^- -deplete and -replete conditions (Table 2) and half saturation constants for urea uptake were much lower than for NO_3^- , NH_4^+ and amino acids, suggesting a higher affinity for urea than for other N compounds tested. Further, during a bloom dominated by *P. donghaiense* in the East China Sea in 2005, maximum specific uptake rates for urea were $11\text{--}28 \times 10^{-3} \text{ h}^{-1}$ and half saturation constants were $0.25\text{--}14.95 \mu\text{mol N L}^{-1}$, which are comparable to results presented here for culture studies [16]. These results suggest that even if present at low concentrations, urea has the potential to be an important source of N fueling the growth of *P. donghaiense* and this may be important for out-competing co-occurring phytoplankton during bloom initiation and/or supporting the persistence of bloom organisms in the environment once NO_3^- has become depleted.

A wide range of maximum uptake rates and half saturation constants for NO_3^- , NH_4^+ , urea, and amino acid uptake have been observed among dinoflagellates both in laboratory cultures and experiments using natural populations during blooms (Table 4). Higher maximum uptake rates have been observed for NH_4^+ compared to NO_3^- , urea or amino acids for most species including: cultures of *P. minimum*, *Heterosigma akashiwo* and *Alexandrium minutum*, and assemblages dominated by *Dinophysis acuminata* (representing ~91% of the phytoplankton biomass), *P. minimum* (representing >93% of the total phytoplankton cells), *Alexandrium catenella* (more than 99% of the phytoplankton) and *Karenia mikimotoi* (cell density up to $8 \times 10^{-6} \text{ cell} \cdot \text{L}^{-1}$) (Table 4). In contrast, in cultures of the diatom *Pseudonitzschia australis*, maximum NO_3^- uptake rates exceeded NH_4^+ uptake rates in populations growing on NO_3^- -deplete condition [54]. The high variability in maximum uptake rates and half saturation constants between species and experiments could be due to differences in the preconditioning of cells prior to kinetic studies. Nutritional history was a significant factor affecting N uptake rates by *P. minimum* culture [55]. N uptake kinetics are known to vary with in response to the physiological status and their nutrient history of cells [36]. Our culture experiments demonstrated that *P. donghaiense* can take up NH_4^+ and other N compounds at high rates even when ambient NO_3^- concentrations are high (e.g., $\sim 43 \mu\text{mol N L}^{-1}$). This nutritional flexibility may contribute to the initiation and long duration of *P. donghaiense* blooms in nature. While blooms initiate when NO_3^- concentrations are high, NO_3^- is rapidly depleted as blooms progress.

Studies have shown that nutrient preconditioning and physiological status [36], cell size [56,57], growth rates [50,55,58], incubation time (minutes versus hours) [51], N substrate interactions [58–61], N preconditioning [53,62], the DIN/DIP ratio [63], and environmental factors such as irradiance and temperature

[59,64,65] can all contribute to variations in nutrient uptake kinetics. Because we conducted short incubations of uniform duration in culture systems acclimated to identical conditions, it is unlikely that these factors contributed to variability in uptake rates observed in this study.

While there are still few studies examining N uptake during blooms of *P. donghaiense*, kinetic parameters for NO_3^- , NH_4^+ , urea, and glycine uptake were compared during successive dinoflagellate blooms in Changjiang River estuary and East China Sea coastal waters in 2005 [16]. In these mixed blooms, *Karenia mikimotoi* was the dominant species initially and then was succeeded by *P. donghaiense*. In most cases, when the bloom was dominated by *P. donghaiense*, the K_s values for urea and glycine uptake (K_s , 5.3 and $1.8 \mu\text{mol N L}^{-1}$ respectively) were higher than when the bloom was dominated by *K. mikimotoi* (K_s , 1.4 and $1.7 \mu\text{mol N L}^{-1}$ respectively) (Table 4). The differences in the K_s of the two bloom species for NO_3^- and NH_4^+ may have been a driver of species succession as concentrations of these compounds were drawn down during the *K. mikimotoi* bloom which preceded the *P. donghaiense* bloom.

Cell-normalized N uptake rates by *P. donghaiense*-dominated assemblages in the Changjiang River estuary were comparable to cell-normalized N uptake rates measured in the culture studies reported here. Uptake rates of NH_4^+ , urea, and glycine were ~ 96 , ~ 22 and $\sim 38 \text{ fmol N cell}^{-1} \text{ h}^{-1}$, respectively, during blooms of *P. donghaiense* (cell density and Chl *a* were $\sim 5.0 \times 10^6 \text{ cells L}^{-1}$ and $8.66\text{--}9.68 \mu\text{g L}^{-1}$, respectively) [11,16] while maximum uptake rates for NH_4^+ , urea and algal amino acids observed in this study were ~ 119 , 78 and $\sim 53 \text{ fmol N cell}^{-1} \text{ h}^{-1}$, respectively. This suggests that uptake may not have been saturated in the environment. The K_s values of NO_3^- and urea for *P. donghaiense* were much lower than the environmental concentrations, but except for NH_4^+ and DFAA, which suggest that *P. donghaiense* have higher affinities on NO_3^- and urea than those for NH_4^+ and DFAA, but the later two also contribute on *P. donghaiense* bloom initiation and duration.

Summary

Results presented here demonstrate that *P. donghaiense* can grow on a diverse array of N compounds as their sole source of N, including inorganic N (NO_3^- and NH_4^+) and organic N compounds such as urea, dissolved free amino acids, small peptides, and even cyanate to support their growth. Maximum specific growth rates varied by a factor of 2 for all of the N compounds tested. In addition, uptake kinetics for regenerated N sources (e.g., NH_4^+ , urea, and amino acids) were similar under NO_3^- -replete and -deplete conditions suggesting that competition for these compounds may contribute to the success of *P. donghaiense* during bloom initiation when NO_3^- concentrations are high, and maintenance, when NO_3^- concentrations have been depleted. The nutritional flexibility exhibited by *P. donghaiense* likely contributes to its success in eutrophic environments where inorganic nutrient concentrations can be high, but where nutrient concentrations and the dominant form of bioavailable N rapidly change in response to stochastic events and the formation of algal blooms.

Acknowledgments

We are pleased to acknowledge Peter W. Bernhardt for contributing isotopic analysis and Drs. Katherine C. Filippino and Yingzhong Tang for their kind assistance in reviewing earlier versions of this manuscript.

Author Contributions

Conceived and designed the experiments: ZH MM NX SD. Performed the experiments: ZH. Analyzed the data: ZH MM. Contributed reagents/materials/analysis tools: ZH NX MM SD. Wrote the paper: ZH MM.

References

- Anderson DM (1997) Turning back the harmful red tide. *Nature* 388: 513–514.
- Wang J, Wu J (2009) Occurrence and potential risks of harmful algal blooms in the East China Sea. *Science of the Total Environment* 407: 4012–4021.
- Anderson DM, Cembella AD, Hallegraeff GM (2012) Progress in understanding harmful algal blooms: paradigm shifts and new technologies for research, monitoring, and management. *Annual Review of Marine Science* 4: 143–176.
- Glibert PM, Burkholder JM, Kana TM (2012) Recent insights about relationships between nutrient availability, forms, and stoichiometry, and the distribution, ecophysiology, and food web effects of pelagic and benthic *Prorocentrum* species. *Harmful Algae* 14: 231–259.
- Heisler J, Glibert PM, Burkholder JM, Anderson DM, Cochlan WP, et al. (2008) Eutrophication and harmful algal blooms: A scientific consensus. *Harmful Algae* 8: 3–13.
- Yan T, Zhou M, Zou J (2002) A national report of HABs in China. In: “Max” Taylor FJR, Trainer VL, editors. *PICES Scientific Report No. 23, Harmful algal blooms in the PICES region of the North Pacific*. pp. 21–33.
- Lu S, Hodgkiss IJ (2004) Harmful algal bloom causative collected from Hong Kong waters. *Hydrobiologia* 512(1–3): 231–238.
- Zhou M (2010) Environmental Settings and Harmful Algal Blooms in the Sea Area Adjacent to the Changjiang River Estuary. *Coastal Environmental and Ecosystem Issues of the East China Sea*, pp. 133–149 (Citable URL: <http://hdl.handle.net/10069/23506>).
- Lu D, Goebel J (2001) Five red tide species in genus *Prorocentrum* including the description of *Prorocentrum donghaiense* Lu Sp. Nov. from the East China Sea. *Chinese Journal of Oceanography and Limnology* 19(4): 337–344.
- Lu D, Goebel J, Qi Y, Zou J, Han X, et al. (2005) Morphological and genetic study of *Prorocentrum donghaiense* Lu from the East China Sea, and comparison with some related *Prorocentrum* species. *Harmful Algae* 4(3): 493–505.
- Li J, Glibert PM, Zhou M, Lu S, Lu D (2009) Relationships between nitrogen and phosphorus forms and ratios and the development of dinoflagellate blooms in the East China Sea. *Marine Ecology Progress Series* 383: 11–26.
- Paerl HW (1997) Coastal eutrophication and harmful algal blooms: importance of anthropogenic deposition and groundwater as ‘new’ nitrogen and other nitrogen sources. *Limnology and Oceanography* 42: 1154–1165.
- Anderson DM, Glibert PM, Burkholder JM (2002) Harmful algal blooms and eutrophication: nutrient sources, composition, and consequences. *Estuaries* 25: 704–726.
- Glibert PM, Burkholder JM (2006) The complex relationships between increasing fertilization of the earth, coastal eutrophication and proliferation of harmful algal blooms. In: Granéli E, Turner J, editors. *Ecology of harmful algae*. Springer-Verlag, Berlin. pp. 341–354.
- Zhou M, Shen Z, Yu R (2008) Responses of a coastal phytoplankton community to increased nutrient input from the Changjiang (Yangtze) River. *Continental Shelf Research* 28: 1483–1489.
- Li J, Glibert PM, Zhou M (2010) Temporal and spatial variability in nitrogen uptake kinetics during harmful dinoflagellate blooms in the East China Sea. *Harmful Algae* 9: 531–539.
- Shen Z, Liu Q, Zhang S, Miao H, Zhang P (2003) A nitrogen budget of the Changjiang River catchment. *Ambio* 32(1): 65–69.
- Chai C, Yu Z, Song X, Cao X (2006) The status and characteristics of eutrophication in the Yangtze River (Changjiang) Estuary and the adjacent East China Sea, China. *Hydrobiologia* 563: 313–328.
- Hu Z, Mulholland MR, Duan S, Xu N (2012) Effects of nitrogen supply and its composition on the growth of *Prorocentrum donghaiense*. *Harmful Algae* 13: 72–82.
- Antia NJ, Harrison PJ, Oliveira L (1991) Phycological reviews: the role of dissolved organic nitrogen in phytoplankton nutrition, cell biology, and ecology. *Phycologia* 30: 1–89.
- Berman T, Bronk DA (2003) Dissolved organic nitrogen: a dynamic participant in aquatic ecosystems. *Aquatic Microbial Ecology* 31: 273–305.
- Bronk DA, Steinberg D (2008) Nitrogen regeneration. In: Capone DG, Bronk DA, Mulholland MR, Carpenter EJ, editors. *Nitrogen in the marine environment*. Academic Press. pp. 385–468.
- Fu M, Wang Z, Li Y, Li R, Sun P, et al. (2009) Phytoplankton biomass size structure and its regulation in the Southern Yellow Sea (China): seasonal variability. *Continental Shelf Research* 29: 2178–2194.
- Yuan X, Glibert PM, Xu J, Liu H, Chen M, et al. (2012) Inorganic and organic nitrogen uptake by phytoplankton and bacteria in Hong Kong waters. *Estuaries and Coasts* 35: 325–334.
- Guillard RRL (1975) Culture of phytoplankton for feeding marine invertebrates. In: Smith WL, Chanley MH, editors. *Culture of Marine Invertebrate Animals*. New York: Plenum Press. pp. 26–60.
- Parsons TR, Maita Y, Lalli CM (1984) A manual of chemical and biological methods for seawater analysis. Pergamon Press. pp. 173.
- Solorzano L (1969) Determination of ammonia in natural waters by the phenylhypochlorite method. *Limnology and Oceanography* 14: 799–801.
- Cowie GL, Hedges JI (1992) Improved amino acid quantification in environmental samples: Charge-matched recovery standards and reduced analysis time. *Marine chemistry* 37: 223–238.
- Welschmeyer NA (1994) Fluorometric analysis of chlorophyll *a* in the presence of chlorophyll *b* and pheopigments. *Limnology and Oceanography* 39: 1985–1992.
- Guillard RRL (1973) Division rates. In: Stein JR, editor. *Handbook of phycolgical methods: culture methods and growth measurements*. Cambridge: Cambridge University Press. pp. 289–311.
- Monod J (1942) *Recherches sur la croissance des cultures Bactériennes*. 2nd ed. Hermann, Paris. pp. 211.
- Mulholland MR, Lee C (2009) Peptide hydrolysis and the uptake of dipeptides by phytoplankton. *Limnology and Oceanography* 54(3): 856–868.
- Anderson DM, Burkholder JM, Cochlan WP, Glibert PM, Gobler CJ, et al. (2008) Harmful algal blooms and eutrophication: Examining linkages from selected coastal regions of the United States. *Harmful Algae* 8: 39–53.
- Kana TM, Lomas MW, MacIntyre HL, Cornwell JC, Gobler CJ (2004) Stimulation of the brown tide organism, *Aureococcus anophagefferens*, by selective nutrient additions to in situ mesocosms. *Harmful Algae* 3: 377–388.
- Dugdale RC, Wilkerson FP, Hogue VE, Marchi A (2007) The role of ammonium and nitrate in spring bloom development in San Francisco Bay. *Estuarine, Coastal and Shelf Science* 73: 17–29.
- Mulholland MR, Lomas MW (2008) N uptake and assimilation. In: Capone DG, Bronk DA, Mulholland MR, Carpenter EJ, editors. *Nitrogen in the marine environment*. Academic Press. pp. 303–384.
- Mulholland MR, Gobler CJ, Lee C (2002) Peptide hydrolysis, amino acid oxidation and nitrogen uptake in communities seasonally dominated by *Aureococcus anophagefferens*. *Limnology and Oceanography* 47: 1094–1108.
- Yu J, Tang D, Oh Im-Sang, Yao L (2007) Response of harmful algal blooms to environmental changes in Daya Bay, China. *Terrestrial Atmospheric and Oceanic Sciences* 18(5): 1011–1027.
- Seeyave S, Probyn TA, Pitcher GC, Lucas MI, Purdie DA (2009) Nitrogen nutrition in assemblages dominated by *Pseudo-nitzschia* spp., *Alexandrium catenella* and *Dinophysis acuminata* off the west coast of South Africa. *Marine Ecology Progress Series* 379: 91–107.
- Jeong HJ, Yoo YD, Kim JS, Seong KA, Kang NS, et al. (2010) Growth, feeding, and ecological roles of the mixotrophic and heterotrophic dinoflagellates in marine planktonic food webs. *Ocean science journal* 45 (2): 65–91.
- Monod J (1950) La technique de la culture continue, théorie et applications. *Annales de l’Institut Pasteur Paris* 79(4): 390–410.
- Guillard RRL, Kilham P, Jackson TA (1973) Kinetics of silicon-limited growth in the marine diatom *Thalassiosira pseudonana* Hasle and Heimdal (= *Cyclotella nana* Hustedt). *Journal of Phycology* 9(3): 233–237.
- Taylor GT, Gobler CJ, Sañudo-Wilhelmy SA (2006) Speciation and concentrations of dissolved nitrogen as determinants of brown tide *Aureococcus anophagefferens* bloom initiation. *Marine Ecology Progress Series* 312: 67–83.
- Gobler CJ, Burson A, Koch F, Tang Y, Mulholland MR (2012) The role of nitrogenous nutrients in the occurrence of harmful algal blooms caused by *Cochlodinium polykrikoides* in New York estuaries (USA). *Harmful Algae* 17: 64–74.
- Healey FP (1980) Slope of the Monod equation as an indicator of advantage in nutrient competition. *Microbial Ecology* 5(4): 281–286.
- Tilman D (1977) Resource competition between planktonic algae: an experimental and theoretical approach. *Ecology* 58: 338–348.
- Hu Z, Xu N, Duan S, Li A, Zhang C (2010) Effects of urea on the growth of *Phaeocystis globosa*, *Scirrhopsiella trochoidea*, *Skeletonema costatum*. *Acta Scientiae Circumstantiae* 30(6): 1265–1270 (In Chinese, with English abstract).
- Allen CM, Jones ME (1964) Decomposition of carbamoyl phosphate in aqueous solution. *Biochemistry* 3: 1238–1247.
- Eppley RW, Thomas WH (1969) Comparison of half-saturation constants for growth and nitrate uptake of marine phytoplankton. *Journal of Phycology* 5: 375–379.
- Goldman JC, Glibert PM (1983) Kinetics of inorganic nitrogen uptake by phytoplankton. In: Carpenter EJ, Capone DG, editors. *Nitrogen in the marine environment*. Academic Press, New York. pp. 233–274.
- Harrison PJ, Parslow JS, Conway HL (1989) Determination of nutrient uptake kinetic parameters: a comparison of methods. *Marine Ecology Progress Series* 52: 301–312.
- Herndon J, Cochlan WP (2007) Nitrogen utilization by the raphidophyte *Heterosigma akashiwo*: Growth and uptake kinetics in laboratory cultures. *Harmful Algae* 6: 260–270.
- Cochlan WP, Harrison PJ (1991) Uptake of nitrate ammonium and urea by nitrogen-starved cultures of *Micromonas pusilla* (Prasinophyceae): transient responses. *Journal of Phycology* 27: 673–679.
- Cochlan WP, Herndon J, Kudela RM (2008) Inorganic and organic nitrogen uptake by the toxic diatom *Pseudo-nitzschia australis* (Bacillariophyceae). *Harmful Algae* 8: 111–118.

55. Fan C, Glibert PM, Burkholder JM (2003) Characterization of the affinity for nitrogen, uptake kinetics, and environmental relationships for *Prorocentrum minimum* natural blooms and laboratory cultures. *Harmful Algae* 2: 283–299.
56. Smith REH, Kalf J (1982) Size dependent phosphorus uptake kinetics and cell quota in phytoplankton. *Journal of Phycology* 18: 275–284.
57. Hein M, Pedersen MF, Sand-Jensen K (1995) Size-dependent nitrogen uptake in micro- and macroalgae. *Marine Ecology Progress Series* 118: 247–253.
58. Maguer J-F, L'Helguen S, Madec C, Labry C, Le Corre P (2007) Nitrogen uptake and assimilation kinetics in *Alexandrium minutum* (Dinophyceae): effects of N-limited growth rate on nitrate and ammonium interactions. *Journal of Phycology* 43: 295–303.
59. Lomas MW, Glibert PM (1999) Interactions between NH_4^+ and NO_3^- uptake and assimilation: comparison of diatoms and dinoflagellates at several growth temperatures. *Marine Biology* 133: 541–551.
60. Cochlan WP, Bronk DA (2003) Effects of ammonium on nitrate utilization in the Ross Sea: implications for f-ratio estimates. In: DiTullio GR, Dunbar RB, editors. *Biogeochemistry of the Ross Sea*. AGU Antarctic Research Series 78. pp. 159–178.
61. Jauzein C, Loureiro S, Garcés E, Collos Y (2008) Interactions between ammonium and urea uptake by five strains of *Alexandrium catenella* (Dinophyceae) in culture. *Aquatic Microbial Ecology* 53: 271–280.
62. Lindehoff E, Granéli E, Glibert PM (2011) Nitrogen uptake kinetics of *Prymnesium parvum* (Haptophyte). *Harmful Algae* 12: 70–76.
63. Li J, Glibert PM, Alexander JA (2011) Effects of ambient DIN:DIP ratio on the nitrogen uptake of harmful dinoflagellate *Prorocentrum minimum* and *Prorocentrum donghaiense* in turbidistat. *Chinese Journal of Oceanology and Limnology* 29(4): 746–761.
64. Cochlan WP, Price NM, Harrison PJ (1991) Effects of irradiance on nitrogen uptake by phytoplankton: comparison of frontal and stratified communities. *Marine Ecology Progress Series* 69: 103–116.
65. Kudela RM, Cochlan WP (2000) Nitrogen and carbon uptake kinetics and the influence of irradiance for a red tide bloom off southern California. *Aquatic Microbial Ecology* 21: 31–47.
66. Collos Y, Gagne C, Laabir M, Vaquer A, Cecchi P, et al. (2004) Nitrogenous nutrition of *Alexandrium catenella* (Dinophyceae) in cultures and in Thau Lagoon, southern France. *Journal of Phycology* 40: 96–103.
67. Kudela R, Ryan J, Blakely M, Lane J, Peterson T (2008) Linking the physiology and ecology of *Cochlodinium* to better understand harmful algal bloom events: a comparative approach. *Harmful Algae* 7: 278–292.