

A Habitat Model for the Detection of Two-lined Salamanders at C. F. Phelps Wildlife Management Area, Fauquier and Culpeper Counties, Virginia

Jay D. McGhee,¹ Randolph-Macon College,
Ashland, Virginia, 23005 and

Michael D. Killian, Department of Biological Sciences,
University of Mary Washington, Fredericksburg, Virginia 22401

ABSTRACT

Aquatic salamanders represent an important component of Virginia river watersheds, but despite potential declines, few specifics are known about their habitat preferences. We surveyed the habitats of the northern two-lined salamander and collected data on an array of habitat variables associated with the species. We used a logistic regression analysis to develop a model predicting its presence or absence for a given 50m-transect. Our final model incorporated the variation in stream depth and direction of stream flow and accounted for 25% of the variation in our data. We conclude that stream depth variation is an important feature of salamander habitat ecology, and surmise that direction of flow is of site-specific importance possibly related to stream order. Both features may be behavioral adaptations to avoid fish predation.

INTRODUCTION

Stream-dwelling salamanders are an important component of aquatic ecosystems. They account for a significant proportion of the biomass of a stream ecosystem, and act as a key trophic link, important as both predators and prey (Spight 1967, Burton and Likens 1975, Rocco and Brooks 2000). Consequently, these salamanders have potential to act as an indicator of stream health (Rocco and Brooks 2000, Barr and Babbitt 2002). This is particularly true for headwater streams where salamanders may act as the dominant vertebrate predator (Davic and Welsh 2004). Accordingly, it would be beneficial to better understand how these species make use of their available habitat. This is especially important in the face of on-going amphibian declines (Alford and Richards 1999). Knowledge of this type may provide better insights into the conservation of these species and their associated ecosystems (Cushman 2005).

Previous surveys of stream and terrestrial amphibian diversity have been carried out in the Rappahannock River watershed of northern Virginia; however, more needs to be done to quantify the habitat preferences of important stream species (Mitchell 1998, McGhee and Killian 2010). To begin addressing this need, we conducted a preliminary study of salamander habitat at C.F. Phelps Wildlife Management Area (WMA) located in the Rappahannock River watershed and developed a simple habitat model for the

¹ Corresponding author. E-mail: jaymcghee@rmc.edu

northern two-lined salamander (*Eurycea bislineata*), a common stream species for the area (McGhee and Killian 2010).

Northern two-lined salamanders are common to northern Virginia forest streams within the Rappahannock River watershed (Mitchell and Reay 1999). While they are considered potentially important components of the local ecosystems in which they occur, few studies have developed predictive models of habitat use (Davic and Welsh 2004). They occupy stream margins and seeps, using submerged rocks and woody debris for cover; but may periodically be found in upland terrestrial sites (Petranka 1998). Females attach eggs beneath submerged rocks of varying surface area in headwater streams (Jakubanis et al. 2008). Larvae of this species are benthic predators associated with stream pools with low silt (Smith and Grossman 2003, Petranka 1998). Two-lined salamanders are able to access low-order streams typically inaccessible to predatory fishes, and have become adapted to these headwater stream environments (Vannote et al. 1980, Davic and Welsh 2004). We hypothesized that two-lined salamanders would be detected in or near cool narrow, shallow streams. From this hypothesis, we predicted that important habitat variables in a logistic regression model would be stream temperature, stream depth, and stream width.

METHODS

We chose sampling sites by randomly selecting a GPS starting location constrained to occur within C. F. Phelps WMA, and moving from that point to the nearest stream. We then moved upstream or downstream a randomly selected distance of up to 50m, and laid a 50m transect running downstream. We sampled stream transects by searching five 1-m² quadrats placed within each of the five 10-m sections of the transect. The particular location of the quadrat within these 10-m sections was randomly selected (Jaeger 1994, Jaeger and Inger 1994). We searched quadrats by looking under larger cover objects such as rocks or decaying logs, leaf pack, leaf litter, and using a standard-mesh aquarium dip net (1/16 inch mesh size) to sample stream bottoms (Mitchell 2000). We identified captured salamanders to species (Petranka 1998). Data were collected at both transect and quadrat levels (Table 1).

We used logistic regression to select models with those predictive variables most associated with salamander captures at the transect level. Variables measured at the quadrat level were averaged and averages and standard deviations were used as separate predictor variables. As synergistic effects may occur between the variables we measured, we created *a priori* multiplicative variables for testing as well (Table 1). We used forward stepwise selection ($P = 0.05$ to enter and 0.10 to remove) in SPSS (SPSS Inc., Chicago IL). We assessed variable coefficients using the change in -2 loglikelihood and evaluated the explanatory value of models using Nagelkerke's r^2 (Ryan 1997, Hosmer and Lemeshow 1989, Nagelkerke 1991). For all statistical analyses $\alpha = 0.05$.

RESULTS

From 13 April 2007 – 21 April 2009, we sampled 78 stream transects with 390 stream quadrats. We located 256 two-lined salamanders, 203 of which were larval. Two-lined salamanders were detected in 45 of the 78 stream transects, for a 58% encounter rate. Logistic regression selected two predictor variables: the standard deviation of maximum stream depth (SDMD: -0.12 ± 0.06 SE, change in -2 log

Table 1. Habitat variables for stream and terrestrial transect sites at C. F. Phelps Wildlife Management Area, Virginia. For variables that had a standard deviation (SD) associated with them, the SD was included in the analysis as a separate predictor.

Transect-Level	Quadrat-Level
Season ^a	Mean Maximum Depth
Relative Humidity	Maximum Depth SD
Vapor Pressure Deficit	Mean Stream Width
Air Temperature (C)	Stream Width SD
Air Pressure	Mean Depth*Width
Weather ^b	Depth*Width SD
Bank Habitat ^c	Mean Water Temperature
Direction of Stream Flow	Water Temperature SD
Slope of Stream Flow	

^a Spring: Mar 20/21, summer: June 20/21, fall: Sep 22/23, winter: Dec 21/22

^b Clear, partly cloudy, overcast, light rain, medium rain

^c Deciduous, coniferous, mixed deciduous/coniferous, open field/shrub

likelihood = 5.331, df = 1, $P = 0.021$), and direction of stream flow (Direction: 0.10 ± 0.01 SE, change in $-2 \log$ likelihood = 4.301, df = 1, $P = 0.038$, Figure 1). The model explained 25% of the variation in data ($r^2 = 0.25$). Probability of predicting the detection of a two-lined salamander within a stream transect was equal to

$$\frac{1}{1 + e^{-(0.10 \text{ Angle} - 0.12 \text{ SD max depth} - 0.753)}}$$

This model would correctly predict the presence of two-lined salamanders in 84% of cases in our study site, and correctly predict the absence in 48% of cases. The standard deviation and the average of the maximum stream depth were positively correlated ($r = 0.75$, $P << 0.0001$), and so the majority of transects with low variability in depth also tended to be shallow. Two-lined salamanders tended to be found in streams flowing both south and west (logistic regression $\beta = 0.10$, $P = 0.05$). No other variables or combinations thereof produced models of significant predictive value.

DISCUSSION

Our model indicated that two-lined salamanders are sensitive to variation in stream depth. As those streams with high depth variation tended to be generally deeper, we interpret this as a preference for shallower sites in avoidance of fish predators (Sih et al. 1992). The majority of our captures were larval, and Barr and Babbitt (2002) found that larval two-lined salamanders occurred in negative association with brook trout (*Salvelinus fontinalis*), a fish predator. Average maximum depth also tended to be chosen by models if depth SD and direction of stream flow were removed, reinforcing the likely importance of depth. Variation in depth may provide refuges for predators to feed on larvae, or larvae and adults may simply tend to avoid deeper sites. No salamanders were found in our study site at depths greater than 20 cm.




FIGURE 1. The relationship between angle of stream flow and stream depth variation for transects at C. F. Phelps Wildlife Management Area, Virginia. Salamanders were typically detected (circles) in streams with relatively low variability, flowing southwest.

The model's selection of stream flow direction as a predictor of the presence of two-lined salamanders is difficult to interpret. Individuals were most easily detected in streams flowing towards the south and west, towards the general direction of the bordering Rappahannock River. South and west flowing streams tended to flow either close to the Rappahannock or to be a 2nd or 3rd order stream, and larvae, which often drift downstream, may be attempting to find slow moving, shallow, or low depth-variance pools with sufficient cover (Petranka 1998, Barr and Babbitt 2002). Bruce (1986) found that first-year two-lined larvae tended to dominate downstream samples compared to upstream samples. Unfortunately, direction of stream flow is unlikely to translate this effect to other sites very well.

Interestingly, the model failed to include stream temperature. Grant et al. (2005) also failed to detect a water temperature effect for two-lined salamanders in the Shenandoah National Park, Virginia. Barr and Babbitt (2002) and Rocco and Brooks (2000), however, detected a positive relationship between two-lined salamander presence and temperature, but they may have found a greater range of temperatures concurrent with the greater elevation variability at their sites (300 – 1200 m and 358 – 752 m compared to our 200 – 400 m).

Our model was able to provide significant information on the habitat used by two-lined salamanders using only two relatively easily acquired variables, and recommends itself for use as a preliminary predictor for presence/absence surveys when relatively few man-hours are available. It does tend to discount sites where the species does occur (false absences) about half the time, however, so more complete models are required to better understand the habitat ecology of the species.

ACKNOWLEDGMENTS

We thank Joe Ferdinandsen, manager of C. F. Phelps WMA for his cooperation, along with R. Hughes & Jerry Sims. We thank University of Mary Washington students Carly Byers, Sarah Almahdali, Jennifer Clary, Hillary Adams, and Ramsey Hanna for their assistance in the field. We thank the students of J. D. McGhee's animal ecology classes for Fall 2007 and 2008 for their help in field sampling. This work was made possible through a grant by the University of Mary Washington.

LITERATURE CITED

- Alford, R. A., and S. J. Richards. 1999. Global amphibian declines: a problem in applied Ecology. *Annual Review of Ecology and Systematics* 30:133–165.
- Barr, G. E. and K. J. Babbit. 2002. Effects of biotic and abiotic factors on the distribution and abundance of larval two-lined salamanders (*Eurycea bislineata*) across spatial scales. *Oecologia* 133:176–185.
- Bruce, R. C. 1986. Upstream and downstream movements of *Eurycea bislineata* and other salamanders in a southern Appalachian stream. *Herpetologica* 42:149–155.
- Burton, T. M. and G. E. Likens. 1975. Energy flow and nutrient cycling in salamander populations in the Hubbard Brook Experimental Forest, New Hampshire. *Ecology* 56: 1068 – 1080.
- Cushman, S.A. 2005. Effects of habitat loss and fragmentation on amphibians: A review and prospectus. *Biological Conservation* 128:231–240.
- Davic, R.D., and H.H. Welsh, Jr. 2004. On the ecological roles of salamanders. *Annual Review of Ecology, Evolution, and Systematics* 35:405–434.
- Grant, E. H. C., R. E. Jung, and K. C. Rice. 2005. Stream salamander species richness and abundance in relation to environmental factors in Shenandoah National Park, Virginia. *American Midland Naturalist* 153: 348 – 356.
- Hosmer Jr., D. W. and S. Lemeshow. 1989. *Applied Logistic Regression*. John Wiley and Sons, New York.
- Jaeger, R. G. 1994. Transect sampling. Pages 103–106 *in* W. R. Heyer, M. A. Donnelly, R. W. McDiarmid, L. C. Hayek, and M. S. Foster, editors. *Measuring and monitoring biological diversity: standard methods for amphibians*. Smithsonian Institution Press, Washington, D. C.
- Jaeger, R. G. and R. F. Inger. 1994. Quadrat sampling. Pages 97–102 *in* W. R. Heyer, M. A. Donnelly, R. W. McDiarmid, L. C. Hayek, and M. S. Foster, editors. *Measuring and monitoring biological diversity: standard methods for amphibians*. Smithsonian Institution Press, Washington, D. C.
- Jakubanis, J., M. J. Dreslik, and C. A. Phillips. 2008. Nest ecology of the southern two-lined salamander (*Eurycea cirrigera*) in eastern Illinois. *Northeastern Naturalist* 15:131–140.
- McGhee, J.D. and M.D. Killian. 2010. Salamander diversity at C. F. Phelps wildlife management area, Fauquier and Culpeper counties, Virginia. *Northeastern Naturalist* 17:629 – 638.
- Mitchell, J. C. 1998. Amphibian decline in the Mid-Atlantic Region: monitoring and management of a sensitive resource. Final Report, Legacy Resource Management Program, U. S. Department of Defense, Arlington, Virginia.
- Mitchell, J. C. 2000. *Amphibian monitoring methods & field guide*. Smithsonian

- National Zoological Park Conservation and Research Center, Front Royal, Virginia.
- Mitchell, J.C., and K.K. Reay. 1999. Atlas of Amphibians and Reptiles in Virginia. Special Publication Number 1, Virginia Department of Game and Inland Fisheries, Richmond, Virginia.
- Nagelkerke, N. J. D. 1991. A note on the general definition of the coefficient of determination. *Biometrika*, 78:691-692.
- Petranka, J. W. 1998. Salamanders of the United States and Canada. Smithsonian Institution Press, Washington, D. C.
- Rocco, G. L. and R. P. Brooks. 2000. Abundance and distribution of a stream Plethodontid salamander assemblage in 14 ecologically dissimilar watersheds in the Pennsylvania central Appalachians. Final Report No. 2000-4. Pennsylvania State Cooperative Wetlands Center, Forest Resources Laboratory, Pennsylvania State University. Prepared for U.S. Environmental Protection Agency, Region III.
- Ryan, T. P. 1997. Modern regression methods. John Wiley and Sons, Inc. New York.
- Sih, A., L. B. Kats, and R. D. Moore. 1992. Effects of predatory sunfish on the density, drift, and refuge use of stream salamander larvae. *Ecology* 73:1418-1430.
- Smith, S., and G.D. Grossman. 2003. Stream microhabitat use by larval Southern Two-lined Salamanders (*Eurycea cirrigera*) in the Georgia Piedmont. *Copeia* 2003:531-543.
- Spight, T. M. 1967. Population structure and biomass production by a stream salamander. *American Midland Naturalist* 78:437-447.
- Vannote, R.L., G. W. Minshall, K.W. Cummins, J.R. Sedell, and C.E. Cushing. 1980. The river continuum concept. *Canadian Journal of Fisheries and Aquatic Sciences* 37:130-137.